

Biochemical Indices in Ecology: A Catalyst in Environmental Study

Nwaichi EO^{1,2*} and Essien EB²

¹The University of Nottingham, Nottingham, UK

²University of Port Harcourt, Port Harcourt Nigeria, Nigeria

Abstract

Environmental challenges has continued to threaten the existing unsustainable global living. The debilitating impact on environment arises from employed techniques or effects of such techniques and requires sensitive and effective monitoring indicators for its evaluation. Borne out of increasing complexity of environmental problems and apparent strained resources of government, researchers, industries and decaying food production and water supply, a review of indices of interactions between organisms and the environmental factors that impact them in a bid to foster environmental protection and nature conservation which are core elements of ecology, is therefore pertinent.

Keywords: Biochemistry; Environmental quality; Microbial community; biological indicators of pollution; Biomarkers.

Commentary

Biochemistry, the study of chemical processes within and relating to living organisms [1] or chemistry of living matter, deals with the structures, functions and interactions of biological macromolecules. Biochemical indices traverses genetics and molecular biology with emerging novel roles for RNA [2]. Ecology is the branch of biology that deals with relationship among organisms and to their physical surroundings. It is a branch of science concerned with the interrelationship of organisms and their environments [3] or simply the totality or pattern of relations between organisms and their environment. Ecological diversity measurements [4] are necessary to understand survival and adaptability of species. The role of biochemistry in interactions between organisms or between an organism and its environment (soil, water and air) cannot be overestimated.

Soil biological processes, such as organic matter degradation, mineralization, and nutrient recycling, are basically driven by microorganisms and enzymes. Frac, et al. [5] reported that community level physiological profiling using Biolog EcoPlates can be used as sensitive and effective indicator for evaluation of microbial functional diversity and community of soil under sewage sludge-treatment. Also, the possibility of early detection of soil physical and chemical properties from modification in biological properties, including microbial community has been documented by Szarlip, et al. [6]. The ratio of total Nitrogen calculated from biochemical soil properties (Nc) (biomass C, mineralised N, phosphomonoesterase, β - glucosidase and urease activities) to that measured by Kjeldahl method (Nk) was used as an index of soil quality in disturbed soils; in all cases, soil degradation was reflected in Nc/Nk values [7]. The researchers recommended the adoption of the Nc: Nk for evaluation of soil degradation as it distinguishes among biochemically balanced, activity transient and degraded soils and its application in the rapid calculation of ecological dose, ED₅₀ of soil pollutants. Soil productivity has been linked to biochemical activity. The quality of soil is related to the maintenance of equilibrium between organic matter content and these five biochemical indices. Nwaichi and Frac [8] recorded correlation of the 5'-terminal restriction fragments generated after Csp digestion of 16S rRNA gene with observed DNA concentrations in the soil community profile and revealed loss of diversity with crude oil pollution. They posited that the herculean nature of the study of complex diversity pattern of the microbial community in parched and inundated soil environment may have arisen from inadequate

biological classification of such environment. Molecular methods have the advantage of obtaining information about uncultivable organisms and can be applied to study complex trophic interactions in the field and to address underlying ecological questions. A wholistic understanding of soil microbial communities has been broadly defined by biochemical-based techniques and molecular-based techniques [9]. Similarly, using Average well colour development in BIOLOG system with 95 different carbon sources used to produce a metabolic profile of microorganisms, Nwaichi and Frac [8] showed wide distribution in soil community spread.

According to Chícharo and Chícharo [10], the ratio of RNA to DNA (known as standardized RNA/DNA ratio) has been widely used as a biochemical index to assess potential survival and growth and in determining the ecophysiological condition of marine organisms. Besides becoming a particularly promising biochemical tool that not only reflects physiological condition, this index also allows for the estimation of instantaneous growth rates in a great variety of organisms [11-13]. Chícharo and Chícharo [10] also reported validation of standardized RNA/DNA ratio for study of marine small animals. Cytochrome P450, a biomarker often used in the studies assessing aquatic environment pollution has proved to be a very suitable method for biochemical and environmental monitoring of aquatic environment quality [14]. In Czech Republic, Marcela, et al. [15] employed liver enzymes of phase I and phase II of xenobiotic transformation, namely cytochrome P450, ethoxyresorufin-O-deethylase, glutathione-S-transferase and tripeptide glutathione to assess the contamination status of different rivers. They reported that these biomarkers are very suitable and useful tool for monitoring of aquatic pollution in the study Czech rivers. The relationship between levels of biochemical markers in fish liver and the ambient level of organic pollutants in fish muscle and bottom sediment were demonstrated in the said study.

The use of Ascorbic acid content [16], chlorophyll content [17],

***Corresponding author:** Nwaichi EO, The University of Nottingham, Nottingham, UK, Tel: 234(0)8056031989; E-mail: nodullm@yahoo.com

Received: November 29, 2015; **Accepted:** January 20, 2016; **Published:** January 23, 2016

Citation: Nwaichi EO, Essien EB (2016) Biochemical Indices in Ecology: A Catalyst in Environmental Study. *Biochem Anal Biochem* 5: 238. doi:10.4172/2161-1009.1000238

Copyright: © 2016 Nwaichi EO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

leaf extract pH [18] and relative water content [19] have been used to determine the sensitivity and tolerance of plants to air pollution. In their study, Tanee, et al. [20] concluded that air pollution tolerance indices of monitored plants were of the sensitive category and hence can be used to monitor air quality of Niger Delta. Based on their results, air pollution tolerance index, APTI obtained using the four leaf biochemical status can serve in predicting air quality condition of an area and these biochemical parameters are important in the study of plant-environment interactions as they can serve as bioindicators of air pollution. In the same vein, Krishnaveni and Magesh [21] investigated air pollution tolerance index using some biochemical indices and submitted that most of study the plants were sensitive to parameters and advanced the added merits of these indices to include simplicity, cost-effectiveness of process and equipment. The plant species which are more sensitive act as biological indicators of air pollution. Increased number of important biochemical parameters gave more accurate and reliable results in air quality and exposure study carried out by Sarika, et al. [22]. Furthermore, their results from calculation of APTI values of twelve different tree species revealed that all monitored biochemical parameters are important and greatly govern in evaluation of APTI values and thus, selection of suitable tree species to remove/reduce air pollution.

Biochemical systematics and ecology has accelerated understanding of environmental assessment, monitoring and management and should be further explored in related field. Environmental biota, processes, function, and the interactions has been to a great extent, revolutionized by biochemical indices. In a time of great global environmental crises, increasing complexity of environmental problems and obvious strained resources of the regulators and polluters, most reliable approach to studying ecology has become an increasingly important topic.

References

1. Biochemistry (2015) acs.org.
2. Ulveling D, Francastel C, Hubé F (2011) When one is better than two: RNA with dual functions. *Biochimie* 93: 633-644.
3. Ecology.
4. Sahney S, Benton MJ, Ferry PA (2010) Links between global taxonomic diversity, ecological diversity and the expansion of vertebrates on land. *Biol Lett* 6: 544-547.
5. Fraç M, Oszust K, Lipiec J, Jezierska-Tys S, Nwaichi EO (2014) Soil microbial functional and fungal diversity as influenced by municipal sewage sludge accumulation. *Int J Environ Res Public Health* 11: 8891-8908.
6. Szarlip P, Włodarczyk T, Brzezinska M, Glinski J (2010) Production and uptake of nitrous oxide (N₂O) as affected by soil conditions. *Acta Agrophysica* 187: 1-66.
7. Leiros MC, Trasar-Cepeda C, Garcia - Fernandez F, Gil - Sotres F (1999) Defining the validity of a biochemical index of soil quality. *Biol Fert Soils* 30: 140-146.
8. Nwaichi EO, Frac M (2015) Evaluation of soil microbial communities as influenced by crude oil pollution. *Afr J Biotechnol* 14: 1632-1639.
9. Abbasi FM, Ahmad H, Perveen F, Inamullah, Sajid M, et al. (2010) Assesment of genomic relationship between *Oryza sativa* and *Oryza australiensis*. *Afr. J. Biotechnol* 9: 1312-1316.
10. Chicharo MA, Chicharo L (2008) RNA:DNA ratio and other nucleic acid derived indices in marine ecology. *Int J Mol Sci* 9: 1453-1471.
11. Amaral V, Cabral HN, Paula J (2008) Implications of habitat specific growth and physiological condition for juvenile crab population structure. *Mar Freshwater Res* 59: 726-734.
12. KÄhler HR, Sandu C, Scheil V, Nagy-PetricÄf EM, Segner H, et al. (2007) Monitoring pollution in river MureÄY, Romania, Part III: biochemical effect markers in fish and integrative reflection. *Environ Monit Assess* 127: 47-54.
13. Buckley LJ, Caldarone EM, Clemmesen C (2008) Multi-species larval fish growth model based on temperature and fluorometrically derived RNA/DNA ratios: results from a meta-analysis. *Mar Ecol-Prog Ser* 371: 221-232.
14. Ikeda T, Sano F, Yamaguchi A, Matsuishi T (2007) RNA/DNA ratios of calanoid copepods from the epipelagic through abyssopelagic zones of the North Pacific Ocean. *Aquatic Biol* 1: 99-108.
15. Havelková M, Randák T, Blahová J, Slatinská I, Svobodová Z (2008) Biochemical markers for the assessment of aquatic environment contamination. *Interdiscip Toxicol* 1: 169-181.
16. Hoque MA, Banu MNA, Oluma E (2007) Exogenous proline and glycinebetaine increase NaCl-induced Ascorbate glythione cycle enzyme activities and praline improves salt tolerance more than glycinebetaine in tobacco bright yellow-2 suspension-cultural cells. *J. Plant Physiol* 164: 1457-1468.
17. Flowers MD, Fiscus EL, Burkey KO (2007) Photosynthesis, chlorophyll fluorescence and yield of snap bean (*Phaseolus vulgaris* L.) genotypes differing in sensitivity to Ozone. *Environmental and Experimental Botany* 61: 190-198.
18. Klumpp G, Furlan CM, Domingos M, Klumpp A (2000) Response of stress indicators and growth parameters of *Tibouchina pulchra* Cogn. exposed to air and soil pollution near the industrial complex of Cubatão, Brazil. *Sci Total Environ* 246: 79-91.
19. Rao DN (1979) Plant leaf as pollution monitoring device. *Fertilizer News* 24: 25-28.
20. Tanee FBG, Albert E, Amadi BR (2014) Biochemical Properties and Air Pollution Tolerance Indices of Plants in Port Harcourt City, Nigeria. *BJAST* 4: 4835-4845.
21. Krishnaveni M, Magesh P (2014) Air pollution tolerance index induced by biochemical components in plants. *Int J Pharm Pharm Sci* 6: 362-364.
22. Sarika K, Kiran P, Harshika K, Gauri R (2015) Air quality status and its effect on biochemical parameters of roadside Trees species in jalgaon city, Maharashtra. *IJPAES* 5: 128-133.

Citation: Nwaichi EO, Essien EB (2016) HBiochemical Indices in Ecology: A Catalyst in Environmental Study. *Biochem Anal Biochem* 5: 238. doi:10.4172/2161-1009.1000238

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>