

## Biological Activities and Valuable Compounds from Five Medicinal Plants

Million Getasetegn<sup>1\*</sup> and Yirefu Tefera<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Natural and Computational Sciences, Bule Hora University, Bule Hora, Ethiopia

<sup>2</sup>Department of Biology, Faculty of Natural and Computational Sciences, Bule Hora University, Bule Hora, Ethiopia

### Abstract

Currently, there are more than 85,000 plant species that have been documented for medical use globally. This implies, plant derived natural products hold great promise for discovery and development of new pharmaceuticals in diverse human ailments. Among these divers medicinal species *Aframomum corrorima*, *Commiphora myrrha*, *Nigella sativa*, *Ocimum lamiifolium* and *Olea europaea* are one of very well known valuable plants in serving the local people as spice, flavor, fumigants, and sources of traditional and modern medicines. In modern science their medicinal activities have been corroborated by phytochemical and biological research investigations. As the biological studies revealed they have antibacterial, antioxidant, antifungal, antimicrobial and anti-inflammatory properties even though further investigation required on *in vivo* and animal model to ascertain those *in vitro* activities are really valuable in life. The phytochemical investigations have also shown that these plants are holder of diversified useful compound classes like terpene, steroid, flavonoid, glycoside, fatty acid and alkaloids. However, their pharmacology and pharmacokinetics activities have to be further investigated in order to make sure their metabolic effect upon their usage because some compounds have ability to be metabolized during *in vivo* analysis, and to see whether these compounds have synergic or additive effects. Therefore, the purpose of this review paper is to present the compiled and unified scientific studies which have been done on the above well known medicinal plants about their description, medicinal uses, volatile and extracted active compounds, and chemical structures of isolated compounds in justification with scientific researches.

**Keywords:** *A. corrorima*; *C. myrrha*; *N. sativa*; *O. lamiifolium*; *O. europaea*

### Introduction

According to the WHO traditional medicine is a health practices, approaches, knowledge and beliefs incorporating plants, animals and mineral based medicine, spiritual therapies, manual techniques and exercises, applied singularly or in combination to treat, diagnose and prevent illnesses and maintain well being. The therapeutic use of natural products is the early medical practice. To date in the world, there are more than 85,000 plant species that have been documented for medical use. This indicates the plant derived natural products hold great promise for the discovery and development of new pharmaceuticals in a diverse human ailments [1]. This further implies many medicines widely in used today incorporate ingredients from plants. Almost 71% of new drugs that have been approved since 1981 have directly or indirectly been derived from natural products [2]. Hence, traditional medicinal plants have been greatly contributing in the development of modern medicines. The study revealed approximately 80% of the world's population relies on these traditional plant based medicines for primary health care. The remaining 20% of the world's population also depends on plant products for health care [3].

Ethnobotanical studies are one of the reliable guides for humans in utilizing the abundance of herbal resources that nature has to offer. Fabrican and Farnsworth reported on 122 compounds that obtained from 94 species of plants are being used worldwide in medicine [4]. In addition, 80% of these have had ethnomedical uses which are identical or related to the current use of active elements of plant. As a result, the authors have suggested that ethnomedicinal information must be preserved as a valuable source for novel drug discovery. Still there is a requisite for the development of scientific technology to enhance usage of medicinal plant products even though indigenous plants are known and commonly used as a source of medicines and spices in different countries. The following five indigenous plants *Aframomum corrorima*, *Commiphora myrrha*, *Nigella sativa*, *Ocimum lamiifolium*, and *Olea europaea* are among well known indigenous plants being as a source of medicine, spice, flavor and traditional healer. These plants are also holder of

diversified compound classes such as flavonoid, terpenoid, steroid and alkaloid.

Therefore, the main aim of this review paper is to present the compile and unified scientific studies which have been done on the above well known five indigenous plants about their description, medicinal uses, volatile and extracted active compounds, and chemical structures of some isolated compounds.

### *Aframomum corrorima* (Braun) Jansen

*Aframomum corrorima* (Braun) Jansen or Ethiopian cardamom is a renowned spice and medicinal crop of Zingiberaceae family. It is an Indigenous plant to Ethiopia and other countries. It is a close relative to the widely known Indian cardamom (*Elettaria cardamomum* Maton). The species is native to Ethiopia as well as some parts of Southwestern Sudan, Western Uganda, Tanzania and Eritrea [5-7]. *A. corrorima* locally in Ethiopia is known as "Korerima". The dried fruits are traded throughout Ethiopia and also into North East Africa, Arabia and India. The word 'Aframomum' is composed from Africa and Amomum, the latter derived from the Arabic "hamma" which means "hot, warm" indicating the pungency of the seeds. The specific epithet 'corrorima' refers in Latin form to the local name of the species, "Korerima" [7,8].

*A. corrorima* is a perennial aromatic herb growing about 1-2 m high with scaly of large size, bearing flowers either terminally on

\*Corresponding author: Million Getasetegn, Department of Chemistry, Faculty of Natural and Computational Sciences, Bule Hora University, Bule Hora, Ethiopia, Tel: 251913212460; E-mail: [noillim2001@gmail.com](mailto:noillim2001@gmail.com)

Received May 16, 2016; Accepted June 14, 2016; Published June 21, 2016

Citation: Getasetegn M, Tefera Y (2016) Biological Activities and Valuable Compounds from Five Medicinal Plants. Nat Prod Chem Res 4: 220. doi:10.4172/2329-6836.1000220

Copyright: © 2016 Getasetegn M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

aerial leaf shoots or from the ground level, red-brown underground rhizomes and leafy stems. It flowers usually from January to December and the fruits mature 2 or 3 months later. It is found in humid or moist forests, slightly shaded, wetter and open places in forests at 1350-2000 m altitude. Naturally, in Ethiopia it grows and cultivated in the forests of south and south western parts of the country such as Gamo Gofa, Debub Omo, Kaffa, Ilubabor, Sidamo, Welega and other regions. Even though farmers are currently collecting this spices plant from the forest and sale at the local market, the destruction of the natural forests for various purposes has significantly reduced the diversity of this shade obligate spices [7,9,10].

### Medicinal uses


Traditionally “*korerima*” has been used as spices, medicine, an income source and means of soil conservation [11]. It is also used as a carminative, purgative and tonic agent. The seeds are ground and mixed with other spices and used to flavor all kinds of sauces coffee, tea, butter, cottage cheese, bread and “*wot*” [9] and for preparation of “*Berberé*” (hot red pepper) [7].

Ethnobotanical surveys were conducted in the three major “*korerima*” growing regions of southern Ethiopia Gamo Gofa, Debub Omo and Kaffa. Plant parts are used as a medicine for treatment of different ailments. This study revealed 83% of key informants replied that seeds are mostly used as traditional medicine followed by leaves (75%) and rhizomes (72%) [11]. The antioxidant activity of seeds (29.59  $\mu\text{mol/g}$  dry weight (DW)) was significantly higher than that of pods (14.23  $\mu\text{mol/g}$  DW). The total phenolic content in seeds (3.98 mg/g DW) was significantly higher than in pods (1.32 mg/g DW). The phenolic substances including flavonoids and phenolic acids are responsible for the antioxidants activities. The seed of this plant has also antimicrobial and antifungal activities [10,12].

### Volatile compounds

Matured green capsules dried on wire mesh bed for 10 days scored maximum oleoresin (10.04% w/w) and essential oil content of dried seeds (5.53%) and essential oil from husks (0.93% v/w) [6]. The composition of essential oil has been investigated by GC and GC/MS. Eighteen components comprising 91% of the oil were identified and majority of them are sesquiterpenes. These are  $\beta$ -caryophyllene (2.11%) (1), nerolidol (13.89%) (2) and caryophyllene oxide (1.25%) (3) [13].

A research was done in Ethiopia to analyze essential oil composition by hydrodistillation. The oil refractive indices were recorded and shown to have 1.494 for leaf and 1.445 for rhizome. The GC and GC/MS analysis of these essential oils identified the presence of 38 (in leaf) and 52 (in rhizome) compounds. The major component of oil from leaf was  $\beta$ -caryophyllene (60.7%) (1). The rhizome oil was dominated by  $\gamma$ -terpinene (21.8%) (4) and  $\beta$ -pinene (17.6%) (5) [11] (Table 1). This essential oil has antifatulent, virustatic and motility-enhancing effects [7].



### Phytochemicals and elements

In the seed part, the levels of essential and non-essential metals were determined using flame atomic absorption spectrometry. As the research revealed the levels of the nutrients were in the following

ranges: Ca (1794-2181); Mg (1626-2067); Mn (141-180); Fe (37-46); Zn (12-18); Ni (6.6-8.5); Cu (5.8-8.3); Cr (3.8-5.8) and Co (2.0-2.3) in  $\mu\text{g/g}$ . The concentration of non-essential cadmium was in the range from 0.9-1  $\mu\text{g/g}$  while that of lead was below the detection limit of the method [14].

According to Tane et al. some of the *Aframomum* species are known for the production of labdane diterpenoids, flavonoids, sesquiterpenoids and arylalkaloids which may act against micro-organisms [15]. Solvent extraction yields about 10% oleoresin and this content depends on the solvent used and the raw material. Cardamom oleoresin contains about 52-58% volatile oil and used as flavoring agent [16].

### *Commiphora myrrha* (Nees) Engl.

Myrrh (*Commiphora myrrha* (Nees) Engl.) a yellow fragrant oleo-gum resin is an indigenous plant to Ethiopia and belongs to Burseraceae family. Its resin is collected commercially from South Ethiopia, Somalia, Sudan, Eritrea, Kenya and Arabia. The two species *C. molmol* and *C. myrrha* are synonymous. *C. myrrha* commonly is known in many countries as African myrrh, Somali myrrh, Common myrrh, Herabol myrrh, myrrh [17-19] and in Ethiopia commonly called “*Kerbe*”.

*C. myrrha* is a shrub or tree which growth up to 4 m high. The plant is mainly distributed in Eastern Africa, Arabia and India. It is found in Acacia-*Commiphora* woodland and bush land on sandy to loamy soil at altitudes 250-1300 m. In Ethiopia it is found in Afar, Bale, Hararge, Sidamo and Somalia. It is also found in north east Kenya and other parts of the world [20].

Myrrh is an aromatic gum resin of the plant stem resinous exudates of different *Commiphora* spp. that is used as medicinally active extract. Even though it has been collected from various species of the genus *Commiphora* such as *C. africana*, *C. erythraea*, *C. gileadensis*, *C. habessinica*, *C. hodai*, *C. kua*, *C. quadricincta*, *C. schimperi* and *C. truncate*, the best quality has been obtained from the specie of *C. myrrha* [21,22]. Myrrh consists of 7%-17% volatile oil, 25%-40% resin, 57%-61% gum, and 3%-4% impurities [22].

### Medicinal uses

Myrrh is used in traditional medicine particularly by women who fumigate their body for health care [22], as spices and flavor [17]. Ethnobotanical survey indicated that it has many medicinal activities and has been used to treat various diseases such as amenorrhea, ache, dysmenorrhhea, tumors, fever, stomach complaints, gall bladder, chest ailments, snake and scorpion bites [23]. The British Herbal Compendium allow the use of myrrh tincture as gargle to treat pharyngitis and tonsillitis, as mouth wash for gingivitis and ulcers, and externally to treat skin inflammations. In France its topical use is approved for treatment of small wounds, for nasal congestin from the common cold. The German Standard License for myrrh tincture indicates its use for inflammations of the gums and mouth mucosa because it contains large amounts of tannins [21,24].

Other than the ethnobotanical survey, the laboratory based researches have revealed myrrh is used as antibacterial, antiseptic, antitumor, antifungal and anaesthetic activity [25]. Antimicrobial activity of natural extracts and pure compounds can be detected by observing the growth response of various micro-organisms to samples that are placed in contact with them. The research findings suggested that extracts and isolated compounds of myrrh resin could be useful for preventing and treating human gynecologic cancer disease. *In vitro* research shown 85% EtOH and petroleum ether extract significantly

Plant name	Compounds name	Comd. No.	References
<i>Aframomum corrorima</i>	$\beta$ -Caryophyllene	1	[11,13]
	Nerolidol	2	
	Caryophyllene oxide	3	
	$\gamma$ -Terpinene	4	
	$\beta$ -Pinene	5	
<i>Commiphora myrrha</i>	Germacrene B	6	[20,21,28-30,32,34]
	Furanodiene	7	
	Lindistrene	8	
	Curzerene	9	
	$\beta$ -Elemene	10	
	Germacrene	11	
	Furanoedesma-1,3-diene	12	
	Curzerenone	13	
	Furanodiene-6-one	14	
	Neomyrrhaol	15	
	Sandaracopimaric acid	16	
	Abietic acid	17	
	2-Methoxy-5-acetoxyfuranogermacr-1(10)-en-6-one	18	
	Dehydroabietic acid	19	
	Myrrheterpenoid K	20	
	Myrrheterpenoid L	21	
	Myrrheterpenoid M	22	
	Myrrheterpenoid N	23	
	Furanogermacr-1(10)Z,4Z-dien-6-one	24	
	Commiferin	25	
	Quercetin	26	
	$\beta$ -Sitosterol	27	
	Commiterpenes A	28	
	Commiterpenes B	29	
	Commiterpenes C	30	
	D-Galactose	31	
	L-Arabinose	32	
	4-Methyl D-glucuronic acid	33	
	6-O-(4-O-Methyl- $\beta$ -D-glucuronosyl)-D-galactose	34	
	4-O-(4-O-Methyl- $\alpha$ -D-glucuronosyl)-D-galactose	35	
	2-Methoxy-8,12-epoxygermacra-1(10),7,11-triene-6-one	36	
	Docosanoic acid	37	
	Myrrhone	38	
Mansumbinone	39		
<i>Nigella sativa</i>	cis-4-Methoxythujane	40	[44,45,47,48,50,51]
	trans-4-Methoxythujane	41	
	Thymoquinone	42	
	Dithymoquinone	43	
	Thymol	44	
	Thymohydroquinone	45	
	Nigellicine	46	
	Nigellidine	47	
	Nigellimine	48	
	Nigellimine N-oxide	49	
	$\alpha$ -Hereditin	50	
	Quercetin	26	
	Kaempferol-3-glucosyl	51	
	Quercetin-3-(6-ferulolyl glucosyl)	52	
	Triglycoside quercetin 3-glucoside	53	
	Kaempferol 3-glucoside	54	
Rutin	55		

<i>Ocimum lamiifolium</i>	Bornyl acetate	56	[61,68-70]
	p-Cymene	57	
	α-Pinene	58	
	β-Pinene	5	
	Camphene	59	
	Sabinene	60	
	α-Phellandrene	61	
	Apigenin	62	
	Genkwanin	63	
	Acacetin	64	
	Apigenin 7,4'-dimethyl ether	65	
	Luteolin	66	
	Ladanein	67	
	5,6-Dihydroxy-7,3',4'-trimethoxyflavone	68	
	5,7-Dihydroxy-6,4'-dimethoxyflavone	69	
	Cirsimaritin	70	
	Salvigenin	71	
	Cirsiliol	72	
	Cirsilineol	73	
	Eupatorin	74	
5-Hydroxy-6,7,3',4'-tetramethoxyflavone	75		
Quercetin3-O-xylosyl(1"→2") galactoside	76		
Flavone 5-O-glycosides	77		
<i>Olea europaea</i>	α-Pinene	58	[73,77,79,80]
	2,6-Dimethyloctane	78	
	2-Methoxy-3-isopropylpyrazine	79	
	Hydroxytyrosol	80	
	Oleuropein	81	
	Verbascoside	82	
	Rutin	55	
	Luteolin-7-Glucoside	83	
	Nüzhenide	84	
	Oleuropein Glucoside	85	
	Maslinic acid	86	
	Oleanolic acid	87	
	Ursolic Acid	88	
	Erythrodil	89	
Uvaol	90		

**Table 1:** Lists of compounds that are exist in the five medicinal plants.

inhibited cell proliferation of human ovarian cancer and endometrial carcinoma cells. The inhibitory effects of these solvent extracts (85% EtOH and petroleum ether) on human ovarian cancer cells were strong and observed  $IC_{50}$  as 15.8 and 26.91  $\mu\text{g/mL}$ , respectively. Similarly, the  $IC_{50}$  values of these extracts on endometrial carcinoma cells were 20.73 and 26.63  $\mu\text{g/mL}$ , respectively [18]. Another biological activity that was investigated on MeOH extract of this plant has shown antifungal activity against *Aspergillus niger* and antibacterial activity at higher concentration. However, the petroleum ether extract has strongly shown antibacterial activity against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and antifungal activity to *Aspergillus niger* and *Candida albicans* [24,26]. Moreover, the resin powder extracts of *C. myrrha* were tested against different gram negative bacteria by disc diffusion method and it has been found that EtOAc and hexane extracts to possess bacterial growth inhibition [26]. In addition, the anti-inflammatory test of EtOH extract significantly inhibited the development of paw swelling induced by formalin and

significantly reduced acetic acid-induced writhing. Furthermore, in comparison of the pharmacological activities, the petroleum ether fraction was stronger than EtOH extract. This is because the petroleum ether fraction showed significant analgesic activity in the model at the dose of 100 mg/kg ( $p < 0.01$ ) and the ethyl acetate fraction exhibited less analgesic activity ( $p < 0.05$ ) [19].

### Volatile compounds


The essential oils of *Commiphora spp.* are rich in furanosesquiterpenoids, which have been found to possess anesthetic, antibacterial, antifungal, antihyperglycemic properties, asthma, bronchitis, for mature and wrinkled skin, gargle for mouth ulcers and thrush and cleanser for the womb, prevention of tooth decay, for treating cancer sores, sore throat, for treatment of mild inflammations of oral and pharyngeal mucosa [21,24].

The minimum inhibitory concentration (MIC) of the essential oil of *C. myrrha* against the toxigenic strain of *Aspergillus flavus* was shown at

3.0  $\mu\text{L/mL}$  concentration [27]. In Somalia the quenching activity against singlet oxygen, an actor of lipid peroxidation and DNA degradation, of the essential oil and resinoid of *C. myrrha* has been studied and compared to DL- $\alpha$ -tocopherol using 1,3-diphenylisobenzofuran (DPBF) as a probe. This research revealed the essential oil and menthofuran has a higher quenching activity than DL- $\alpha$ -tocopherol [25].

Volatile compounds from the oleo-gum resin of *C. myrrha*, which were collected from Ethiopia, were isolated by supercritical extraction method with  $\text{CO}_2$  and hydrodistillation. The main components are identified and quantified by GC/MS and has shown the following compounds germacrene B 4.3% (6), furanodiene 19.7% (7), lindestrene, 12.9% (8), curzerene, 8.5% (9),  $\beta$ -elemene 8.7% (10), germacrone, 5.8% (11), and furanoeudesma-1,3-diene 34.9% (12) [21,28,29] (Table 1). The irritant potentials of essential oil and seven sesquiterpenoid compounds from this oleo-gum-resin were investigated by open mouse ear assay. The essential oil curzerenone (13), furanodiene-6-one (14) and furanoeudesma-1,3-diene (12) showed potent and persistent irritant effects while others possess least irritant potentials [30].

Another research on the antioxidant and antimicrobial potential of MeOH and EtOAc crude extracts and the essential oil of *C. myrrha* resin were also investigated. As the research revealed the MeOH extract exhibited the highest antioxidant and antimicrobial activity as compared to EtOAc extract and essential oil [31].




## Phytochemicals

Investigation on the resin a new cycloartane type triterpene named cycloartane-1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ ,25-tetraol (neomyrrhaol) (15), along with four known terpenes, sandaracopimaric acid (16), abietic acid (17), 2-methoxy-5-acetoxyfuranogermacr-1(10)-en-6-one (18), and dehydroabietic acid (19) have been isolated. Compounds (18) and (19) exhibited significant aromatase inhibiting activity with  $\text{IC}_{50}$  values at 0.2  $\mu\text{M}$  and 0.3  $\mu\text{M}$ , respectively. However, compounds (15-19) did not inhibit contraction of the isolated uterine [32]. Compounds 16, 17, 18 and 19 from petroleum ether extract in the concentration range of 6.25-100  $\mu\text{M}$  were determined about their antiproliferative activity on various cancer cell lines. The results elucidated that compounds 17 and 19 significantly inhibited cancer cells proliferation of human ovarian cancer cells and cervical carcinoma cells. Compound 19 also significantly inhibited endometrial carcinoma cells. In another comparison while compound 19 inhibited human ovarian cancer cell growth with  $\text{IC}_{50}$  of 26.93  $\mu\text{M}$ , compound 17 significantly inhibited with  $\text{IC}_{50}$  of 46.89  $\mu\text{M}$ . This indicates compound 19 has strong effect than 17. However, it is clear that the two compounds showed remarkably antiproliferative activity on human ovarian cancer cells [18].

In the investigation of the gum exudates and resin of *C. myrrha* has led in the isolation of six sesquiterpenoids Myrrhterpenoid K (20), Myrrhterpenoid L (21), Myrrhterpenoid M (22), Myrrhterpenoid N (23), furanogermacr-1(10)Z,4Z-dien-6-one (24) and curzerenone

(13). Compound (20) exhibited weak cytotoxic activity against breast tumor cell line in a clonogenic assay while the other five compounds were inactive [28,33] (Table 1). A sesquiterpen which is commiferin (25) was also isolated by another researcher [29].

A further phytochemical investigation afforded quercetin (26) and  $\beta$ -sitosterol (27) from the aqueous and petroleum ether extracts, respectively [34]. Three cadinane sesquiterpenes, commiterpenes A (28), commiterpenes B (29), commiterpenes C (30) were also isolated from the resinous exudates of *C. myrrha*. All these sesquiterpenes showed neuroprotective effects [28,20]. Extractions with 90% EtOH the resins are largely removed and crude polysaccharides are obtained. After hydrolysis 15 amino acids were detected and in the fractions D-galactose (31), L-arabinose (32), and 4-methyl D-glucuronic acid (33) (in proportions 4:1:3) were identified [29]. Hydrolysis of purified polysaccharides of gum myrrh gave high yields of a mixture of neutral sugars and acidic oligosaccharides. The latter after isolation and purification gave 1:6 mixture of two aldobiuronic acids and identified as 6-O-(4-O-methyl- $\beta$ -D-glucuronosyl)-D-galactose (34) and 4-O-(4-O-methyl- $\alpha$ -D-glucuronosyl)-D-galactose (35) [29]. The UPLC-MS/MS chromatogram analysis of petroleum ether extract identified the following compounds 2-methoxy-8,12-epoxygermacra-1,7,11-triene-6-one (36), docosanoic acid (37), myrrhone (38) and mansumbinone (39) [18].



## *Nigella sativa* L.

The genus *Nigella* belongs to Ranunculaceae family which contains about 20 species. These species are mostly occurring in Europe, the Mediterranean, central Asia and Africa. Even though there are many species in the genus *Nigella*, in Ethiopia only one species (*Nigella sativa* L.) is widely cultivated and the others are an ornamental. *N. sativa* is also cultivated and have been used to treat various diseases in other parts of Africa such as in Kenya, Somalia, Djibouti, Sudan, northern Africa, and outside Africa in Afghanistan, central and southern Europe, India, Iran, Syria, the Soviet Union and other countries [35]. The seed of this plant has various names as black seed, black cumin, black caraway, funnel flower, roman coriander and kalunji. It is found in most local markets and assumed to be grown through-out the agricultural highlands. The local name of this plant in Ethiopia is called "Tikur azmmud" [17].

*N. sativa* is annual herbs and usually profusely branched herb up to 0.7 m tall with finely divided leaves. The leaf segments narrowly linear to threadlike. The stem is ribbed, sometimes hollow when old. The flowers are pale blue or pale purple, with 5-10 petals. The fruit is a capsule composed of several united follicles, each containing numerous seeds [35,36]. Its seeds are small, black and possess aromatic odor and taste [37]. It is an indigenous plant to Ethiopia and most other countries. It is a cultivated plant mostly grown in back gardens of homesteads, on heavier soils as a field crop, tolerating a wide range of soil types, occasionally found and growing wild between 1500-2500 m height [35].

## Medicinal uses

*N. sativa* is an important spice in the culinary to add as flavor. The main uses of this seeds in Ethiopia and other countries are as bread flavor, to prepare hot pepper sauce and dishes. It is mixed with melted butter, wrapped in a piece of cloth and sniffed to relieve some types of headaches. Medicinally it has been used in the folk practice as a natural remedy for a number of diseases such as asthma, hypertension, diabetes, inflammation, cough, eczema, fever and gastrointestinal disturbances. Its seed oil has also antipyretic, analgesic and antineoplastic activity. In Ethiopia it is used to treat headache, stomachache and to induce abortion [35,37]. Bhatti et al. [38] reported that *N. sativa* is a common drug used in Tibbe-Nabvi (Prophet's Medicine) throughout the world because of its therapeutic efficacy and potential to cure disease. Even though *N. sativa* L. used as medicine in different parts of the country, the DNA fingerprints of four Taxa of black seeds from Qassim (Saudi Arabia), Ethiopia, Egypt, and Syria are not identical. The study revealed there are several genetic differences in black seeds Taxa among in these countries [39].

There are ample of scientific evidences for its hypoglycemic, antioxidant, liver protecting, analgesic, antithrombotic, spasmolytic, anti-tumor and antibacterial effects. For example, the research that was done based on identification of the effective of extraction method; the seed was extracted by supercritical carbon dioxide (SCCO<sub>2</sub>) and Soxhlet extraction using various organic solvents. The antibacterial activities of these extracts are investigated by the agar dilution method against gram-positive bacteria (*Bacillus cereus* and *Staphylococcus aureus*) and gram-negative bacteria (*Escherichia coli* and *Yersinia enterocolitica*). As the result revealed the SCCO<sub>2</sub> (at 120 bar/40°C) extract showed effective growth inhibition than solvent extracts against in all the tested bacteria [37]. The result of this investigation has been supported by other research using organic solvent. The MeOH extract exhibited a significant antibacterial activity at the concentration of 50 mg/mL ( $p \leq 0.01$ ) against gram-positive bacteria (*Streptococcus pyogenes*) with a greater inhibition zone of 19 mm than gram negative bacteria (*Pseudomonas aeruginosa*, *Klebsiella pneumonia* and *Proteus vulgaris*) that has 15 mm zone of inhibition. However, *Klebsiella pneumonia* and *Proteus vulgaris* showed resistance against aqueous extract at 20 mg/mL [40]. The ethanol extract of the seed part can also generate antioxidants, possess antitumor activity, and ameliorate and prolong the life span of mice bearing Ehrlich ascites tumor [41].


The study that was conducted to evaluate the *in vitro* and *in vivo* anti-cancer effect of *N. sativa* seed extracts, the essential oil (IC<sub>50</sub>=0.6%, v/v) and EtOAc (IC<sub>50</sub>=0.75%) extracts were more cytotoxic against the murine mastocytoma cell line than the butanol extract (IC<sub>50</sub>=2%). Similar results were also obtained with the kidney carcinoma cell lines of monkeys (Vero). Although all extracts had a comparable cytotoxic effect against the sheep heart carcinoma cell line, with IC<sub>50</sub> values ranging from 0.2 to 0.26% (v/v), tests on the kidney carcinoma cell lines of hamsters revealed a high cytotoxic effect of the EtOAc (IC<sub>50</sub>=0.2%) as compared to the essential oil (IC<sub>50</sub>=1.2%). These data show that

the cytotoxicity of each extract depends on the tumor cell type [42]. The powder of seed of *N. sativa* was also investigated *in vivo* by oral administration to the hypercholesterolemic patients (n=10) at dose of 1 g before breakfast for two months to observe its activity. The result revealed it reduces total cholesterol ( $p=0.002$ ), triglyceride ( $p=0.0001$ ), HDL-cholesterol ( $p=0.0003$ ) and LDL-cholesterol level ( $p=0.002$ ) [38].

In the determination of the toxic effect of *N. sativa* seed powder on the liver function of rats which was evaluated by measuring liver enzymes and through histopathological examination of liver tissue with supplementation up to the dose of 1 g/kg for a period of 28 days, resulted with no changes in liver enzymes level and did not cause any toxicity effect on the liver function [43].


## Volatile compounds

The seed part of *N. sativa* contains up to 40% oil and is used in medicine and perfume industry [17]. Using GC/MS instrument forty-eight compounds have been identified in the essential oil from seed part which is cultivated in Poland. The two monoterpenoids *cis*-4-methoxythujane (40) and *trans*-4-methoxythujane (41) occur as 0.3% and 4%, respectively [44]. Even though in another GC/MS analysis of the volatile oils gave many compounds, the pharmacologically active constituents are thymoquinone (42), dithymoquinone (43), thymol (44) and thymohydroquinone (45) [45]. Comparison analysis of solvent extracts was investigated to show their biological activities. As the research revealed the ether and methanol volatile oil extracts have significant antifungal and antibacterial activities as compared to tetracycline, cefuroxime and ciprofloxacin positive controls, the ether extracts showed stronger activity than methanolic one [46].



## Phytochemicals

According to phytochemical investigation four alkaloids have been reported as the major constituent of *N. sativa* seeds such as nigellidine (46) and nigellidine (47) have an indazole nucleus whereas nigellimine (48) and nigellimine N-oxide (49) are isoquinolines [47-49]. Recently, a triterpene saponin  $\alpha$ -Heredin (50) was isolated from the seeds of *N. sativa*.  $\alpha$ -Heredin is known to have antitumor activity [50]. The ethanolic extract of seed was found to contain three flavonoids namely quercetin (26) and kaempferol 3-glucosyl (51) and quercetin-3-(6-ferululoyl glucosyl) (52) [51]. Other than those triglycoside quercetin 3-glucoside (53), kaempferol 3-glucoside (54) and rutin (55) were also isolated from this seeds [50].



## *Ocimum lamiifolium* Hochst. ex Benth.

The genus *Ocimum* comprises more than 150 species and it is considered as one of the largest genera of the Lamiaceae family [52]. *Ocimum* is a versatile aromatic genus well known for medicinal properties and for economically important essential oils. The genus is very variable and possesses wide range of intra and inter-specific genetic diversity [53].

*Ocimum lamiifolium* Hochst. ex Benth. is an indigenous plant in Ethiopia and locally is called “*dama-kassie*” (in Amharic) and “*anchabi*” (in Oromifa). It is among the commonly occurring *Ocimum spp.* that distributed in different regions of Ethiopia [54]. It is also cultivated and used as medicine in East Africa from Kenya to Malawi, Dem. Rep. Congo, Cameroon and many other countries [55].

*O. lamiifolium* is an erect, robust branching sub shrub or shrub growing about 0.7-3 m tall and indumentum of simple hairs. Leaves are ovate and opposite, and flowers are pinkish in racemes. It grows beside roads and streams, in bush land and at forest edged and on grassland between 1200-2900 m [54]. It has petiolate leaves; which is 0.1-0.5 times as long as the blade, up to 40 mm long [54,55].

### Ethnobotanical uses

Traditionally the leaves part of *O. lamiifolium* is used against eye diseases and headaches [56]. It is also used to relieve pain and fever in a special local preparation in Ethiopia known as “*Yemich medhanit*” [54,55]. The fresh leaves water extract is drunk to treat diarrhoea, amoeba (diarrhoea with blood) and cough/cold [57]. The plant have synergic activity on these diseases when the water extracts of fresh leaves of *Vernonia amygdalina* and *Clutia abyssinica* mixed with root of *O. lamiifolium* [58]. A research which was done in Sheko ethnic group in Southwest Ethiopia, *O. lamiifolium* is used to treat skin and gastrointestinal ailments. In this study the plant has been assigned with the highest fidelity level values, a possible indication of their better healing potential [55]. An ethnomedicinal survey that was conducted on the investigation of knowledge and practice of traditional antimalarial plants, the leaf part of *O. lamiifolium* has been traditionally used for antimalarial activity [59]. The thermal expulsion and direct burning of the leaves part revealed significant repellency against the main vectors of malaria [60].

### Medicinal uses

As the study about the use of medicinal plants in self-care in rural central Ethiopia revealed, *O. lamiifolium* is among the most frequently used plants [57]. The above ethnobotanical uses of this plant were also corroborated by a scientific laboratory based research. *In vitro* research was done by agar disc diffusion method on the aqueous, ethanol and methanol extracts of *O. lamiifolium*. The result revealed the three extracts have antibacterial activity against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Shigella boydii* [61,62]. Evaluation research was done on the antipyretic effects of aqueous and ethanol extracts in mice and the result revealed the extracts have antipyretic property [53]. These extracts were also investigated on anti-inflammatory activities. Both have significant activity but greater anti-inflammatory activity was observed for the aqueous extract. Aqueous extracts of the plant allowed animals to recover barbiturate sleep duration in proportions of 88% [63]. The dried methanolic extract was also tested *in vitro* for its protection activity against acetaminophen-induced hepatotoxicity. The result of this study revealed the plant has activity [64].


### Volatile compounds

*Ocimum spp.* and their essential oils have been traditionally used to kill or repel insects, and also to flavor foods and oral products, in fragrance, in folk medicine and as condiments [65]. Nerio et al. studied and reported that *Ocimum spp.* have promising essential oils and used as insect repellents [66].

The essential oil of *O. lamiifolium* was analyzed by GC and GC/MS and revealed to have major compounds such as Bornyl acetate (56) (30.3%), p-cymene (57) (11.4%),  $\alpha$ -pinene (58) (5.7%),  $\beta$ -pinene (5) (4.2%) and camphene (59) (5.9%). The oils were also evaluated for antimicrobial activity and have showed moderate activity [67]. The antioxidant activity of essential oils and methanolic extracts of *O. lamiifolium* was evaluated by DPPH and linoleic acid peroxidation assays. As the result revealed sabinene (60) (12.2%) and  $\alpha$ -phellandrene (61) (11.6%) volatile oils scavenge DPPH and inhibit linoleic acid peroxidation. This inhibition property suggested that the plant has hepatoprotective activity [68].

### Phytochemicals

The phytochemical compounds are investigated by HPLC, APCI-MS, UV and NMR method and the following flavonoid compounds are isolated and characterized: apigenin (62), genkwanin (63), acacetin (64), apigenin 7,4'-dimethyl ether (65), luteolin (66), ladanein (67), 5,6-dihydroxy-7,3',4'-trimethoxyflavone (68), 5,7-dihydroxy-6,4'-dimethoxyflavone (69), cirsimaritin (70), salvigenin (71), cirsililol (72), cirsilioneol (73), eupatorin (74), 5-hydroxy-6,7,3',4'-tetramethoxyflavone (75) [69]. Using NMR spectroscopy quercetin 3-O-xylosyl(1 $\rightarrow$ 2 $\rightarrow$ ) galactoside (76) was characterised from the leaf extract. *O. lamiifolium* produced larger amounts of flavone 5-O-glycosides (77) [70].



### *Olea europaea* L.

The genus *Olea* belongs to Oleaceae family which comprises about 30 species. Olive (*Olea europaea* L.) is one of the species and native to tropical and warm temperate regions of the world. The tree is usually distributed in many parts of the world and used as table oil in the coastal regions of the Eastern Mediterranean Basin, the neighboring coastal areas of south Eastern Europe, western Asia, northern Africa, Tunisia, California, Chile, Croatia, Argentina, South Africa, Saudi Arabia, Australia as well as northern Iran at the south end of the Caspian and other countries. *O. europaea* is an indigenous plant to Ethiopia and locally known as “*Woirra*” which yield up to 40% olive oil. The fruit of this wild African olive tree is small and the yield of its oil is low. It has been declined in abundance due to over-utilization and exploitation that is not accompanied by a sound conservation strategy [71].

*O. europaea* is a small tree growing up to 5-15 m. It grows between the altitudes of 800-2500 m. Bark is grey to brownish-blackish, smooth to rough with old. Leaves are narrowly oblong-elliptic, 2-10 cm  $\times$  7-17 mm, grey-green to shiny dark green above, greyish or yellowish with a dense covering of silvery, golden or brown scales on the under surface; apex and base narrowly tapering, apex sharp tipped; margin entire,

rolled under and curved back from the midrib, petiole slender, up to 10 mm long, so the leaves tend to droop. Flowers greenish-white or cream, 6-10 mm long, sweetly scented, in loose axillary or occasionally terminal heads, 5-6 cm long [72].

### Medicinal uses

*O. europaea* is a highly valued plant and its leaves, twigs and woods are used to fumigate pots for milk and for the local beverages in Ethiopia known as “Tella” and “Tej”. Twigs are used as toothbrushes (tooth sticks) and hard wood for carving. The roots are used to treat hemorrhoids. More uses are for preventing heart disease, for treating gout, malaria, stomach problems, hemorrhoids, wrinkles, and to treat malaria [73].

Alcoholic olive leaves extract revealed antibacterial effect [24]. Methanol and chloroform extracts of the leaves part have dose-dependent anti-inflammatory and antinociceptive activities [74]. Another research revealed using disc diffusion and microdilution methods the antimicrobial activity of olive leaf extracts was screened against gram-positive and gram-negative bacteria. According to this study the extract revealed antibacterial activity against *Listeria monocytogenes*, *Bacillus cereus*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Proteus vulgaris*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Enterobacter sakazakii*, and *Escherichia coli*. However, the extract has low antibacterial activity against *Salmonella typhimurium* [75]. The olive leaf is useful as an antioxidant [76].


### Volatile components

GC and GC-MS analysis of the essential oil resulted in the identification of 32 compounds representing 99.44% of the oil;  $\alpha$ -pinene (58) (52.70%), 2,6-dimethyloctane (78) (16.57%) and 2-methoxy-3-isopropylpyrazine (79) (6.01%) were the main components. This essential oil showed radical scavenging, anti-inflammatory and analgesic activities [77].

### Phytochemicals

Hydroxytyrosol (80) is mainly found in olive leaves after hydrolysis of oleuropein (81) and has antioxidant, antibacterial and anti-inflammatory properties. An investigation was done to see the effect of hydroxytyrosol for alleviating the pain in patients with gonarthrosis. The result revealed administered hydroxytyrosol has significant improvement [78]. Oleuropein is the major active compound in olive leaf and well known for its benefits for human health. It is considered as premier cooking oil, used for body care. The research study revealed oleuropein (81) is found in the leaves and has significant antispasmodic, antioxidant, diuretic and lipid lowering properties [73]. Extracts of leaves, fruits and seeds of olive tree were analysed by reverse phase HPLC with diode array detection and mass spectrometry. This methodology allowed the identification of verbascoside (82), rutin (55), luteolin-7-glucoside (83), oleuropein (81), nüzhenide (84), oleuropein glucoside (85) and hydroxytyrosol (80) [79].

Pentacyclic triterpenes from *O. europaea* are gaining interest due to their beneficial health effects, as anti-inflammatory, anti-diabetic and anti-tumoral. By LC-MS, APCI and LC-MS instruments the following triterpenes: maslinic (86), oleanolic (87) and ursolic acids (88) along with erythrodiol (89) and uvaol (90) were identified. These compounds are the main triterpenes compounds that present in the fruits and leaves of *O. europaea* [80-82].



### Conclusion

Nowadays, many medicines that have been widely in used come from plant resources. Traditionally, the vast majority of people rely on traditional plant based medicines for primary health care [3]. There are many indigenous plants in the world which have high medicinal activities. To mention some of them *A. corrorima*, *C. myrrha*, *N. sativa*, *O. lamiifolium*, and *O. europaea* are among very well known plants in serving the people as spice, flavor, fumigants, and traditional medicines. Some of their biological activities have been corroborated by scientific researches and have shown antibacterial, antioxidant, antifungal, antimicrobial and anti-inflammatory activities. Based on the literature review, majority of the biological researches on these plants relied on *in vitro* analysis which is pivotal as precursor for further investigation. However, it is pertinent further to investigate on *in vivo* and animal model about their pharmacological and pharmacokinetic activities. This is because by sticking only *in vitro* analysis we cannot be sure about their potent effect on metabolic pathway. Some *in vitro* active plant extract may not be active *in vivo* and animal model because the nature of some active compounds is being metabolized.

The phytochemical investigation also revealed these plants are holder of copious kinds of active compound classes and showed wide spectrum of biological activities. To point out, *A. corrorima* have terpene compounds; *C. myrrha* holder of terpenoid, steroid, flavonoid, glycoside and fatty acid. In *N. sativa* terpenoid, steroid, alkaloid compounds are present. *O. lamiifolium* also afforded in extraction terpenoid and flavonoid compounds while *O. europaea* gave terpenoid and steroidal compounds (Table 1). Hence, from these all kinds of compound classes, terpene compound class is a common type among these five medicinal plants. Moreover, even though each plant gives different kinds of active compounds, there are some chemical compounds that are co-exist in one another in a different proportion such as compound 5 is found in *O. lamiifolium* in 4.2% whereas in *A. corrorima* it is found in a larger amount 17.6%. Compound 58 is found in *O. europaea* in larger amount 52.70% than *O. lamiifolium* 5.70%. Compound 26 is also found in both *N. sativa* and *C. myrrha*. Compound 55 is also found in *O. europaea* and *N. sativa* (Table 1). Therefore, from the point of view of ethno-medicinal usage, still it needs further investigations on the isolation of compounds and cross checking of their pharmacology and pharmacokinetics activity on *in vivo* and model to ascertain those traditional practices are really true and also to see whether these compounds have synergic or additive effects when the local people use these plants as a blend form with other medicinal plants.

### References

1. Qurishi Y, Hamid A, Zargar MA, Singh SK, Saxena AK (2010) Potential role of natural molecules in health and disease: Importance of boswellic acid. Journal of Medicinal Plants Research 4: 2778-2785.
2. Newman DJ, Cragg GM (2012) Natural products as sources of new drugs over the 30 years from 1981 to 2010. Journal of natural products 75: 311-335.

3. Raksha B, Pooja S, Babu S (2014) Bioactive compounds and medicinal properties of Aloe vera L. An update. *Journal of Plant Sciences* 2: 102-107.
4. Fabricant DS, Farnsworth NR (2001) The value of plants used in traditional medicine for drug discovery. *Environ Health Perspect* 109 Suppl 1: 69-75.
5. Wannakrairoj S, Tefera W (2013) In vitro chromosome doubling in korarima (*Aframomum corrorima*) using colchicine and oryzalin. *Kasetsart J. Nat. Sci.* 47: 684-694.
6. Gebreegziabher FG, Mohammed A, Hailemichael G, Jansen PCM (2014) Influence of harvesting stages, drying structures and drying durations on oleoresin and essential oil content of korarima (*Aframomum corrorima*) capsules grown in Ethiopia. *Agricultural Science, Engineering and Technology Research* 2: 1-8.
7. Edwards S, Demissew S, Hedberg I (1997) The Flora of Ethiopia and Eritrea. National Herbarium, Addis Ababa University, Uppsala 6: 324-329.
8. Demissew S, Asfaw N (2009) *Aframomum corrorima* (Braun) Jansen. Aromatic Plants of Ethiopia. Addis Ababa, Shama Books, pp: 178-180.
9. Mitiku H, Mohammed A, Tilahun D, Jansen PCM (2015) Physical and biochemical characteristics of some korarima accessions (*Aframomum corrorima*) collected from major growing regions of southern and southwestern Ethiopia. *Sky Journal of Food Science* 4: 001-009.
10. Eyob S, Martinsen BK, Tsegaye A, Appelgren M, Skrede G, et al. (2008) Antioxidant and antimicrobial activities of extract and essential oil of korarima (*Aframomum corrorima*). *African Journal of Biotechnology* 7: 2585-2592.
11. Eyob S, Appelgren M, Rohloff J, Tsegaye A, Messele G, et al. (2008) Traditional medicinal uses and essential oil composition of leaves and rhizomes of korarima (*Aframomum corrorima*) from southern Ethiopia. *South African Journal of Botany* 74: 181-185.
12. Rice-Evans CA, Miller NJ, Paganga G (1996) Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med* 20: 933-956.
13. Molla BA, Asfaw N, Lwande W (1994) Chemical constituents of the essential oil of *Aframomum corrorima* from Ethiopia. *Ethiop. J. Sci.* 17: 145-148.
14. Chandravanshi BS, Mekassa B (2015) Levels of selected essential and non-essential metals in seeds of korarima (*Aframomum corrorima*) cultivated in Ethiopia. *Braz J Food Technol* 18: 102-111.
15. Tane P, Simplicie DT, Godfred AA, Joseph DC (2005) Bioactive metabolites from *Aframomum* species. Proceedings of the 11th NAPRECA Symposium Hotel Panorama, Antananarivo, Madagascar, pp: 214-223.
16. Nair KPP (2006) The agronomy and economy of Cardamom (*Elettaria cardamomum* M.): The "Queen of spices". *Indian Council of Agricultural Research, New Delhi, India, Advances in Agronomy* 9: 179-470.
17. Seidemann J (2005) *World Spice plants*. Springer, pp: 115-253.
18. Su S, Wang T, Chen T, Duan J, Yu L, et al. (2011) Cytotoxicity activity of extracts and compounds from *Commiphora myrrha* resin against human gynecologic cancer cells. *Journal of Medicinal Plants Research* 5: 1382-1389.
19. Su S, Wang T, Duan JA, Zhou W, Hua YQ, et al. (2011) Anti-inflammatory and analgesic activity of different extracts of *Commiphora myrrha*. *J Ethnopharmacol* 134: 251-258.
20. Xu J, Guo Y, Zhao P, Xie C, Jin DQ, et al. (2011) Neuroprotective cadinane sesquiterpenes from the resinous exudates of *Commiphora myrrha*. *Fitoterapia* 82: 1198-1201.
21. Hedberg I, Edwards S (1989) Flora of Ethiopia and Eritrie: Pittosporaceae to Araliaceae 3. Addis Ababa University, Asmara and Uppsala University, Sweden 3: 442-447.
22. Al-Mobeeriek A (2011) Effects of myrrh on intra-oral mucosal wounds compared with tetracycline- and chlorhexidine-based mouthwashes. *Clin Cosmet Investig Dent* 3: 53-58.
23. Shen T, Lou HX (2008) Bioactive constituents of myrrh and frankincense, two simultaneously prescribed gum resins in chinese traditional medicine. *Chem Biodivers* 5: 540-553.
24. Ahmed AM, Rabii NS, Garbaj AM, Abolghait SK (2014) Antibacterial effect of olive (*Olea europaea* L.) leaves extract in raw peeled undeveined shrimp (*Penaeus semisulcatus*). *International Journal of Veterinary Science and Medicine* 2: 53-56.
25. Nomicos EY (2007) Myrrh: medical marvel or myth of the Magi? *Holist Nurs Pract* 21: 308-323.
26. Chandrasekharnath N, Mahlakshmi YV, Jayalakshmi L, Venkanna B, Uma A (2013) Screening and isolation of bioactive factors from *Commiphora myrrha* and evaluation of their antimicrobial activity. *International Journal of Engineering Research and Applications* 3: 1291-1294.
27. Prakash B, Singh P, Kedia A, Dubey NK (2012) Assessment of some essential oils as food preservatives based on antifungal, anti-aflatoxin, antioxidant activities and in vivo efficacy in food system. *Food Research International* 49: 201-208.
28. Marongiu B, Piras A, Porcedda S, Scorciapino A (2005) Chemical composition of the essential oil and supercritical CO<sub>2</sub> extract of *Commiphora myrrha* (Nees) Engl. and of *Acorus calamus* L. *J Agric Food Chem* 53: 7939-7943.
29. Hanus LO, Rezanka T, Dembitsky VM, Moussaieff A (2005) Myrrh--*Commiphora* chemistry. *Biomed Pap Med Fac Univ, Palacky Olomouc Czech Repub* 149: 3-27.
30. Saeed MA, Sabir AW (2004) Irritant potential of some constituents from oleogum-resin of *Commiphora myrrha*. *Fitoterapia* 75: 81-84.
31. Mohameda AA, Ali SI, EL-Baz FK, Hegazy AK, Kord MA (2014) Chemical composition of essential oil and in vitro antioxidant and antimicrobial activities of crude extracts of *Commiphora myrrha* resin. *Industrial Crops and Products* 57: 10-16.
32. Su SL, Duan JA, Tang YP, Zhang X, Yu L, et al. (2009) Isolation and biological activities of neomyrrh and other terpenes from the resin of *Commiphora myrrha*. *Planta Med* 75: 351-355.
33. Xu J, Guo Y, Zhao P, Guo P, Ma Y, et al. (2012) Four new sesquiterpenes from *Commiphora myrrha* and their neuroprotective effects. *Fitoterapia* 83: 801-805.
34. Ahmed IS, Tohami MSEI, Almagboul AZ, Verpoorte R (2014) Quercetin and  $\beta$ -sitosterol from *Commiphora myrrha* Fum Extracts. *University of Africa Journal of Sciences*, pp: 57-70.
35. Edwards S, Tadesse M, Demissew S, Hedberg I (2000) Flora of Ethiopia and Eritrie. Addis Ababa University and Uppsala University, Sweden 2: 32.
36. Ansari Z, Satish T (2013) Traditional uses of *Nigella sativa*, in Malegaon region of Nashik: A Review. *International Journal of Pure & Applied Bioscience* 1: 19-23.
37. Kumar TVS, Negi PS, Sankar KU (2010) Antibacterial activity of *Nigella sativa* L. Seed extracts. *British Journal of Pharmacology and Toxicology* 1: 96-100.
38. Bhatti IU, Rehman F, Khan MA, Marwat SK (2009) Effect of Prophetic medicine Kalonji (*Nigella sativa* L.) on lipid profile of human beings: an in vivo approach. *World Applied Science Journal* 6: 1053-1057.
39. Al-Huqail A, Al-Saad F (2010) DNA Fingerprinting and Genotyping of Four Black Seed (*Nigella sativa* L.) Taxa. *JKAU: Met. Env and Arid Land Agric Sci* 21: 93-108.
40. Hasan NA, Nawahwi MZ, Malek HAB (2013) Antimicrobial activity of *Nigella sativa* seed extract. *Sains Malaysiana* 42: 143-147.
41. Musa D, Dilsiz N, Gumushan H, Ulakoglu G, Bitiren M (2004) Antitumor activity of an ethanol extract of *Nigella sativa* seeds. *Biologia, Bratislava* 59: 735-740.
42. Mbarek LA, Mouse AH, Elabbadi N, Bensalah M, Gamouh A, et al. (2007) Antitumor properties of black seed (*Nigella sativa* L.) extracts. *Braz J Med Biol Res.* 40: 839-847.
43. Dollah MA, Parhizkar S, Latiff LA, Bin Hassan MH (2013) Toxicity effect of *nigella sativa* on the liver function of rats. *Adv Pharm Bull* 3: 97-102.
44. Wajs A, Bonikowski R, Kalembe D (2008) Composition of essential oil from seeds of *Nigella sativa* L. cultivated in Poland. *Flavour Fragr. J.* 23: 126-132.
45. Hajhashemi V, Ghannadi A, Jafarabadi H (2004) Black cumin seed essential oil, as a potent analgesic and anti-inflammatory drug. *Phytother Res* 18: 195-199.
46. Haloci E, Manfredini S, Toska V, Vertuani S, Ziosi P, et al. (2012) Antibacterial and antifungal activity assesment of *Nigella sativa* essential oils. *International Journal of Medical, Health, Biomedical, Bioengineering and Pharmaceutical Engineering* 6: 270-272.
47. Rehman AS, Malik S, Ahmed I, Rehman H (1985) Nigellimine-N-Oxide, a new isoquinoline alkaloid from seeds of *Nigella sativa*. *Heterocycles* 23: 953-955.
48. Rehman AS, Malik S, Cun HH, Clardy J (1985) Isolation and structure determination of nigellicine, a novel alkaloid from seeds of *Nigella sativa*. *Tetrahedron Lett* 26: 2759-2762.

49. Rehman AS, Malik S, Sadiq H, Choudhary MI, Clardy J (1995) Nigellidine, a new indazole alkaloid from seeds of *Nigella sativa*. *Tetrahedron Lett* 36: 1993-1996.
50. Kumara SS, Huat BT (2001) Extraction, isolation and characterization of antitumour principle, alpha-hedrin, from the seeds of *Nigella sativa*. *Planta Med* 67: 29-22.
51. Merfort I, Wary V, Barakat H, Hussain A, Nawwar AM (1997) Flavonoltriglycosides from seeds of *Nigella sativa*. *Phytochem*. 46: 359-363.
52. Ololade ZS, Fakankun OA, Alao FO, Udi OU (2014) *Ocimum basilicum* var. *purpureum* Floral essential oil: Phytochemicals, phenolic content, antioxidant, free radical scavenging and antimicrobial potentials. *Global Journals Inc*. 14: 2-9.
53. Makonnen E, Debella A, Zerihun L, Abebe D, Tekla F (2003) Antipyretic properties of the aqueous and ethanol extracts of the leaves of *Ocimum suave* and *Ocimum lamifolium* mice. *Journal of Ethnopharmacology* 88: 85-91.
54. Hedberg I, Kelbessa E, Edwards S, Demissew S, Persson E (2006) Flora of Ethiopia and Eritrea: Gentianaceae to Cyclocheilaceae. Addis Ababa, Ethiopia, Uppsala, Sweden 5: 573-575.
55. Giday M, Asfaw Z, Woldu Z (2010) Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *J Ethnopharmacol* 132: 75-85.
56. Fichtl R, Adi A (1994) Honeybee Flora of Ethiopia. Margraf Verlag, Weikersheim, pp: 110-112.
57. Gedif T, Hahn HJ (2003) The use of medicinal plants in self-care in rural central Ethiopia. *J Ethnopharmacol* 87: 155-161.
58. Stark TD, Mtui DJ, Balemba OB (2013) Ethnopharmacological Survey of Plants Used in the Traditional Treatment of Gastrointestinal Pain, Inflammation and Diarrhea in Africa: Future Perspectives for Integration into Modern Medicine. *Animals (Basel)* 3: 158-227.
59. Karunamoorthi K, Tsehay E (2012) Ethnomedicinal knowledge, belief and self-reported practice of local inhabitants on traditional antimalarial plants and phytotherapy. *Journal of Ethnopharmacology* 141: 143-150.
60. Wano M (2006) Evaluation of Essential oils of some local plants for their repellency against *Anopheles arabiensis* and *Aedes aegypti*. Addis Ababa University, Addis Ababa, Ethiopia, pp: 13-16.
61. Gebrehiwot A, Unakal CG (2013) Effect of aqueous and ethanol extracts of *Ocimum lamifolium* and *Amaranthus dubius* against bacteria isolated from clinical specimen. *Int J Pharm and Int Res* 3: 10-14.
62. Damtie D, Mekonnen Y (2015) In vitro evaluation of the antibacterial activities of the methanol, aqueous and n-hexane extracts of *Ocimum lamifolium* from Ethiopia. *African Journal of Pharmacy Pharmacology* 9: 91-97.
63. Woldesellassie M, Eyasu M, Kelbessa U (2011) In vivo anti-inflammatory activities of leaf extracts of *Ocimum lamifolium* in mice model. *J Ethnopharmacol* 134: 32-36.
64. Mukazayire MJ, Allaey V, Calderon PB, Ste'vigny C, Bigendako MJ, et al. (2010) Evaluation of the hepatotoxic and hepatoprotective effect of Rwandese herbal drugs on in vivo (guinea pigs barbiturate-induced sleeping time) and in vitro (rat precision-cut liver slices, PCLS) models. *Experimental and Toxicologic Pathology* 62: 289-299.
65. Padilha de PJ, Gomes CMR, Paumgarten FJ (2003) Chemical composition, toxicity and mosquito repellency of *Ocimum selloi* oil. *J Ethnopharmacology* 88: 253-260.
66. Nerio LS, Olivero-Verbel J, Stashenko E (2010) Repellent activity of essential oils: a review. *Bioresour Technol* 101: 372-378.
67. Runyoro D, Ngassapa O, Vagionas K, Aliannis N, Graikou K, et al. (2010) Chemical composition and antimicrobial activity of the essential oils of four *Ocimum* species growing in Tanzania. *Food Chemistry* 119: 311-316.
68. Mukazayire MJ, Tomani JC, Sté'vigny C, Chalchat JC, Conforti F, et al. (2011) Essential oils of four Rwandese hepatoprotective herbs: Gas chromatography-mass spectrometry analysis and antioxidant activities. *Food Chem* 129: 753-760.
69. Grayer RJ, Veitch NC, Kite GC, Price AM, Kokubun T (2001) Distribution of 8-oxygenated leaf-surface flavones in the genus *Ocimum*. *Phytochemistry* 56: 559-567.
70. Grayer RJ, Kite GC, Veitch NC, Eckerta MR, Marin PD, et al. (2002) Leaf flavonoid glycosides as chemosystematic characters in *Ocimum*. *Biochemical Systematics and Ecology* 30: 327-342.
71. Tadesse M, Mesfin B (2010) A review of selected plants used in the maintenance of health and wellness in Ethiopia. *Ethiopian e-Journal for research and innovative foresight* 2: 85-102.
72. Hedberg I, Edwards S, Nemomissa S (2003) Flora of Ethiopia and Eritrea: Apiaceae to Dipsacaceae. Addis Ababa, Ethiopia, Uppsala University, Sweden 4: 79-80.
73. Aouidi F, Dupuy N, Artaud J, Roussos S, Msallemd M, et al. (2012) Rapid quantitative determination of oleuropein in olive leaves (*Olea europaea*) using mid-infrared spectroscopy combined with chemometric analyses. *Industrial Crops and Products* 37: 292-297.
74. Mahjoub RC, Khemiss M, Dhidah M, Della A, Bouraoui A, et al. (2011) Chloroformic and methanolic extracts of *Olea europaea* L. leaves present anti-inflammatory and analgesic activities. *International Scholarly Research Network Pharmacology*, pp: 1-5.
75. Gökmen M, Kara R, Akkaya L, Torlak E, Önen A (2014) Evaluation of antimicrobial activity in olive (*Olea europaea*) leaf extract. *American Journal of Microbiology* 5: 37-40.
76. Ferreira ICFR, Barros L, Soares ME, Bastos ML, Pereira JA (2007) Antioxidant activity and phenolic contents of *Olea europaea* L. leaves sprayed with different copper formulations. *Food Chemistry* 103: 188-195.
77. Haloui E, Marzouk Z, Marzouk B, Bouftira I, Bouraoui A, et al. (2010) Pharmacological activities and chemical composition of the *Olea europaea* L. leaf essential oils from Tunisia. *Journal of Food, Agriculture and Environment* 8: 204-208.
78. Takeda R, Koike T, Taniguchi I, Tanaka K (2013) Double-blind placebo-controlled trial of hydroxytyrosol of *Olea europaea* on pain in gonarthrosis. *Phytomedicine* 20: 861-864.
79. Silva S, Gomes L, Leitão F, Coelho AV, Boas LV (2006) Phenolic compounds and antioxidant activity of *Olea europaea* L. Fruits and Leaves. *Food Sci Tech Int* 12: 385-396.
80. Giménez E, Juan ME, Melià SC, Barbosa J, Nebot VS, et al. (2015) Pentacyclic triterpene in *Olea europaea* L: A simultaneous determination by high-performance liquid chromatography coupled to mass spectrometry. *Journal of Chromatography A* 1410: 68-75.
81. Solomon MA (2009) Ethiopian herbal medicine practice and the recognition with modern medicine. *Phcog Rev* 3: 44.
82. Balch A (2014) Medicinal plants used in traditional medicine by Oromo people, Ghimbi District, Southwest Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 10: 40.