

Original Research Article

BIOLOGICAL EVALUATION OF SOME NOVEL THIAZOLE , THIAZOLO[3,2-*a*]PYRIDINE AND THIAZOLO[3',2':1,6]PYRIDINE RERIVATIVES CONTAINING DIPHENYL MOIETY AS ANTIMICROBIAL AGENTS

Abd El-Haleem M. Hussein¹, Abu-Bakr A. El-Adasy¹, Ibrahim S. A. Hafi², Esam A. Ishak¹, Emad H. Gawish¹, Mohamed S. A. El-Gaby^{1*}

1. Department of Chemistry, Faculty of Science, Al-Azhar University at Assiut, Assiut 71524, Egypt.
2. Department of Chemistry, Faculty of Education, Suez Canal University, Al-Arish, Egypt.

ABSTRACT

Purpose: The main objective of the present research study is to synthesize some novel thiazole , thiazolo [3,2-*a*] pyridine and thiazolo-[3',2':1,6]pyrido[2,3-*d*]pyrimidine derivatives and evaluate them for their Antimicrobial effect.

Methods: Condensation of 4-phenylmercaptobenzaldehyde **2** with thiosemicarbazide afforded the new thiosemicarbazone derivative **3**. Heterocyclization of thiosemicarbazone **3** with various α -halocarbonyl compounds furnished the novel thiazole derivatives **5**, **6** and **7**. Compound **2** was condensed with 2-cyanomethylene-4-thiazolidinone **8** to yield the 4-thiazolidinone derivative **9**. Cyclocondensation of compound **9** with arylidenemalononitrile (1:1 molar ratio) afforded the novel thiazolo [3,2-*a*] pyridine derivatives **10a-c**. Ternary condensation of compound **2**, malononitrile and thioglycolic acid (2:2:1 molar ratio) yielded the thiazolo[3,2-*a*]pyridine derivative **12**. Compound **12** was cyclized with formic acid and formamide to yield the thiazolo-[3',2':1,6]pyrido[2,3-*d*]pyrimidine derivatives **13** and **14**, respectively. The structures of the newly synthesized compounds were confirmed on the basis of analytical and spectral data. Some of the prepared compounds **3**, **5**, **7**, **9**, **10a,b** and **12** were tested *in vitro* against bacterial and fungal strains.

Results: The results of antimicrobial screening data revealed that most of the synthesized compounds showed varying degrees of inhibition against both bacteria and fungi.

Conclusion: New thiazole and thiazolo[3,2-*a*]pyridine derivatives were prepared from easily accessible starting materials. Some newly synthesized compounds were evaluated for their antibacterial and antifungal activities *in vitro* against four bacteria and two fungi.

Keywords: Thiosemicarbazone, 2-Cyanomethylene-4-thiazolidinone Thiazole, Thiazolo[3,2-*a*]pyridine, Antimicrobial agents.

***Correspondence: Mohamed S A El-Gaby** * Department of Chemistry, Faculty of Science, Al-Azhar University at Assiut, Assiut 71524, Egypt ;Email : m_elgaby@hotmail.com

Running Title-Synthesis and antimicrobial screening of some novel thiazoles.

INTRODUCTION

Aryl sulfide derivatives are a common functionality found in numerous pharmaceutically active compounds. Indeed, a number of drugs in therapeutic areas such as diabetes and inflammatory, immune, Alzheimer's, and Parkinson's diseases contain the aryl sulfide functionality [1-12]. On the other hand, the heterocyclic scaffold comprising thiazole are present in compounds possessing a variety of biological activities such as antibacterial [13], antifungal [14], antioxidant [15], cytotoxic [16], analgesic, anti-inflammatory [17,18], anticonvulsant [19] anti YFV (yellow fever virus), anti-HIV [20] and anti-tubercular [21] and antiviral [22] activities. Also, our search of the literature revealed that, some

thiazolo[3,2-*a*]pyridine derivatives have been reported to possess antibacterial [23], Anti-virulence [24] and anti-hypertensive [25], antioxidant and cytotoxic [26] activities. Prompted by the above facts and in continuation of our efforts in the field of biologically active heterocyclic compounds [27-29], we hereby report the synthesis and antimicrobial evaluation of some novel thiazole and thiazolo[3,2-*a*]pyridine derivatives containing diphenyl sulfide moiety from easily accessible starting materials.

MATERIALS AND METHODS

General:

Melting points were determined on a Gallenkamp apparatus and are uncorrected. The purity of the compounds was checked by TLC. IR spectra were recorded in a Pye-Unicam SP300 instrument in potassium bromide discs. ^1H and ^{13}C NMR spectra were recorded in a Varian Mercury VXR-300 spectrometer (300 MHz for ^1H NMR and 75 MHz for ^{13}C NMR) in $\text{DMSO-}d_6$ and the chemical shifts were related to that of the solvent. Mass spectra were recorded in a GCMS-QP 1000 EX Shimadzu Spectrometer, the ionizing voltage was 70 eV. Elemental analyses were carried out in the Microanalytical Laboratory of Cairo University, Giza, Egypt. 2-Cyanomethylene-4-thiazolidinone **8** was synthesized using method previously published [30]. Antimicrobial activities were carried out at the Regional Center for Mycology and Biotechnology at Al-Azhar University, Cairo, Egypt.

Experimental:

a. 4-Phenylmercaptobenzaldehyde **2**

A mixture of 4-fluorobenzaldehyde (0.01 mol) and thiophenol (0.01 mol) in dimethyl sulfoxide (20 mL) was refluxed in the presence of anhydrous potassium carbonate (2 gm) for 1 hr, left to cool and poured into crushed ice. The obtained solid product was collected by filtration, washed with water and crystallized from acetic acid / water to give **2** as colorless crystals. Yield 92%; m.p. 52-53°C [Lit. 53-54°C] [30]. Anal. Calcd. For $\text{C}_{13}\text{H}_{10}\text{OS}$: C, 72.87; H, 4.72; S, 14.96. Found: C, 72.97; H, 4.65; S, 14.91.

b. 2-(4-(Phenylthio)benzylidene)hydrazine-1-carbothioamide (**3**)

A mixture of 4-phenylmercaptobenzaldehyde **2** (0.01 mol) and thiosemicarbazide (0.01 mol) in ethanol (30 mL) was heated under reflux for 1 h. A crystalline solid was obtained on cooling. It was crystallized from ethanol. Yield 83% as yellow crystals; mp 180–181°C; IR (KBr; cm^{-1}): 3417, 3274, 3135 (NH_2/NH), 3030 (CH_{arom}), 2980 (CH_{aliph}), 1578 ($\text{C}=\text{N}$). ^1H NMR (300 MHz, $\text{DMSO-}d_6$, δ/ppm): 6.23 (br, 2H, NH_2 , exchangeable with D_2O), 7.32 - 7.70 (m, 9H, Ar -H), 8.17 (s, 1H, $\text{CH}=\text{N}$) and 11.40 ppm (s, 1H, NH, exchangeable with D_2O); MS (m/z) 60 (100%), 109 (25%), 211 (52.3%), 287 (21%), 288 (5.2%; M^+). Anal. Calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_3\text{S}_2$ (287.40): C, 58.51; H, 4.56; N, 14.62. Found: C, 58.30; H, 4.60; N, 14.80.

c. General Procedure for the Synthesis of compounds (**5**, **6** and **7**)

A mixture of compound **3** (0.01 mol) and α -halocarbonyl reagents (0.01 mol) **4a-c** and fused sodium acetate (2 gm) in ethanol (30 mL) was refluxed for 3 hrs, left to cool and poured into crushed ice. The obtained solid product was collected by filtration, washed with water and crystallized from the proper solvents.

4-Methyl-2-(2-(4-(phenylthio)benzylidene)hydrazinyl)-thiazole (**5**)

Yield 73% as gray crystals (from 1,4-Dioxane- H_2O); mp 320–321°C; IR (KBr; cm^{-1}): 3432 (NH), 3027 (CH_{arom}), 2986 (CH_{aliph}), 1639 ($\text{C}=\text{N}$). ^1H NMR (300 MHz, $\text{DMSO-}d_6$, δ/ppm): ^1H NMR (300 MHz, $\text{DMSO-}d_6$, δ/ppm): 2.29 (s, 3H, CH_3), 6.18 (s, 1H, thiazole-H), 7.27-7.56 (m, 9H, Ar-H), 7.83 (s, 1H, $\text{CH}=\text{N}$) and 8.56 ppm (s, 1H, NH, exchangeable with D_2O). ^{13}C NMR (75 MHz, $\text{DMSO-}d_6$, δ/ppm): 15.12 (CH_3), 141.16, 138.11, 135.14, 132.34, 131.15, 130.57, 129.25, 129.44, 127.33, 108.15 (Ar-C and Ar-CH),

158.24 (C=N), 153.13 (C=N). MS (m/z): 114 (100%), 184 (18.2%), 325 (16%), 326 (8%; M+1). Anal. Calcd. for C₁₇H₁₅N₃S₂ (325.45): C, 62.74; H, 4.65; N, 12.91. Found: C, 62.90; H, 4.49; N, 12.80.

4-Phenyl-2-(2-(4-(phenylthio)benzylidene)hydrazinyl)thiazole (6)

Yield 75% as brown crystals (from 1,4-Dioxane-H₂O); mp 340–341°C; IR (KBr; cm⁻¹): 3464 (NH), 3045 (CH_{arom}), 2976 (CH_{aliph}), 1638 (C=N). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 6.52 (s, 1H, thiazolo- H), 7.14-7.81 (m, 14 H, Ar-H), 8.7 (s, 1H, CH=N) and 8.96 ppm (s 1H, NH, exchangeable with D₂O). MS (m/z): 77 (100%), 132 (24%), 295 (26.3%), 387 (23%), 388 (5%; M⁺). Anal. Calcd. for C₂₂H₁₇N₃S₂ (387.52): C, 68.19; H, 4.42; N, 10.84. Found: C, 68.10; H, 4.30; N, 10.90.

2-(2-(4-(Phenylthio)benzylidene)hydrazinyl)thiazol-4-amine (7)

Yield 78 % as yellow crystals; mp 310–311°C; (KBr; cm⁻¹): 3434, 3320, 3110 (NH₂/NH), 3048 (CH_{arom}), 2982 (CH_{aliph}), 1638 (C=N). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.60 ppm (2H, br, NH₂, exchangeable with D₂O), 6.20 (s, 1H, thiazole-H), 7.30-8.35 (m, 9H, Ar-H), 8.62 (s, 1H, CH=N) and 11.42 ppm (s, 1H, NH, exchangeable with D₂O). ¹³CNMR (75 MHz, DMSO-*d*₆, δ/ppm): 139.25, 137.45, 134.26, 132.71, 131.30, 130.15, 129.56, 128.69, 127.53, 110.65 (Ar-C and Ar-CH), 156.13 (C=N), 152.28 (C=N). Anal. Calcd. for C₁₆H₁₄N₄S₂ (326.43): C, 58.87; H, 4.32; N, 17.16. Found: C, 58.76; H, 4.26; N, 17.10.

2-(4-Oxo-5-(4-(phenylthio)benzylidene)-4,5-dihydrothiazol-2-yl)-acetonitrile (9)

A mixture of 2-cyanomethylene-4-thiazolidinone **8** (0.01 mol) with 4-phenylmercaptobenzaldehyde **2** (0.01 mol) in ethanol (30 mL) and a few drops of piperidine was refluxed for 1 hour. After cooling, the resulting solid product was collected by filtration, washed with water and the crude product crystallized from ethanol to give **9**. Yield 68 % as orange crystals; mp 310–311°C; (KBr; cm⁻¹): 3040 (CH_{arom}), 2950 (CH_{aliph}), 2200 (C≡N), 1695 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 4.20 (s, 2H, CH₂CN), 7.33-7.78 ppm (m, 10H, Ar-H and methylidene-H). Anal. Calcd. for C₁₈H₁₂N₂OS₂ (336.43): C, 64.26; H, 3.60; N, 8.33. Found: C, 64.19; H, 3.49; N, 8.24.

d. General Procedure for the Synthesis of thiazolo[3,2-a]-pyridines (10a-c)

A mixture of compound **9** (0.01 mol) and arylidenemalononitrile (0.01 mol) in ethanol (30 mL) was treated with a few drops of piperidine and refluxed for 3 hrs. A crystalline solid was obtained on cooling. It was crystallized from an appropriate solvent.

5-Amino-3-oxo-7-phenyl-2-(4-(phenylthio)benzylidene)-2,3-dihydro-7H-thiazolo[3,2-a]pyridine-6,8-dicarbonitrile (10a).

Yield 72% as orange crystals (from Ethanol); mp 214–215°C; IR (KBr; cm⁻¹): 3346, 3221 (NH₂), 3059 (CH_{arom}), 2927 (CH_{aliph}), 2210 (C≡N), 1725 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.60 (s, 1H, pyridine-H), 6.70 (br, 2H, NH₂, exchangeable with D₂O), 7.12-7.48 (m, 14H, Ar-H and methylidene-H). ¹³CNMR (75 MHz, DMSO-*d*₆, δ/ppm): 35.25 (pyridine-C4), 56.14, 76.36, 124.92, 127.45, 128.15, 128.66, 129.51, 131.16, 131.76, 133.26, 133.52, 135.23, 135.64, 143.36, 149.55, 158.19 (Ar-C and Ar-CH), 116.65, 118.75 (2CN), 162.48 (C=O). Anal. Calcd. for C₂₈H₁₈N₄S₂O (490.60): C, 68.55; H, 3.70; N, 11.42. Found: C, 68.41; H, 3.84; N, 11.28.

5-Amino-7-(4-nitrophenyl)-3-oxo-2-(4-(phenylthio)benzylidene)-2,3-dihydro-7H-thiazolo[3,2-a]pyridine-6,8-dicarbonitrile (10b).

Yield 68% as yellow crystals (from Ethanol); mp 228–229°C; IR (KBr; cm⁻¹): 3400, 3340 (NH₂), 3042 (CH_{arom}), 2951 (CH_{aliph}), 2215, 2209 (2C≡N), 1719 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.40 (s, 1H, pyridine-H), 6.50 (br, 2H, NH₂, exchangeable with D₂O), 7.23-7.52 (m, 14H, Ar-H and methylidene-H). MS (m/z): 66 (100%), 109 (43%), 242 (19.5%), 360 (26%), 535 (20%), 536 (8.3%; M⁺). Anal. Calcd. for C₂₈H₁₇N₅S₂O₃ (535.60): C, 62.79; H, 3.20; N, 13.08. Found: C, 62.63; H, 3.26; N, 12.87.

5-Amino-7-(4-methoxyphenyl)-3-oxo-2-(4-(phenylthio)benzylidene)-2,3-dihydro-7H-thiazolo[3,2-a]pyridine-6,8-dicarbonitrile (10c)

Yield 65% as yellow crystals (from Ethanol); mp 216–217°C; IR (KBr; cm⁻¹): 3401, 3364 (NH₂) 3054 (CH_{arom}), 2937 (CH_{aliph}), 2200 (C≡N), 1690 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 2.36 (s, 3H, CH₃), 5.30 (s, 1H, pyridine-H), 6.80 (br, 2H, NH₂, exchangeable with D₂O), 7.12-7.48 (m, 14H, Ar-H, and methylydene-H). Anal. Calcd. for C₂₉H₂₀N₄OS₂ (504.63): C, 69.03; H, 4.00; N, 11.10. Found: C, 68.92; H, 3.84; N, 11.15.

e.5-Amino-3-oxo-2-(4-(phenylthio)benzylidene)-7-(4-(phenylthio)phenyl)-2,3-dihydro-7H-thiazolo[3,2-a]pyridine-6,8-dicarbonitrile (12)

A mixture of 4-phenylmercaptobenzaldehyde **2** (0.02 mol), malononitrile (0.02 mol) and thioglycolic acid (0.01 mol) in ethanol (30 mL) was treated with a few drops of piperidine and refluxed for 3 hrs. The solid product so formed was collected by filtration and crystallized from ethanol to give **12**. Yield 80% as orange crystals; mp 237–238°C; IR (KBr; cm⁻¹): 3383, 3200 (NH₂), 3035 (CH_{arom}), 2950 (CH_{aliph}), 2211, 2197 (2C≡N), 1716 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.62 (s, 1H, H-pyridine) 6.40 (br, 2H, NH₂, exchangeable with D₂O) and 7.30 - 7.78 ppm (m, 19H, Ar-H, and methylydene-H). Anal. Calcd. for C₃₄H₂₂N₄S₃O (598.76): C, 68.20; H, 3.70; N, 9.36. Found: C, 68.11; H, 3.52; N, 9.19.

f.4,9-Dioxo-8-(4-(phenylthio)benzylidene)-5-(4-(phenylthio)phenyl)-3,5,8,9-tetrahydro-4H-thiazolo-[3',2':1,6]pyrido[2,3-d]pyrimidine-6-carbonitrile (13)

A mixture of **12** (0.01 mol) in absolute ethanol (30 mL) and formic acid (5 mL) was refluxed for 5 hrs. After cooling, the solid product was collected by filtration and crystallized from ethanol to give **13**. Yield 62% as brown crystals; mp 210–211 °C; IR (KBr; cm⁻¹): 3444 (NH), 3059 (CH_{arom}), 2215 (C≡N), 1710, 1659 (2C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.61 (s, 1H, pyridine-H), 6.91-7.67 (m, 19H, Ar-H and methylydene-H), 7.90 (s, 1H, pyrimidine-H) and 11.6 (s, 1H, NH, exchangeable with D₂O). Anal. Calcd. for C₃₅H₂₂N₄S₃O₂ (626.77): C, 67.07; H, 3.54; N, 8.94. Found: C, 66.93; H, 3.39; N, 8.68.

g.4-Amino-9-oxo-8-(4-(phenylthio)benzylidene)-5-(4-(phenylthio)phenyl)-8,9-dihydro-5H-thiazolo[3',2':1,6]pyrido-[2,3-d]pyrimidine-6-carbonitrile (14)


A mixture of **12** (0.01 mol) in absolute ethanol (20 mL) and formamide (5 mL) was refluxed for 5 hrs and poured into ice-water. The solid product was collected by filtration, washed with water and crystallized from ethanol to give **14**. Yield 68% brown solid; mp 220–221°C; IR (KBr; cm⁻¹): 3420, 3407 (NH₂), 3068 (CH_{arom}), 2929 (CH_{aliph}), 2211 (C≡N), 1700 (C=O). ¹HNMR (300 MHz, DMSO-*d*₆, δ/ppm): 5.63 (s, 1H, pyridine-H), 6.40 (s, 2H, NH₂, exchangeable with D₂O), 7.20-8.02 ppm (m, 19H, Ar-H and methylydene-H), 8.43 (s, 1H, pyrimidine-H). Anal. Calcd. for C₃₅H₂₃N₅S₃O (625.78): C, 67.18; H, 3.70; N, 11.19. Found: C, 67.10; H, 3.56; N, 11.07.

Antimicrobial screening

Some of the prepared compounds **3**, **5**, **7**, **9**, **10ab**, and **12** were tested *in vitro* against four bacterial strains, *Bacillus cereus*, *Micrococcus luteus*, *Escherichia coli* and *Serratia marcesens* and two strains of fungi, *Candida albicans* and *Aspergillus niger* using the agar cup diffusion (8 mm diameter) assay against the microbial organisms, listed in Table 1 [31]. A 1 mg/mL solution in DMSO was used. The bacteria and fungi were maintained on nutrient agar and Czapek's-Dox agar media, respectively. DMSO showed no inhibition zones. The agar media were inoculated with different microorganism cultures tested. After 24h of incubation at 30 °C for bacteria and 48 h of incubation at 28°C for fungi. Zones of inhibition were recorded in millimeters. Ampicillin in a concentration 25 µg mL⁻¹ and Mycostatine (30 µg mL⁻¹) used as a reference for antibacterial and antifungal activities, respectively. The minimal inhibitory concentration (MIC) of some of the tested compounds was measured by a serial plate dilution method [31].

DISCUSSION


Szmant and co-workers [30] have performed the synthesis of 4-phenylmercaptobenzaldehyde **2** from the reaction of diphenyl sulfide **1** with zinc cyanide in presence of aluminum chloride and stream hydrogen chloride in low yield (45.7%). In our lab, we synthesize 4-phenylmercaptobenzaldehyde **2** via nucleophilic substitution reaction of 4-fluorobenzaldehyde with thiophenol in refluxing dimethyl sulfoxide in the presence of anhydrous potassium carbonate. The product was isolated in 92 % yield (Scheme 1). Thiosemicarbazone derivative **3** was obtained by condensation of 4-phenylmercaptobenzaldehyde **2** with thiosemicarbazide in refluxing ethanol. The molecular structure of thiosemicarbazone **3** was readily established by analytical and spectral data. The infrared spectrum of **3** showed the presence of NH₂ group stretching at 3417, 3274 cm⁻¹ as well as C=N group at 1578 cm⁻¹. The ¹H NMR spectrum was also in accordance with the proposed structure. Also, the mass spectrum of compound **3** showed a molecular ion peak at m/z 288 (5.2 %; M⁺) corresponding to the molecular formula C₁₄H₁₃N₃S₂ and the base peak was found in the spectrum at m/z 60 (100 %) characteristic for the thiocarbamoyl moiety.



Scheme 1 Synthesis of 4-phenylmercaptobenzaldehyde **2** and thiosemicarbazone **3**

The behavior of thiosemicarbazone derivative **3** toward some α -halocarbonyl reagents was investigated with respect to the synthesis of highly substituted thiazoles. Heterocyclization of thiocarbamoyl functional group in compound **3** with chloroacetone **4a** at reflux temperature in the presence of fused sodium acetate furnished the 4-methylthiazole derivative **5** (Scheme 2). The structure of compound **5** was confirmed on the basis of its elemental analysis and spectral data. The infrared spectrum of compound **5** indicated the absence of the thiocarbamoyl group absorption band, indicating the formation of thiazole derivative **5**. In addition, the ¹H NMR spectrum (DMSO-*d*₆) of compound **5** revealed singlet signal at δ 2.29 ppm (3H) for methyl protons, 6.18 ppm (1H) corresponding to thiazole proton at 5-position, 7.83 ppm (1H) attributable to the methine proton and the presence of multiplet signal at δ 7.27-7.56 ppm (9H) for aromatic protons, while the NH proton appears as a broad singlet at δ 8.56 ppm. Also, The mass spectrum of compound **5**, showed a molecular ion peak at m/z 326 which is in agreement with the molecular formula C₁₇H₁₅N₃S₂. The formation of thiazole derivative **5**


is assumed to proceed through initial alkylation by loss of sodium chloride to form intermediate **A**, followed by intramolecular cyclization via elimination of water [28]. In a similar manner, cyclocondensation of thiocarbamoyl functional group in compound **3** with phenacyl bromide **4b** in refluxing ethanol in the presence of fused sodium acetate afforded 4-phenylthiazole derivative **6** (Scheme 2). The mass spectrum of **6** is in accordance with the proposed structure which showed a molecular ion peak at m/z 388 (5%) corresponding to the molecular formula $C_{22}H_{17}N_3S_2$ and the base peak was found in the spectrum at m/z 77 (phenyl moiety). Cycloalkylation of thiocarbamoyl functional group in compound **3** with chloro acetonitrile **4c** by refluxing in ethanol in the presence of fused sodium acetate [28] furnished 4-aminothiazole derivative **7**. The infrared spectrum of the reaction product showed the characteristic absorption bands at 3434, 3320, 3110 cm^{-1} for the NH_2/NH groups and 1578 cm^{-1} for the $C=N$ group. The 1H NMR spectrum (DMSO- d_6) showed a broad singlet at δ 5.60 ppm (br, 2H) assigned to the NH_2 protons, 6.2 (s, 1H), 7.30-8.35 (m, 9H), 8.62 (s, 1H) and 11.42 (s, 1H) ppm which were assigned to the thiazole proton at 5-position, aromatic, methine, and NH protons, respectively. The formation of **7** is assumed to proceed via initial alkylation to form the intermediate **B** followed by nucleophilic addition to the cyano group (Scheme 2).



Scheme 2 Synthesis of thiazole derivatives **5**, **6** and **8**.

Condensation of 2-cyanomethylene-4-thiazolidinone **8** with 4-phenyl-mercaptobenzaldehyde **2** in ethanolic piperidine and reflux yielded 2-(4-oxo-5-(4-(phenylthio)benzylidene)-4,5-dihydrothiazol-2-yl)acetonitrile **9**, (Scheme 3). The molecular structure of **9** was elucidated on the basis of analytical and spectral data. The infrared spectrum showed the appearance of absorption bands at 2200 and 1695 cm^{-1} for $C\equiv N$ and $C=O$ (4-thiazolidinone), respectively. Its 1H NMR spectrum (DMSO- d_6) showed the appearance of a signal for methylene protons at δ 4.20 ppm and 7.33-7.78 ppm for aromatic and methylenidene protons. Cyclization of 4-thiazolidinone derivative **9** with arylidenemalononitrile (1:1 molar ratio) in ethanol containing a catalytic amount of piperidine afforded the novel thiazolo[3,2-*a*]pyridine

derivatives **10a-c** (Scheme 3). The structures of **10** were established on the basis of their elemental analysis and spectral data (IR, ^1H NMR, ^{13}C NMR and MS). The infrared spectra of compounds **10a-c** showed the presence of the NH_2 , $\text{C}\equiv\text{N}$ and $\text{C}=\text{O}$ absorption bands. The ^1H NMR spectrum (DMSO- d_6) of compound **10a** revealed a singlet at δ 5.60 ppm which was assigned to the proton at the 7-position of the thiazolo[3,2-*a*]pyridine ring, a broad singlet at δ 6.70 ppm assigned to the NH_2 protons and a multiplet at δ 7.12-7.48 (m, 14H) assigned to the aromatic and methyldene protons. Moreover, the mass spectrum for the thiazolo[3,2-*a*]pyridine **10b** showed a molecular ion peak at m/z 536 (8.3%) corresponding to the molecular formula $\text{C}_{28}\text{H}_{17}\text{N}_5\text{S}_2\text{O}_3$ with base peak at m/z 66 (malononitrile fragment) The formation of thiazolopyridine **10** is assumed to proceed via *Michael* addition of methylene functional group in compound **9** to the benzylidene moiety to yield *Michael* adduct **C** (Scheme 3), followed by intramolecular cyclization at the cyano group and aromatization [23].




Scheme 3 Synthesis of 4-thiazolidinone **9** and thiazolo[3,2-*a*]pyridines **10a-c**


Ternary condensation of 4-phenylmercaptobenzaldehyde **2**, malononitrile and thioglycolic acid (2: 2: 1 molar ratio) in ethanol at reflux temperature in the presence of piperidine yielded thiazolo[3,2-*a*]pyridine derivative **12** (Scheme 4). The assignment of the structure **12** was based on elemental analysis and spectral data. The infrared spectrum of **12** displayed stretching bands at 3383, 3200 cm^{-1} for the formed amino group and two absorption bands for two $\text{C}\equiv\text{N}$ groups of 2211, 2197 cm^{-1} , in addition to the presence of absorption at 1716 cm^{-1} for the $\text{C}=\text{O}$ group. Its ^1H NMR spectrum (DMSO- d_6) showed appearance of a signal at δ 5.62 ppm for the proton at the 7-position of the thiazolo[3,2-*a*]pyridine ring and broad signal at δ 5.62 ppm for new formed amino protons, in addition to broad signal at δ 7.30-7.78 ppm for aromatic and methyldene protons. The formation of thiazolopyridine derivative **12** can be explained by the reaction pathway depicted in Scheme 5.

Thiazolo[3',2':1,6]pyrido[2,3-*d*] pyrimidine derivative **13** was obtained by refluxing of compound **12** with formic acid [32] (Scheme 6). The structure of compound **13** was confirmed by its analyses and spectral data. The infrared spectrum displayed absorption bands for the NH , $\text{C}\equiv\text{N}$ and two carbonyl functions at 3444, 2215, 1710 and 1659 cm^{-1} , respectively. Its ^1H NMR spectrum (DMSO- d_6) revealed a singlet signal at δ 5.61 ppm for the proton at the 5-position of the thiazolopyridopyrimidine ring, a multiplet at δ 6.91-7.67 ppm assigned to the aromatic and methyldene protons, a singlet signal at δ 7.90 ppm assigned to the proton at the 2-position of the pyrimidine ring while the NH was

appeared at δ 11.6 ppm. The formation of **13** was assumed to take place via the intramolecular cyclization of the intermediate **E** with formic acid (Scheme 1). Cyclocondensation of compound **12** with formamide under reflux furnished the corresponding thiazolo-[3',2':1,6]pyrido[2,3-*d*]pyrimidine derivative **14** (Scheme 6). The structure of this product was established on the basis of its elemental analysis and spectroscopic data.




Scheme 4 One-pot synthesis of thiazolo[3,2-*a*]pyridine **12**



Scheme 5: Possible mechanism for one-pot synthesis of thiazolo[3, 2-*a*]pyridine **12**

The infrared spectrum of the products showed absorption bands corresponding to NH_2 , nitrile and carbonyl groups. The ^1H NMR spectrum was also in accordance with the proposed structure. The

Formation of **14** was assumed to proceed through initial formation of the intermediate **F** followed by intramolecular cyclization via nucleophilic addition of amino group to the cyano group [23] (Scheme 6).



Scheme 6 Synthesis of thiazolo[2,3':6,1]pyrido[2,3-d]-pyrimidines **13&14**

Table 1: Antimicrobial activity of some of the newly synthesized compounds.

Comp. no.	Gram positive bacteria		Gram negative bacteria		Fungi	
	<i>Bacillus Cereus</i>	<i>Micrococcus luteus</i>	<i>Escherichia coli</i>	<i>Serratia marcesens</i>	<i>Candida albicans</i>	<i>Aspergillus niger</i>
3	++	++	++	+++	+	+
5	++	++	++	+++	+	+
7	+++	+	++	+	+	+
9	+	+	++	+	+	+++
10b	+	-	++	+	+	-
10c	+	-	++	++	++	-
12	++	+	++	++	++	+
Standard	++++	++++	++++	++++	++++	++++

+ : Less active (0.1-0.5 cm)

++ : Moderately active (0.6-1.4 cm)

+++ : Highly active (1.5-3.0 cm)

++++ : Very highly activity (over 3.0 cm)

Standard: For Gram positive and Gram negative bacteria: Ampicillin 25 µg mL⁻¹; for fungi: Mycostatine 30 µg mL⁻¹.

Antimicrobial screening

The antibacterial screening revealed that some of the tested compounds showed good inhibition against various tested microbial strains. The result indicated that among the synthesized compounds, **3** and **5** showed good activity against *Serratia marcescens*. Compound **7** also possessed good antibacterial activity against *Bacillus Cereus*. The remaining compounds showed moderated activity against the tested bacterial strains.

The in vitro antifungal data indicated that the compound **10b** exhibited good antifungal activity against the fungi tested *Candida albicans*. Also, compound **9** with a diphenyl sulfide moiety showed excellent activity against *Aspergillus niger*. Compounds **10b,c** were found to be weakly active against both *Micrococcus luteus* and *Aspergillus niger*. The results of the antimicrobial screening of selected new compounds are summarized in Table 1.

REFERENCES

- 1 Youssef M S K, Ahmed R.A. Recent trends in the chemistry of heterocyclic sulfides, Phosphorus, Sulfur and Silicon and the Related Elements, 2006; 181:1123-1199.
- 2 Alcaraz M L, Atkinson S , Cornwall P, Foster A C , Gill D M , Humphries LA , Keegan P S , Kemp R , Merifield E, Nixon R A , Noble A J , O'Beirne D , Patel Z M, Perkins J , Rowan P, Sadler P , Tornos J T , Singleton J, Watts A J, Woodland I A . Efficient syntheses of AZD4407 via thioether formation by nucleophilic attack of organometallic species on sulphur. Org. Process Res. Dev. 2005; 9: 555-569.
- 3 Clader J W, Billard W , Binch H , Chen L Y , Crosby G , Duffy R A , Ford J , Kozlowski J A , Lachowicz J E , Li S J, Liu C , McCombie S W , Vice S, Zhou G W, Greenlee W J. Muscarinic M2 antagonists: anthranilamide derivatives with exceptional selectivity and in vivo activity, Biorg. Med. Chem. 2004; (12) : 319-326.
- 4 Gangjee A, Zeng Y B, Talreja T, McGuire J J, Kisliuk R L, Queener S F. Design and synthesis of classical and nonclassical 6-arylthio-2,4-diamino-5-ethylpyrrolo[2,3-d]-pyrimidines as antifolates , J. Med. Chem .2007; (50):3046-3053.
- 5 Labelle M , Belley M , Gareau Y, Gauthier J Y, Guay D, Gordon R, Grossman S G, Jones T R , Leblanc Y, Mcauliffe M, Mcfarlane C, Masson P, Metters K M , Ouimet N, Patrick D H, Piechuta H, Rochette C, Sawyer N, Xiang Y B, Pickett C B, Fordhutchinson A W, Zamboni R J, Young R N. Discovery of Mk-0476, a potent and orally-Active leukotriene D-4 receptor antagonist devoid of peroxisomal enzyme-induction, Biorg. Med. Chem. Lett. 1995; (5): 283-288.
- 6 Liu G, Huth J R, Olejniczak E T, Mendoza R, DeVries P, Leitz S, Reilly E B, Okasinski G F, Fesik S W, Von Geldern T W. Novel p-arylthio cinnamides as antagonists of leukocyte function-associated antigen-1/intracellular adhesion molecule-1 interaction. 2. Mechanism of inhibition and structure-based improvement of pharmaceutical properties, J. Med. Chem. 2001; 44: 1202-1210.
- 7 Llauger L, He H Z, Kim J, Aguirre J, Rosen N, Peters U, Davies P, Chiosis G. Evaluation of 8-arylsulfanyl, 8-arylsulfoxyl, and 8-arylsulfonyl adenine derivatives as inhibitors of the heat shock protein, J. Med. Chem. 2005; (48): 2892-2905.
- 8 Nielsen S F, Nielsen E O, Olsen G M, Liljefors T, Peters D. Novel potent ligands for the central nicotinic acetylcholine receptor: Synthesis, receptor binding, and 3D-QSAR analysis, J. Med. Chem. 2000 ; 43: 2217-2226.
- 9 Otzen T, Wempe E G, Kunz B, Bartels R , Lehwerk-Yvetot G, Hansel W, Schaper K J, Seydel J K . Folate synthesizing enzyme system as target for development of inhibitors and inhibitor combinations against *Candida albicans*-synthesis and biological activity of new 2,4-

- diaminopyrimidines and 4'-substituted 4-aminodiphenyl sulfones, *J. Med. Chem.* 2004; 47: 240-253.
- 10 Pasquini S, Mugnaini C, Tintori C, Botta M, Trejos A, Arvela R K, Larhed M, Witvrouw M, Michiels M, Christ F, Debyser Z, Corelli F. Investigations on the 4- quinolone-3-carboxylic acid motif. 1. Synthesis-activity relationship of a class of human immunodeficiency virus type 1 integrase inhibitors, *J. Med. Chem.* 2008; 51: 5125-5129.
 - 11 Sun Z Y, Botros E, Su A D, Kim Y, Wang E J, Baturay N Z, Kwon C H. Sulfoxide containing aromatic nitrogen mustards as hypoxia-directed bioreductive cytotoxins, *J. Med.Chem.* 2000; 43:4160-4168.
 - 12 Wang Y G, Chackalamannil S, Hu Z Y, Clader J W, Greenlee W, Billard W, Binch H, Crosby G, Ruperto V, Duffy R A, McQuade R, Lachowicz J E. Design and synthesis of piperidinyl piperidine analogues as potent and selective M-2 muscarinic receptor antagonists, *Biorg. Med. Chem. Lett.* 2000 ; 10:2247-2250.
 - 13 El-Gaby, M S A, El-Hag Ali G A M, El-Maghraby A A, Abd El-Rahman M T, Helal M H M. Synthesis, characterization and in vitro antimicrobial activity of novel 2- thioxo-4-thiazolidinones and 4,4'-bis(thioxo-4-thiazolidin-one-3-yl) diphenylsulfones, *European Journal of Medicinal Chemistry*, 2009; 44 (10): 4148-4152.
 - 14 Aydogan F, Yolacan C. Transformations of adiminies derivatives from pyrrole-2-carbaldehyde.synthesis of thiazolidino-fused compounds, *Bull Korean Chem Soc.* 2001; 22: 476-480.
 - 15 Manojkumar P, Subbuchettiar G. Synthesis of coumarin heterocyclic derivatives with antioxidant activity and in vitro cytotoxic activity against tumor cells, *Acta Pharm.* 2009 ; 59:159- 70.
 - 16 Jubie S, Gowramma B, Nitin K M, Jawahar N, Kalirajan R, Gomathy S. Synthesis and biological evaluation of some 3-(methoxy phenyl)-2-aryl-thiazolidin-4-one derivatives, *Indian. J. Pharm. Sci.*, 2009 ; 1 (1):32-38.
 - 17 Helal M H M, Salem M A, El-Gaby M S A, Aljahdali M. Synthesis and biological evaluation of some novel thiazole compounds as potential anti-inflammatory agents, *European Journal of Medicinal Chemistry*, 2013 ;(65): 517- 526.
 - 18 Chaudhary M, Parmar S, Chaudhary S K, Chaturvedi A K, Ramasastry B V. CNS depressant activity of pyrimidylthiazolidinones. *J. Pharm. Sci.* 1976; 64: 443.
 - 19 Sriram D, Yogeeshwari P, Kumar A. TG. microwave-assisted synthesis and anti-YFV activity of 2,3-diaryl-1,3-thiazolidin-4-one, *J. Pharm. Sci.* 2005 ; 8 (3):426-29.
 - 20 Rawal R. 2-(Aryl)-3-furan-2-ylmethyl-thiazolidin-4-ones as selective HIV-RT inhibitors, *Bioorg. Med. Chem.* 2005 ; 13: 6771-6776.
 - 21 Saeed A, Abbas N, Florke U. Synthesis and antibacterial activity of some novel 2-arylimino-3-aryl-thiazolidin-4-one, *J. Braz. Chem Soc.* 2007;18 (3):559 - 565.
 - 22 Terzioolu N, Karaly N. Synthesis and primary antiviral activity evaluation of 3-hydrazono-5-nitro-2- indolinone derivatives, *Arkivoc*, 2006 ; 2: 109-118.
 - 23 El-Gaby, M S A, Al-Sehemi A G, Mohamed Y A, Ammar Y A, Recent trends in the chemistry of thiazolopyridines. *Phosphorus, Sulfur and Silicon and the Related Elements*, 2006;181(3): 631-674.
 - 24 Chorell E, Pinkner J S, Phan G, Edvinsson S, Buelens F, Remaut H, Waksman G, Hultgren S J, Almquist F. Design and synthesis of C-2 substituted thiazolo and dhydrothiazolo ring-fused 2-pyridones; Pilicides with increased anti-virulence activity, *J. Med. Chem.* 2010; 53(15):5690–5695.
 - 25 Feliciano A S, Caballero E, Puebla P, Pereira J A P, Gras J, Valenti C. Synthesis and pharmacological activities of some pyrido[2,1-b]oxazines, *Eur. J. Med. Chem.* 1992; 27: 527-535.

- 26 Shi F , Li C, Xia M, Miao K, Zhao Y, Tu S, Zheng W, Zhang G, Ma N. Green chemoselective synthesis of thiazolo[3,2-a]pyridine derivatives and evaluation of their antioxidant and cytotoxic activities, *Bioorganic & Medicinal Chemistry Letters*, 2009; 19: 5565–5568.
- 27 El-Gaby M S A, Saleh N M, Micky J A, Ammar Y A, Mohamed H S A. The syntheses of new cylopenta[d][1,3]thiazine derivatives and their use as antimicrobial agents , *Phosphorus, Sulfur and Silicon and the Related Elements*, 2006; 181 (7): 1655-1663 .
- 28 El-Gaby M S A. Syntheses of hitherto unknown thiazole, ylidene and pyridinethione derivatives having a piperidin-1-yl moiety and their use as antimicrobial agents , *Journal of the Chinese Chemical Society*, 2004; 51 (1) :125-134 .
- 29 El-Gaby M S A, Micky J A, Taha N M, El-Sharief M A M Sh. Antimicrobial activity of some novel thiourea, hydrazine, fused pyrimidine and 2-(4- substituted)anilino-benzoazole derivatives containing sulfonamido moieties, *Journal of the Chinese Chemical Society*, 2002;49 (3) :407-414.
- 30 Isidor J L , Mckee R L. Synthesis of 2-methylene -4-thiazolidinone, *J.Org.Chem.* 1973;38(20): 3615-3617.
- 31 Singh A, Wahi A K. Synthesis, characterization and antimicrobial activity of some new Mannich bases derived from 4- amino triazoles. *Research Journal of Pharmaceutical. Biological and Chemical Sciences*, 2011 ; 2: 896–903.