

Biosorption of some Heavy Metals by *Deinococcus radiodurans* Isolated from Soil in Basra Governorate-Iraq

Jaafar R¹, Al-Sulami A¹, Al-Tae A^{2*}, Aldoghachi F³, Suhaimi N⁴ and Mohammed S⁵

¹College of Education for Pure Science, Basra University, Basra, Iraq

²Marine Science Center, Basra University, Basra, Iraq

³College of Science, Basra University, Basra, Iraq

⁴College of Biotechnology, Putra Malaysia University, Malaysia

⁵College of Medicine, Putra Malaysia University, Malaysia

Abstract

The bacterium *Deinococcus radiodurans* has been isolated from Basra soil. On the basis of morphological, biochemical, 16S rRNA gene sequencing and phylogeny analysis, the isolates were authentically identified as *D. radiodurans*. The biosorption capabilities of *D. radiodurans* for cadmium (Cd²⁺) and lead (Pb²⁺) were monitored at different concentrations and contact times. The characterization of heavy metals around the cells of bacterial strains was observed using the Fourier Transform Infrared (FT-IR) spectrophotometer, X-ray powder diffraction analysis (XRD) and transmission electron microscope (TEM).

Keywords: *Deinococcus radiodurans*; Biosorption; Fourier Transform Infrared; X-ray powder diffraction; Transmission electron microscope

Introduction

Bioremediation is a natural process which depends on bacteria, fungi, and plants to alternating pollutants as these organisms carry out their normal life functions. Thus, bioremediation offers a substitute tool to destroy or reduce the harmful pollutants through biological activity with an effective cost. In the early 1980s, certain microorganisms were found to accumulate metallic elements at a high capacity [1,2].

There are a lot of studies about the bioremediation ability of *D. radiodurans* and its genetic engineering, for cleaning up heavy metals in nuclear waste contaminated sites [3-7]. The development of bioremediation strategies using *Deinococcus* spp is therefore vital for the cleanup of contaminated site with radioactive waste. Additional advantages of deinococci are that they are vegetative, easily cultured, and nonpathogenic. As these sites rarely contaminated by a single chemical, it is necessary to bio remediating strain to be multi resistant to various toxic agents. The present work describes the use of a combination of spectroscopic and microscopic methods to characterize the heavy metals around the cells of bacterial strains isolated from extreme habitats as well as to elucidate the interaction mechanisms of these bacteria with these metals.

Materials and Methods

Source of bacterial isolate

The tested isolates, *D. radiodurans* used in this study was isolated previously from the Um - Qasr district, south of Basra city- Iraq. The isolate was identified by biochemical tests and sequencing 16S rRNA gene and comparing the sequences online with GenBank database (www.ncbi.nlm.nih.gov).

Biosorption experiment

The equilibrium, kinetics data of the biosorbent *D. radiodurans* were obtained by performing batch experiments

Characterization study

FTIR analysis: The Fourier Transform Infrared (FT-IR) analysis was done with Perkin Elmer spectrometer model 100 series (sample

preparation UATR). This analysis was done in the Chemistry department, University Putra Malaysia.

X-ray powder diffraction analysis (XRD): The powder X-ray diffraction analysis was performed using a Shimadzu diffractometer model XRD 6000. The diffractometer employed Cu-K α radiation to generate diffraction patterns from powder crystalline samples at ambient temperature. The Cu-K α radiation was generated by Philips glass diffraction, X-ray tube broad focus 2.7KW type. The crystallite size D of the samples was calculated by using the Debye-Scherrer's relationship. Where D is the crystallite size, λ is the incident X-ray wavelength, β is the (FWHM) Full Width at Half-Maximum, and θ is the diffraction angle, Deby -Scherrer equation: $D = K \lambda / \beta \cos \theta$. This analysis was done in the Chemistry department, University Putra Malaysia.

Transmission Electron Microscope: By centrifuging samples broth culture for 10 min at 3000 rpm, and decanting the supernatant, fixing pallet with 4% glutaraldehyde for 4 h at 4°C and centrifuged again, decanting fixative and adding an appropriate quantity of animal serum to submerge sample then allowing serum to clot. Washing three times with 0.1M Cacodylate buffer for 10 min. Postng fix in 1% Osmium tetroxide for 2h at 4°C. Washing again three times with 0.1M Cacodylate buffer for 10 min. dehydrating in series of acetone at 35, 50, 75, 95, and 100% for 10, 10, 10, 10 and 15 min respectively. (Table 1).

Embedding: Specimens were placed into the beam capsule fill with resin and polymerized in oven at 60 °C for 24-48h. Ultra sectioning was realized by choosing an area of interest, for ultrathin section, i.e.

*Corresponding author: Al-Tae A, Marine Science Center, Basra University, Basra, Iraq, E-mail: amrtaee@yahoo.com

Received January 18, 2016; Accepted February 06, 2016; Published February 10, 2016

Citation: Jaafar R, Al-Sulami A, Al-Tae A, Aldoghachi F, Suhaimi N, et al. (2016) Biosorption of some Heavy Metals by *Deinococcus radiodurans* Isolated from Soil in Basra Governorate-Iraq. J Bioremediat Biodegrad 7: 332. doi: [10.4172/2155-6199.1000332](https://doi.org/10.4172/2155-6199.1000332)

Copyright: © 2016 Jaafar R, et al. This is an open-a ccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Acetone	Resin	Time
1	:	1
1	:	3
100%resin		Overnight
100%resin		2hr

Table 1: Finally infiltrating the specimen with acetone and resin.

selecting the silver section, picking up sections with a grid, then drying with filter paper. Finally the sections were stained with Uranyl acetate for 15 min, and washed by double distilled water. Lead sections were stained for 10 min, washed twice with distilled water. This analysis was done in the Electron Microscope Laboratory, Institute of Bioscience, University Putra Malaysia.

Results and Discussion

FT-IR

One of the most important characteristics of a biosorbent is the presence of its surface functional groups, which are largely characterized by the FTIR spectroscopy method. This technique can only provide a qualitative description for biosorbent functional groups. The studies of FTIR spectra on *Deinococcus* provided the basis to interpret the results with *Fucus vesiculosus* [8], as shown in Figure 1 there was a shift in the bands corresponding to carboxyl (COOH) groups after the biosorption process. Following metal binding, the asymmetric carboxyl stretching band shifted from 1630 to 1636 and 1632 Cm^{-1} for Cd and Pb respectively [9], and there was an increase in the distance between this band and the symmetric stretching of the same groups at 1418 Cm^{-1} , to lower wave numbers after biosorption, indicating that chelating complexes were formed [10,11]. Therefore, chelation was another important mechanism involved in the biosorption of Cd and Pb with *Deinococcus*; as reported by Sheng et al. [12] for *Sargassum* with FTIR spectroscopy studies. These changes in the FTIR spectra have also been observed in other biosorption studies [13,14]. Also the peak at 1387 attributed to C-O bond shifted to 1392 and 1390 Cm^{-1} when exposed to Cd and Pb respectively.

Other studies stated that cadmium biosorption was achieved by the formation of ionic bridges between the metal and two carboxyl groups or a bidentate chelating complex with one carboxyl group [15,16]. The symmetric carboxyl-stretching band's intensity decreased after Cd and Pb binding. The bacteria formed stronger bonds with these two metals. Carboxyl groups are the most abundant functional groups and are the main functional groups involved in the biosorption of heavy metals with *Deinococcus* and other biomasses [17-19]. Cadmium biosorption with *Deinococcus* reduced after blocking these groups, the majority of these groups are located in the *Deinococcus* cell wall and their negative charge can attract metal cations [20,21].

XRD

X-ray diffraction is a non-destructive technique used to provide detailed information on the crystallographic structure of materials. This method offers several advantages e.g., non-destructive, high accuracy, capability to detect single crystals, polycrystalline or amorphous materials. Moreover, standards are readily accessible for thousands of material systems. Due to its versatility, XRD has been widely employed to assist in the characterization of biosorbent and in the verification of heavy metals biosorption mechanisms [22]. Figure 2 shows the biosorbent of *D. radiodurans* un-loaded and loaded with Cd (II) and Pb (II). The XRD profile of the unloaded biosorbent shows typical diffraction peaks. Broad peaks were obtained instead of sharp peaks




Figure 1: The FTIR Spectra of *D. radiodurans* with Pb (II), and Cd (II) loaded and unloaded.




Figure 2: XRD analysis of *D. radiodurans* biosorbent before and after Pb and Cd biosorption.

indicating the sample was poorly crystalline. The peaks at 2-theta 30.2, 44, 64.2, and 77.5° corresponding to (201), (114), (641), and (811) planes. The XRD spectrum is compared with the existed spectrums of control that have been published by the Joint Committee on Powder Diffraction Standards (JCPDS file no. 00-001-0166). The XRD spectra of Cd (II) and Pb (II) exhibit strong peaks at 2-theta value 29.9, 44.1, 64.4 and 77.8° corresponding to (200), (114), (640) and (822) planes, respectively, for Cd (II), whereas, the value 44.2, 64.2 and 77.8° corresponding to (112), (211), and (422) for Pb (II). The XRD spectrum is compared with the existed spectrums of control that have been published by the Joint Committee on Powder Diffraction Standards (JCPDS file no. 00-002-097) and (JCPDS file no. 00-002-085) respectively.

The average crystal size of the un-loaded control, loaded with Cd (II) 50 mg l^{-1} , and Pb (II) 50 mg l^{-1} nanoparticle estimated from the highest peak by using the Debye-Scherrer Eq. is 20.0, 27.8, and 14.0 respectively. The results of XRD patterns of *D. radiodurans* show three sharp peaks with a decrease in the intensity of the these peaks for un-loaded biosorbent after exposing the bacteria to Pb (II) and Cd (II), which could explain the immobilization process. This is in agreement with the results reported in the literatures of [23-25]. And this may suggest that Pb (II) and Cd (II) immobilized on surface of *D. radiodurans* [26].

Transmission electron microscope

One of the most important of TEM techniques has revealed the internal structure of the bacteria, especially when embedding bacteria in a resin and staining thin sections of the resulting block with uranyl

acetate. Figure 3a shows the cells of *D. radiodurans* before interaction with metals while Figure 3b,3c show the bacteria after exposure to Pb and Cd solutions. Metals were mostly seen on the cell wall, looking almost like a crust around the cell in addition to the change in the size and the shape of cells. The cellular localization of the metals bound by the cells of study bacteria was located mainly within the cell wall. However, some intracellular metal accumulates associated were also identified in the cytoplasm of the bacterial cells [27]. Reported that, the cellular localization of the uranium bound by the cells was located mainly within the cell wall of three types of *A. ferrooxidans* using




Figure 3a: Transmission electron micrographs of *D. radiodurans* control.




Figure 3b: Transmission electron micrographs of *D. radiodurans* treated with Pb.




Figure 3c: Transmission electron micrographs of *D. radiodurans* treated with Cd showing some unequal cell products (daughter cells) (Scale of bar 0.5μ).

TEM. Electron microscopic observation carried out by Panak et al. [28] revealed the presence of Ag^{+2} as discrete particles at or near the cell wall of both gram positive and gram negative bacteria. Sinha and Mukherjee [29] showed that, *Bacillus* sp. cell wall components with phosphate residues i.e polysaccharides, teichoic and teichuronic acids or phospholipid layers of the membranes can bind U. While cellular functional groups can be responsible for the extracellular association of [30] showed localized intracellular site of accumulated Cd^{+2} by *P. aeruginosa* and showed electron dense grains in the cytosol and towards the cell envelope. Also, Sultan et al. and Singh et al. [31,32] reported that, the cell surfaces of cultures treated with cadmium chloride tended to be rough, suggesting that the cell increased its surface to improve the interaction of toxic substances with the cell surface. The results of the present study also indicate morphology and size changes of bacteria exposed to heavy metals, such changes may due to cell surface changes as a result of heavy metals exposure reported that, the cell surface morphological changes in *Cryptococcus* sp. after exposing to heavy metals, appeared as shrunken and distorted cell wall in the presence of Cd and depressions in the presence of Pb and Zn. Various factors may be responsible for such alterations in cell surface morphology of microbial biomass in the presence of heavy metals. Secretion of extracellular polymeric substance by *Desulfovibrio desulfuricans* during biosorption of Zn and Cu was reported to modify its cell surface morphology [34] Similarly, Murray [35] reported high dark dense cytoplasm due to Co^{+2} precipitations; partially emptied with a very thick cell wall; changing in the morphology of vegetative cells of *B. firmus* and *B. subtilis*.

Conclusion

Deinococcus radiodurans isolated from polluted soils in Basra city south of Iraq, showed high tolerance for Pb and Cd with the possibility of being exploited in bioremediation in two routes, bioaccumulation and biosorption. *D. radiodurans* survive the adverse effect of high concentration of Pb and Cd as expressed in morphological changes.

References

1. Vijayaraghavan K, Yun YS (2008a) Bacterial biosorbents and biosorption. *Biotechnol Adv* 26: 266-291.
2. Vijayaraghavan K, Yun YS (2008b) Biosorption of C.I. Reactive Black 5 from aqueous solution using acid-treated biomass of brown seaweed *Laminaria* sp. *Dyes Pigm* 76: 726-732.
3. Mann H (1990) Removal and recovery of heavy metals by biosorption. In: Volesky B, ed. *Biosorption of heavy metals*: 93-137.
4. Appukuttan D, Rao AS, Apte SK (2006) Engineering of *Deinococcus radiodurans* R1 for bioprecipitation of uranium from dilute nuclear waste. *Appl Environ Microbiol* 72: 7873-7878.
5. Brim H, Osborne JP, Kostandarithes HM, Fredrickson JK, Wackett LP, et al. (2006) *Deinococcus radiodurans* engineered for complete toluene degradation facilitates Cr (VI) reduction. *Microbiol* 152: 2469-2477.
6. Ginn BR, Fein JF (2008) The effect of species diversity on metal adsorption onto bacteria. *Geo Cosmo Acta* 72: 3939-3948.
7. Feng G, Hong CL, Feng XD, Cong Z (2008) Biosorption of radiocesium by *Deinococcus radiodurans* influenced by cations. *J Nucl Agric Sci* 22: 209-212.
8. Sheng PX, Ting YP, Chen JP, Hong L (2004) Sorption of lead, copper, cadmium, zinc, and nickel by marine algal biomass: characterization of biosorptive capacity and investigation of mechanisms. *J Colloid Interface Sci* 275: 131-141.
9. Sethuraman P, Kumar (2011) *Bacillus subtilis* on Pb^{2+} ions removal from aqueous solution by Biosorption. *Res. J Pharmaceut. Biol Chem Sci* 2: 247.
10. Fuks L, Filipiuk D, Majdan M (2006) Transition metal complexes with alginate biosorbent. *J Mol Struct* 792: 104-109.
11. Figueira MM, Volesky B, Mathieu HJ (1999) Instrumental analysis study of iron species biosorption by *Sargassum* biomass. *Environ Sci Technol* 33: 1840-1846.

12. Raize O, Argaman Y, Yannai S (2004) Mechanisms of biosorption of different heavy metals by brown marine macroalgae. *Biotechnol Bioeng* 87: 451-458.
13. Lodeiro P, Barriada JL, Herrero R, Sastre de Vicente ME (2006) The marine macroalga *Cystoseira baccata* as biosorbent for cadmium(II) and lead(II) removal: kinetic and equilibrium studies. *Environ Pollut* 142: 264-273.
14. Chen JP, Yang L (2005) Chemical modification of *Sargassum* sp. for prevention of organic leaching and enhancement of uptake during metal biosorption. *Ind Eng Chem Res* 26: 9931-9942.
15. Nakamoto K (1997) *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, John Wiley & Sons, New York.
16. Fourest E, Volesky B (1996) Contribution of sulfonate groups and alginate to heavy metal biosorption by the dry biomass of *Sargassum fluitans*. *Environ Sci Technol* 30: 277-282.
17. Mehta SK, Gaur JP (2005) Use of algae for removing heavy metal ions from wastewater: progress and prospects. *Crit Rev Biotechnol* 25: 113-152.
18. Volesky B (2003) *Sorption and Biosorption*, BV-Sorbex, Inc., St. Lambert, Quebec.
19. Davis TA, Volesky B, Mucci A (2003) A review of the biochemistry of heavy metal biosorption by brown algae. *Water Res* 37: 4311-4330.
20. Hashim MA, Chu KH (2004) Biosorption of cadmium by brown, green, and red seaweeds. *Chem. Eng. J* 97: 249-255.
21. Liu Y, Xu H (2007) Equilibrium, thermodynamics and mechanisms of Ni²⁺ biosorption by aerobic granules. *Biochem Eng J* 35: 174-182.
22. Chang YC, Chen DH (2005) Preparation and adsorption properties of monodisperse chitosan-bound Fe₃O₄ magnetic nanoparticles for removal of Cu (II) ions. *Coll Interface Sci* 283: 446-451.
23. Li GY, Huang KL, Jiang YR, Yang DL, Ding P (2008) Preparation and characterization of *Saccharomyces cerevisiae* alcohol dehydrogenase immobilized on magnetic nanoparticles. *Int J Biol Macromol* 42: 405-412.
24. Zhu A, Yuan L, Liao T (2008) Suspension of Fe₃O₄ nanoparticles stabilized by chitosan and o-carboxymethylchitosan. *Int J Pharm* 350: 361-368.
25. Li GY, Yu RJ (2008) Preparation and properties of magnetic Fe₃O₄-chitosan nanoparticles. *J Alloys Compd* 466: 451-456.
26. Merroun ML, Raff J, Rossberg A, Hennig C, Reich T, et al. (2005) Complexation of uranium by cells and S-layer sheets of *Bacillus sphaericus* JG-A12. *Appl Environ Microbiol* 71: 5532-5543.
27. Mullen MD, Wolf DC, Ferris FG, Beveridge TJ, Flemming CA, et al. (1989) Bacterial sorption of heavy metals. *Appl Environ Microbiol* 55: 3143-3149.
28. Panak PJ, Raff J, Selenska-Pobell S, Geipel G, Bernhard G, et al. (2000) Complex formation of U (VI) with *Bacillus* isolates from a uranium mining waste pile. *Radiochem Acta* 88: 71-76.
29. Sinha S, Mukherjee SK (2009) *Pseudomonas aeruginosa* KUCD1, a possible candidate for cadmium bioremediation. *Braz J Microbiol* 40: 655-662.
30. El-Helow ER, Sabry SA, Amer RM (2000) Cadmium biosorption by a cadmium resistant strain of *Bacillus thuringiensis*. Regulation and optimization of cell surface affinity for metal cations. *BioMetals* 13: 273-280.
31. Soltan EM, Mohamed RM, Shoreit AA (2008) Behavioral response of resistant and sensitive *Pseudomonas aeruginosa* S22 isolated from Sohag Governorate, Egypt to cadmium stress. *Afr J Biotechnol* 7: 2375-2385.
32. Singh P, Raghukumar C, Parvatkar RR, Mascarenhas-Pereira MB (2013) Heavy metal tolerance in the psychrotolerant *Cryptococcus* sp. isolated from deep-sea sediments of the Central Indian Basin. *Yeast* 30: 93-101.
33. Chen BY, Utgikar VP, Harmon SM, Tabak HH, Bishop DF, et al. (2000) Studies on biosorption of zinc (II) and copper (II) on *Desulfovibrio desulfuricans*. *Int Biodeter Biodegr* 46: 11-18.
34. El-Meleigy MA, Abed NN, Sari IP (2011) Fate of cobalt and nickel in *B. firmus* and *B. subtilis*. *Nat Sci* 9: 175-189.
35. Murray RGE (1992) The family *Deinococcaceae*. *The Prokaryotes Handbook of the Biology of Bacteria: Ecophysiology, Isolation Identification, Application*. Springer-Verlag 4: 3732-3744.

Citation: Jaafar R, Al-Sulami A, Al-Tae A, Aldoghachi F, Suhaimi N, et al. (2016) Biosorption of some Heavy Metals by *Deinococcus radiodurans* Isolated from Soil in Basra Governorate-Iraq. J Bioremediat Biodegrad 7: 332. doi: [10.4172/2155-6199.1000332](https://doi.org/10.4172/2155-6199.1000332)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>