


## Brain Centers Involved in Schizophrenia

Felix-Martin Werner\*

Institute of Neuroscience of Castilla and León, c/o Euro Akademie Pößneck, Germany

Neuronal pathways in the brain regions involved in schizophrenia, the mesolimbic system, the prefrontal cortex and the hippocampus. In this figure, classical neurotransmitters and neuropeptides with a postsynaptic excitatory impulse (a plus mark) and a presynaptic inhibitory impulse (a minus mark) and the corresponding subreceptors are included. Besides the agonistic (a plus mark) and the antagonistic (a minus mark) effects of the antipsychotic drugs: ARP=aripiprazole; ASN=asenapine; CAP=cariprazine; CLZ=clozapine; LUR=lurasidone; OLZ=olanzapine; OSU=OSU-6162; RISP=risperidone; QTP=quetiapine; ZIP=ziprasidone are indicated.



\*Corresponding author: Werner FM, Institute of Neuroscience of Castilla and León, c/o Euro Akademie Pößneck, Germany, Tel: 3647505520; E-mail: [felixm-werner@versanet.de](mailto:felixm-werner@versanet.de)

Received June 29, 2014; Accepted July 01, 2014; Published July 03, 2014

Citation: Werner FM (2014) Brain Centers Involved in Schizophrenia. J Cytol Histol 5: e101. doi:10.4172/2157-7099.1000i101

Copyright: © 2014 Werner FM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.