

## Bulked Segregant Analysis to Detect Main Effect of QTL Associated with Sheath Blight Resistance in BPT-5204/ARC10531 Rice (*Oryza sativa* L)

Shailesh Yadav<sup>1\*</sup>, Ghanta Anuradha<sup>1</sup>, Ravi Ranjan Kumar<sup>3</sup>, Vemireddy LR<sup>1</sup>, Ravuru Sudhakar<sup>2</sup>, Balram M<sup>1</sup> and Siddiq EA<sup>1</sup>

<sup>1</sup>Institute of Biotechnology, ANGRAU, Rajendranagar, Hyderabad-500030 India

<sup>2</sup>Seed Research and Technology Centre, Rajendranagar, Hyderabad-500030 India

<sup>3</sup>Department of Molecular Biology and Genetic Engineering, Bihar Agricultural University, Sabour-813210, India

### Abstract

The population consisting of 210 F2:3 individuals from the cross between BPT-5204 (highly susceptible to sheath blight) and ARC-10531 a land race from Assam (moderately resistant to sheath blight) was analyzed to identify the markers associated with sheath blight resistance and to study any association of any morphological trait to disease incidence. The frequency distribution curve of F2:3 progenies for disease trait were continuous, indicating the polygenic control over the trait. The range of relative lesion height was 21-75% with a mean of 38.59%. No significant association between sheath blight disease and other morphological traits were detected in F2:3 populations. Parental polymorphisms were surveyed with 500 primer pairs of simple sequence repeats (SSR), revealed 70 polymorphic markers between the parents. In order to detect the major effect, QTL associated with sheath blight resistance, a strategy of combining the DNA pooling from selected segregants and genotyping was adopted. The association of putative markers identified based on DNA pooling from selected segregant was established by Single Marker Analysis (SMA). The results of SMA revealed that SSR markers, RM336 (chromosome#7) and RM205 (chromosome#9) showed significant association with sheath blight and accounted for 21.8% and 17.3% of total variation respectively. The results obtained from the DNA pooling of phenotypic extremes could be a useful strategy to detect the genetic loci with major effects of the complex trait such as disease resistance in rice.

**Keywords:** ARC10531; Bulked segregant analysis (BSA); Rice; Sheath blight

### Introduction

In the present scenario of increasing global human population, decreasing arable land, predicted increases of water scarcity, soil salinity, severe diseases, emerging resistance of pests and pathogens to pesticides and climate change pose significant challenges to modern rice research. The biotic stresses *viz.*, blast, stem rot, sheath blight, and bacterial blight diseases causes severe economic losses to rice productivity. Among them Sheath blight (ShB) is an important fungal disease caused by *Rhizoctonia solani* Kuhn causing up to 25% of yield loss and degrades rice quality. In hot and very high humid condition, yield loss can even reach as high as 50%. With the increasing application of nitrogenous fertilizers and the popularization of semi dwarf cultivars with more tillers, ShB is becoming the most serious disease in many rice-producing areas in the world [1-5]. The fungus *R. solani* Kuhn is soil borne pathogen which survives either as sclerotia or mycelia in plant debris. After the initial infection, the pathogen moves on the plant through surface hyphae and develops new infection structures over the entire plant, causing significant necrotic damage. The architecture of the canopy and the associated microclimate has strong effects on both the mobilization of primary inoculums and the further spread of the disease. Absolute resistance to *R. solani* is not available in any of the rice germplasm grown worldwide. However, it has been reported that resistance to *R. solani* is a typical quantitative trait controlled by polygenes in rice [6-12]. In rice because of availability of high resolution molecular maps, complete sequence information and extensive germplasm collections, mapping of quantitative trait loci (QTLs) for disease resistance such as sheath blight is feasible in crop improvement programme. In this context has reported for the first time the identification of rice QTL resistant to ShB using RFLP markers. To date, around 50 ShB resistance QTLs (ShBR QTLs) have been detected over all 12 rice chromosomes in cultivated varieties,

deep-water varieties and wild species. Some of them were identified in multiple mapping populations and/or environments and not associated with either heading date (HD) or morphological traits, and they are believed to be stable ShB QTLs [9,10,12-14]. However QTL mapping is usually carried out by genotyping large number of progenies which is labor intensive, time consuming and cost-ineffective. Several strategies have been proposed to identify molecular markers near a gene/QTL of interest with reduced number of plants to be genotyped. The two main strategies are selective genotyping and bulk segregant analysis (BSA). Selective genotyping is relatively a low-cost approach to detect QTL with large effects by genotyping individuals from the two tails of the phenotypic distribution. Bulk Segregant Analysis (BSA) serves as an affordable strategy for mapping large effect QTLs by genotyping only the extreme phenotypes instead of the entire mapping population. BSA has been successfully used in rice for identifying markers linked to QTL associated with grain quality parameters blast resistance heat tolerance, drought tolerance gall midge resistance and sheath blight resistance. In the present study, bulked segregant analysis approach was used to identify large effect QTLs for sheath blight resistance and to observe the disease association with any of morphological traits [15-26].

**\*Corresponding author:** Shailesh Yadav, Institute of Biotechnology, Acharya N G Ranga Agricultural University, Rajendranagar, Hyderabad, India, Tel: +91-7893473471; E-mail: [shaileshagri9@gmail.com](mailto:shaileshagri9@gmail.com)

**Received** August 22, 2015; **Accepted** September 28, 2015; **Published** October 01, 2015

**Citation:** Yadav S, Anuradha G, Kumar RR, Vemireddy LR, Sudhakar R, et al. (2015) Bulk Segregant Analysis to Detect Main Effect of QTL Associated with Sheath Blight Resistance in BPT-5204/ARC10531 Rice (*Oryza sativa* L). J Rice Res 3: 149. doi:10.4172/2375-4338.1000149

**Copyright:** © 2015 Yadav S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Materials and Methods

### Plant materials

40 rice germplasm lines including 8 wild, 4 land races, 26 cultivated and 2 advanced breeding lines were screened using typha bits method to identify the resistance source for sheath blight disease. A moderate level of resistance to this disease was identified in Tetep and ARC10531, a land race with the relative lesion height percentage of 21-30%. Molecular mapping of QTLs using Tetep as a source for moderate resistance has already been carried out by several research groups. Hence ARC10531 was selected as male parent with highly susceptible elite variety BPT-5204. Crossing program and generation advancement to F<sub>2</sub> was performed at the glass house facility, IBT, ANGRAU, Hyderabad during 2011 and 2012. The polymorphic SSR marker RM 205 and RM22565 were used to fix F<sub>1</sub> progenies and true F<sub>1</sub>'s were forwarded to F<sub>2</sub> generation [27,28] (Table 1).

### DNA extraction and SSR analysis

Genomic DNA was isolated from frozen fresh leaf tissue of 210 F<sub>2:3</sub> progenies along with both the parents (BPT-5204 and ARC10531) with the procedure described by [29]. The quality and quantity of DNA were estimated spectrophotometrically using a Nano Drop (ND-1000, Wilmington, USA). The final DNA concentration was adjusted to 50 ng/μl. Parental polymorphism survey involving 500 SSR markers spanning all twelve chromosomes was carried out. These SSR markers were selected based on uniform distribution across the 12 rice chromosomes. Out of the 500 SSR markers screened, seventy were found polymorphic among the parental lines (BPT-5204 and ARC10531). The PCR reaction for SSR analysis was performed in volumes of 15 μl containing 50 ng genomic DNA, 0.2 μM each primers, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 100 mM each of dATP, dGTP, dCTP and dTTP, 1.5 mM MgCl<sub>2</sub> and 0.5 unit of Taq polymerase. The PCR amplification was performed on Eppendorf Mastercycler<sup>®</sup> Germany with a PCR profile of 94°C for 5 min. followed by 35 cycles of 1 min. at 94°C, 45 sec at 55°C and 1 min at 72°C followed by final extension at 72°C for 10 min. The PCR products were separated on 3.5% metaphor-agarose gel and documented using a gel documentation system (BIORAD Gel Doctm XR, USA).

### F<sub>2:3</sub> phenotyping for sheath blight resistance and other attributes

The 210 individuals of F<sub>2:3</sub> population of the cross (BPT-5204 × ARC10531) were inoculated with the local Rajendranagar isolate (AG1-IA) of *Rhizoctonia solani* which was obtained from Division of Plant Pathology, Directorate of Rice Research, Rajendranagar, Hyderabad-500030 in the year of 2011. The inoculums of virulent 00 isolate were multiplied in Typha stem bits of 4-5 cm long soaked in typha medium (Peptone-10g, Sucrose- 20 g, K<sub>2</sub>HPO<sub>4</sub>-0.1 g, MgSO<sub>4</sub>-0.1 g dissolved in 1000 ml DDH<sub>2</sub>O pH6-6.5) [30]. Rice plants at maximum tillering stage were inoculated with *R. solani* by placing the typha pieces between tillers in the central region of rice hills 5-10 cm above the ground level. *Rhizoctonia solani* infected plants were kept in a humid chamber made of clear plastic for 2 weeks to allow disease development. Plants were grown at 28°C under 14-hrs day light in the humid chamber in the greenhouse. The humidity was maintained between 80 to 100% from the time of inoculation to disease evaluation. To evaluate sheath blight resistance or susceptibility of rice cultivars, the relative lesion height of inoculated plants were recorded 14 days after inoculation. The relative lesion height (RLH) was calculated by the following formula for scoring disease reaction:

$$RLH\% = \frac{\text{Lesion height}(cm)}{\text{Plant height}(cm)} \times 100$$

Observations were recorded 14 days after inoculation and graded as per 0-9 Standard Evaluation System (SES) scale. Following morphological attributes were recorded in segregating populations for a cross BPT-5204 × ARC-10531: Days to heading (days), plant height (cm), panicle length (cm), tiller no (number), effective tiller no (number), relative lesion height (percentage) [30].

### Marker-phenotype association analysis

Bulked segregant analysis (BSA) has been proposed as an efficient

Sr. No.	Genotype
<b>Wild Accessions</b>	
1	<i>Oryza rufipogon</i> AC100488
2	<i>O. rufipogon</i> AC 100368
3	<i>O. rufipogon</i> AC 100490
4	<i>O. rufipogon</i> AC100483
5	<i>O. nivara</i> AC100456
6	<i>O. nivara</i> AC100396
7	<i>O. nivara</i> AC 100395
8	<i>O. nivara</i> AC 100110
<b>Landraces</b>	
9	N-22
10	Tetep
11	Moroberekan
12	ARC 10531
<b>Cultivated</b>	
13	Swarna
14	Rajeswari
15	Swarnadhan
16	Kavya
17	IR-64
18	Lalnakandha
19	Naveen
20	MTU1061
21	Vandana
22	Pusa basmati
23	MTU1001
24	Sonasali
25	BPT-5204
26	Jaya
27	TKM-6
28	Nilagiri
29	Jyothi
30	WGL-32100
31	Ghanteswari
32	MTU-1010
33	WGL-14
34	Chandan
35	Surekha
36	Khandagiri
37	JGL-3844
38	Jaganath
<b>Advanced Breeding Lines</b>	
39	RIL-45
40	RIL-140

**Table 1:** List of rice germplasm screened for Sheath Blight resistant.

strategy for identifying DNA markers linked to the genes or genomic regions of interest [17]. Polymorphic markers may represent markers that are linked to a gene or QTL of interest [32]. DNA bulks of plants with extreme resistance and those with extreme susceptibility were prepared from F<sub>2:3</sub> phenotyped progenies. This was done by pooling aliquots, containing equivalent amounts of total DNA approximately, 50 ng/μl from each of ten highly resistant and ten highly susceptible plants of the F<sub>2:3</sub> based on phenotypic observations. 70 polymorphic SSR primer pairs were used for screening of parents and two bulk DNA samples. DNA of individual F<sub>2:3</sub> plants that were included in bulks were also analyzed with co-segregating markers to confirm their linkage to the sheath blight disease resistance. The SSR markers found polymorphic among the parents and the bulks were used for F<sub>2:3</sub> progeny analysis. DNA of 210 F<sub>2:3</sub> progenies and parents were analyzed to study co-segregation of these markers.

### Data Analysis

The clearly resolved amplicons of SSR were scored manually as homozygote for the allele for susceptible parent (B), homozygote for the allele for resistant parent (A) and heterozygote carrying the alleles from both parents (H) in the data sheet. Chi-square ( $\chi^2$ ) test was performed to test the goodness of fit of the F<sub>2:3</sub> population for the phenotyping and marker data by comparing an observed frequency distribution with an expected one. Marker-trait association was analyzed by simple linear regression method to know the association between the markers and the sheath blight score using software GenStatv14.1 Frequency distribution curve for sheath blight resistance of 210 F<sub>2:3</sub> progenies were drawn separately using Microsoft Office 2010 Excel utility. Variability parameters were performed in F<sub>2:3</sub> population viz., mean, range, skewness, kurtosis, standard deviation and simple correlation coefficients were worked out by software= GENSTAT v14.1[33].

### Results

#### Phenotyping of F<sub>2:3</sub> progenies for sheath blight resistance and other morphological traits

The F<sub>2:3</sub> progenies of the cross ARC10531 and BPT-5204 (210 progenies) were phenotyped for sheath blight during wet season 2012.

The frequency distribution curve of F<sub>2:3</sub> progenies for disease were continuous and near to normal distribution. The range of relative lesion height in percentage was 21-75%. In the F<sub>2:3</sub> progenies, more individuals were distributed towards 40-50% of relative lesion height and population appears to be skewed more towards susceptible side. The mean value recorded for relative lesion height was 38.51%. The results revealed that variability for the morphological traits viz., plant height, heading date, number of tillers, panicle length and disease score (RLH %) ranges from 70- 127, 69-33, 5-31, 5-25, 14-25 and 21-75 respectively. The traits such as plant height, number of productive tillers and relative lesion height exhibit enough variability (Figure 1 and Table 2).

#### Rice microsatellite markers associated with sheath blight reaction using bulk segregant analysis:

Using the BSA method, two bulks having distinct and often contrasting phenotypes for the trait of interest are generated from a segregating population from a single cross. Seventy polymorphic markers were used for screening of parents ARC10531, BPT-5204, resistant bulk (RB), and susceptible bulk (SB) along with individuals of F<sub>2:3</sub> populations used in respective bulks. Two markers RM 205 (chromosome#9) and RM 336 (chromosome#7) clearly distinguished susceptible bulks from resistant bulks. The F<sub>2:3</sub> progenies were genotyped with these two primers (RM336 and RM205) to study their possible association with sheath blight resistance. Segregation pattern with marker RM336 recorded a resistant allele of donor in 43 plants, susceptible allele of recipient was amplified in 57 plants while 110 plants exhibited both the alleles (heterozygous). Similarly for marker RM205 observed 40 plants showing donor allele, 50 plants of recipient allele while remaining 120 plants depicted as heterozygotes. Genetic analysis with chi-square test indicated goodness of fit to the expected ratio of 1:2:1 for co-dominant marker indicating the association of RM336 and RM205 with sheath blight resistant gene in the present population. To determine the strength of association between the putative markers and the respective phenotypes, linear regression analysis was carried out using marker genotype as groups. The simple regression analysis between phenotypic data of sheath blight resistance and the genotypic data of SSR marker RM336 and RM 205 indicated that the marker




Figure 1: Frequency distribution curve of Sheath Blight disease incidence among 210 F<sub>2:3</sub> Progenies derived from the cross (ARC10531 × BPT-5204).

	Plant height (cm)	Heading date(day)	Number of tillers	Number of productive tillers	Panicle length (cm)	Disease Score (RLH %)
Mean	96.53	83.98	15.98	12.42	19.53	38.59
Standard Error	0.98	0.46	0.36	0.31	0.21	1.32
Standard Deviation	13.12	6.22	4.87	4.21	2.78	17.64
Sample Variance	172.17	38.68	23.69	17.71	7.73	311.32
Kurtosis	-0.84	-0.48	-0.06	0.37	-0.57	-0.61
Skewness	0.10	-0.68	0.48	0.79	0.14	0.93
Range	70-127	69-93	5-31	5-25	14-25	21-75

**Table 2:** Variability parameters for different traits of an F2:3 population derived from the cross (BPT-5204x ARC10531).

Source	Degree of freedom (d.f.)	Sum of squares (s.s.)	Mean sum of squares (m.s.s.)	F	P	Percentage phenotypic Variance
Regression	1	28.0	28.0156	44.01	<0.001	17.1
Residual	208	132.4	0.6366			
Total	209	160.4	0.7676			

**Table 3:** Simple linear regression analysis of SSR marker RM205 with phenotypic data of ShB incidence in rice.

Source	Degree of freedom (d.f.)	Sum of squares(s.s.)	Mean sum of squares(m.s.s.)	F	P	Percentage phenotypic Variance
Regression	1	37.5	37.5324	66.11	<0.001	21.8
Residual	208	118.1	0.5677			
Total	209	155.6	0.7446			


**Table 4:** Simple linear regression analysis of SSR marker RM336 with phenotypic data of ShB incidence in rice.

was significantly linked with ShB resistance. The two tagged marker RM205 and RM 336 associated with ShB resistance was mapped too, using Mapmaker 3.0 (Cambridge, MA, USA) based on F2:3 genotyping data of all polymorphic markers identified on those two respective chromosomes. These results indicated the possible detection of two genetic loci for sheath blight resistance on chromosome 7 (based on map location of RM 336) and on chromosome 9 (based on map location of RM 205) (Tables 3 and 4) (Figures 2 and 3).


## Discussion

Several groups have attempted to identify sources of ShB resistance by screening local accessions, cultivars, landraces, and/or advanced breeding lines. The genotypes which were most promising as sources of ShB resistance have been screened for ShB reaction in different rice growing regions [11]. Tetep is a well reported source of resistance to sheath blight and several QTLs have already been mapped by number of researchers [28,29]. The land race ARC 10531 identified in the present study was observed with equal levels of resistance as that of Tetep, used as an alternate source for ShB resistance in process of development of mapping population. The importance of land races in exploring valuable QTLs for sheath blight resistance is previously discussed by several researchers. The F2:3 progenies of the cross ARC10531 and BPT-5204 were phenotyped for sheath blight during Kharif 2012 following standard method of screening in a hot and humid micro-chamber using the typha bits method reported by [31,34,35]. The frequency distribution curve of F2:3 progenies for disease was continuous and

near to normal distribution. Similar frequency distribution curve showing continuous variation involving the F2:3 progeny derived from Japonica cross was reported for ShB resistance. They also observed the curve depicting skewness towards susceptible reaction to disease [36]. Third and fourth degree statistics viz., skewness and kurtosis of an F2:3 progenies suggests that as plant height is positively skewed (0.10) indicates that trait is associated with complementary interactions whereas heading dates (days) negatively skewed (-0.68) and associated with duplicate interaction. The remaining traits also exhibited positive skewness which is associated with complementary interactions. The skewed distribution of sheath blight trait in present study shows that it is under the control of non-additive gene action, especially epistasis and influenced by environmental variables. In the present study, no significant effects on sheath blight disease were detected from all other measured traits in F2:3 populations. Peng and his coworkers, 2003 concluded that for most of the morphological traits, their inheritance was independent of sheath blight resistance, and the correlations between them and sheath blight resistance were not significant or they were inconsistent in a population derived from a Jasmine85/Lemont and by [28] in a population derived from a HP2216/Tetep. All these studies clearly showed that there are many ShB QTLs that were mapped independent of PH (plant height) and HD (heading



**Figure 2:** A): Alleles showing co-segregation of the SSR marker RM205 among the parents, bulks and F2:3 individuals; B): Alleles showing co-segregation of the SSR marker RM336 among the parents, bulks and F2:3 individuals (M-100 bp ladder; P1-ARC10531; P2-BPT- 5204 RB- Resistant bulk; SB-Susceptible bulk).



**Figure 3:** SSR markers RM336 and RM 205 associated with ShB resistance positioned on the rice chromosome #7 and #9.

	Plant height (cm)	Heading date (days)	Number of tillers	Number of productive tillers	Panicle length (cm)	(RLH %)
Plant height (cm)	1.0000					
Heading date (days)	-0.3055**	1.0000				
Number of tillers	0.0076	-0.0253	1.0000			
Number of productive tillers	-0.0167	0.0437	0.8861**	1.0000		
Panicle length (cm)	0.1141**	-0.0203	0.0839	0.0926	1.0000	
Disease Score (RLH %)	-0.0199	-0.0323	-0.0073	-0.0270	0.0266	1.0000

**Table 5:** Correlation coefficients among different plant attributes of an F<sub>2</sub> population (\*\* P<0.01).

date) on all rice chromosomes Some reports have shown correlation of sheath blight with plant height or heading date studied that sheath blight resistance as they measured it in Tetep might be due to the molecular mechanisms involved in host-pathogen interaction and not due to any morphological adaptation to avoid disease. Similarly in our study with ARC10531 we did not find any morphological correlation with disease score [12,28,37]. In our study the SSR markers RM205 and RM336 were found associated with sheath blight resistance gene/locus in the rice cultivar ARC10531 using bulked segregant analysis. Similarly identified three major QTLs for sheath blight resistance in an F<sub>2</sub> population of Jasmine 85/Lemont through bulked segregant analysis in their study performed the BSA using F<sub>2</sub> population from the cross between Lemont and Teqing. The two major QTLs for rice sheath blight identified on chromosome 9 and 11 with R<sup>2</sup> value of 12.9% and 15.3% developed an F<sub>2</sub> population from a cross between 4011 and Xiangzaoxian19 to identify molecular markers linked with the sheath blight resistance using BSA. The dominant resistant gene/locus named as *Rsb1* was mapped on rice chromosome 5 linked with SSR marker RM164 [19] have discussed the prospects of bulk segregant analysis in a broad range of applications in gene mapping [26,27,38]. Furthermore the genetic map of chromosome #7 and chromosome #9 was constructed to verify the existence of previous reports of QTL region linked with these two markers RM 336 and RM 205 associated with ShB resistance identified in the present study. On chromosome #7 the linked marker RM336 are in agreement with the previously reported result of [28]. Chromosome #9 of rice has been reported to contain many major effect ShB QTLs, most of which are closely mapped to each other. In the present study, SSR marker RM205 found associated with ShB resistance has been mapped earlier by [39]. The consistent QTL for ShB resistance on chromosome #9 has verified by several other researchers which indicate its authenticity and stability. It once again signifies the importance of BSA in establishing marker-trait association in a rapid way. The findings of this study could be directly useful in molecular analysis of segregating generations, breeding lines and varieties having ARC10531 as a parent [10,40] (Table 5).

## References

1. Khush GS (2005) What it will take to feed 5.0 billion rice consumers in 2030. See comment in PubMed Commons below Plant Mol Biol 59: 1-6.
2. Gao JP, Chao DY, Lin HX (2007) Understanding abiotic stress tolerance mechanisms: recent studies on stress response in rice. J Integrated Plant Biol 49:742-750.
3. Zhang Q (2007) Strategies for developing Green Super Rice. See comment in PubMed Commons below Proc Natl Acad Sci USA 104: 16402-16409.
4. Meng QZ, Liu ZH, Wang HY, Zhang SS, Wei SH (2001) Research progress in rice sheath blight. J Shenyang Agricultural University 32:376-381.
5. Liao HN, Xiao LS, Wang HS (1997) Analysis of sheath blight developing history and evolving matter Guangxi. Plant Protection 3: 35-38.
6. Ou SH (1985) Rice Diseases, 2nd edn. Commonwealth mycological institute.
7. Castilla NP, Leano RM, Elazegui FA, Teng PS, Savary S (1996) Effects of plant contact, inoculation pattern, leaf wetness regime and nitrogen supply on inoculum efficiency in rice sheath blight. J Phytopathology 144:187-192.
8. Savary S, Castilla N, Elazegui FA, McLaren C, Ynalvez MA, et al. (1995) Direct and indirect effects of nitrogen supply and disease source structure on rice sheath blight spread. Phytopathology 85: 959-965.
9. Jia Y, Liu G, Costanzo S, Lee S, Dai Y (2009) Current progress on genetic interactions of rice with rice blast and sheath blight. Frontier Agric China. 3:231-239.
10. Zuo SM, Zhang YF, Chen ZX, Chen XJ, Pan XB (2010) Current progress on genetics and breeding in resistance to rice sheath blight. Scientia Sin Vitae 40:1014-1023.
11. Srinivasachary WL, Savary S (2011) Resistance to rice sheath blight (*Rhizoctonia solani*) disease: current status and perspectives. Euphytica 178: 1-22.
12. Li Z, Pinson SR, Marchetti MA, Stansel JW, Park WD (1995) Characterization of quantitative trait loci (QTLs) in cultivated rice contributing to field resistance to sheath blight (*Rhizoctonia solani*). See comment in PubMed Commons below Theor Appl Genet 91: 382-388.
13. Xu Q, Yuan XP, Yu HY, Wang YP, Tang SX, et al. (2011) Mapping quantitative trait loci for sheath blight resistance in rice using double haploid population. Plant Breeding 130: 404-406.
14. Wang Y, Pinson SRM, Fjellstrom RG, Tabien RE (2012) Phenotypic gain from introgression of two QTL, qSB9-2 and qSB12-1 for rice sheath blight resistance. Mol Breeding 30: 293-303.
15. Navabi A, Mather DE, Bernier J, Spaner DM, Atlin GN (2009) QTL detection with bidirectional and unidirectional selective genotyping: Marker-based and trait-based analyses. Theo Appl Genet 118: 347-358.
16. Sun Y, Wang J, Crouch JH, Xu Y (2010) Efficiency of selective genotyping for genetic analysis of complex traits and potential applications in crop improvement. Mol Breeding 26: 493-511.
17. Michelmore RW1, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. See comment in PubMed Commons below Proc Natl Acad Sci USA 88: 9828-9832.
18. Salunkhe AS, Poornima R, Prince KSJ, Kanagaraj P, Sheeba JA, et al. (2011) Fine mapping QTL for drought resistance traits in rice (*Oryza sativa* L) using bulk segregant analysis. Molecular Biotechnology doi: 10.1007/s12033-011-9382-x.
19. Govindaraj P, Arumugachamy S, Maheswaran M (2005) Bulk segregant analysis to detect main effect QTL associated with grain quality parameters in Basmati 370/ASD 16 cross in rice (*Oryza sativa* L) using SSR markers. Euphytica 144:61-68.
20. Yang Q, Lin F, Wang L, Pan Q (2009) Identification and mapping of Pi41, a major gene conferring resistance to rice blast in the *Oryza sativa* subsp. indica reference cultivar, 93-11. See comment in PubMed Commons below Theor Appl Genet 118: 1027-1034.
21. Kumbhar SD, Kulwal PL, Patil JV, Gaikwad AP, Jadhav AS (2013) Inheritance of blast resistance and identification of SSR marker associated with it in rice cultivar RDN 98-2. See comment in PubMed Commons below J Genet 92: 317-321.
22. Zhang GL, Chen LY, Xiao GY, Xiao YH, Chen XB, et al. (2009) Bulk segregant analysis to detect QTL related to heat tolerance in rice (*Oryza sativa* L) using SSR markers. Agric Sci China 8(4):482-487.
23. Venuprasad R, Zhao D, Espiritu M, Amante M, Kumar A, et al. (2009) Identification and characterization of large-effect quantitative trait loci for grain

- yield under lowland drought stress in rice using bulk-segregant analysis. *Theor Appl Genet* 120:177-190.
24. Lima JM, Dass A, Sahu SC, Behera L, Chauhan DK (2007) A RAPD marker identified a susceptible specific locus for gall midge resistance gene in rice cultivar ARC5984. *Crop Protection* 26:1431-1435.
25. Xuemei J, Pan X (2001) Quantitative trait loci controlling sheath blight resistance in two rice varieties. Masters dissertation. Yangzhou University 1-89.
26. Che KP, Zhan QC, Xing QH, Wang ZP, Jin DM, et al. (2003) Tagging and mapping of rice sheath blight resistant gene. See comment in PubMed Commons below *Theor Appl Genet* 106: 293-297.
27. Sato HI, Deta O, Audo J, Kunihiro Y, Hirabayashi H, et al. (2004) Mapping QTLs for sheath blight resistance in the rice line WSS2. *Breeding Sci* 54: 265-271.
28. Channamallikarjuna V, Sonah H, Prasad M, Rao GN, Upreti HC, et al. (2010) Identification of major quantitative trait loci qSBR11-1 for sheath blight resistance in rice. *Mol Breeding* 25:155-166.
29. Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. See comment in PubMed Commons below *Nucleic Acids Res* 8: 4321-4325.
30. IRRI (2002) Standard Evaluation System for Rice. 5th ed. INGER Genetic Resources Centre, IRRI, Manila, Philippines, 20-21.
31. Bhaktavatsalam G, Satyanarayana K, Reddy APK, John VT (1978) Evaluation of sheath blight resistance in rice. *Int Rice Res Newsletter* 3:9-10.
32. Collard BCY, Jahufer MZZ, Brouwer JB, Pang EK (2005) An introduction to markers, quantitative trait loci (QTL) mapping and marker assisted selection for crop improvement: the basic concepts. *Euphytica* 142: 169-196.
33. Payne RW, Murray DA, Harding SA, Baird DB, Soutar DM (2011) *An Introduction to GenStat for Windows (14th Edition)* VSN International, Hemel Hempstead, UK.
34. Manian S, Rao KM (1979) Resistance to sheath blight disease in India. *Int Rice Res Newsletter* 4:5-6.
35. Taguchi-Shiobara F, Ozaki H, Sato H, Maeda H, Kojima Y, et al. (2013) Mapping and validation of QTLs for rice sheath blight resistance. See comment in PubMed Commons below *Breed Sci* 63: 301-308.
36. Sharma A, McClung AM, Pinson SRM, Kepiro JL, Shank AR, et al. (2009) Genetic mapping of sheath blight resistance QTL within tropical japonica rice cultivars. *Crop Sci* 49:256-264.
37. Zou JH, Pan XB, Hen JY, Xu JF, Lu WX, et al. (2000) Mapping quantitative trait loci controlling sheath blight resistance in two rice cultivars (*Oryza sativa* L). *Theor Appl Genet* 101:569-573.
38. Pan XB, Rush MC, Sha XY, Xie QJ, Linscombe SD, et al. (1999) Major gene, non-allelic sheath blight resistance from the rice cultivars Jasmine85 and Teqing. *Crop Sci* 39: 338-346.
39. Tan CX, Ji XM, Yang Y, Pan XY, Zuo SM, et al. (2005) [Identification and marker-assisted selection of two major quantitative genes controlling rice sheath blight resistance in backcross generations]. See comment in PubMed Commons below *Yi Chuan Xue Bao* 32: 399-405.
40. Pinson SRM, Capdevielle FM, Oard JH (2005) Confirming QTLs and finding additional loci conditioning sheath blight resistance in rice using recombinant inbred lines. *Crop Sci* 45: 503-510.

**Citation:** Yadav S, Anuradha G, Kumar RR, Vemireddy LR, Sudhakar R, et al. (2015) Bulk Segregant Analysis to Detect Main Effect of QTL Associated with Sheath Blight Resistance in BPT-5204/ARC10531 Rice (*Oryza sativa* L). J Rice Res 3: 149. doi:[10.4172/2375-4338.1000149](https://doi.org/10.4172/2375-4338.1000149)

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>