

Can We Slowdown Early Vascular Aging and Cognitive Disabilities in High Risk Young Adults?

Rajajeyakumar Manivel^{1*}, Chenniappan M², Shanmugavelu M³, Balachander Jayaraman⁴, Basco B⁵, Ravindranath V⁶ and Janitha Alagarsamy⁷

¹ Assistant Professor, Department of Physiology, Chennai medical college hospital & research Centre, Trichy, Tamilnadu, India

² Consultant Cardiologist, Ramakrishna Hospitals & Adjunct Professor, Dr.MGR. Medical University, Trichy, Tamilnadu, India

³ Chairman, Trichy Diabetes Speciality Centre, Tamilnadu, India

⁴ Professor, Department of Cardiology & Medical Superintendent, Jawaharlal Institute of Postgraduate Medical Education & Research, Pondicherry, India

⁵ Vice chairman, Trichy Diabetes Speciality Centre, Tamilnadu, India

⁶ Managing Director, Trichy Diabetes Speciality Centre, Tamilnadu, India

⁷ Senior Medical Officer, Bharat Heavy Electricals Limited, Tamilnadu, India

*Corresponding author: Rajajeyakumar Manivel, Department of Physiology, Chennai medical college hospital & research Centre, Trichy, Tamilnadu, India, Tel: 09751382650; E-mail: rajakumar60@gmail.com

Rec date: September 25, 2014, Acc date: September 27, 2014, Pub date: September 30, 2014

Copyright: © 2014. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

In recent years, the new concept of early vascular aging (EVA) has emerged as a useful tool to aid understanding of how cardiovascular risk increases in relation to the biological aging process. The core feature of EVA is arterial stiffness, which can be measured as increased pulse wave velocity in relation to a subject's chronological age and sex [1]. EVA is the early arterial aging observed in young patients with essential hypertension compared with age- and gender-matched normotensive individuals [2]. Pulse pressure (an indirect measure of arterial stiffness) is a robust predictor of cardiovascular events. One independent predictor of cardiovascular morbidity and mortality in hypertensive patients is pulse pressure [3]. Although pulse pressure derives from the interaction of cardiac ejection (stroke volume) and the properties of arterial circulation (arterial stiffness and wave reflection), elevated pulse pressure is thought to be largely associated with increased arterial stiffness due to aging, arteriosclerosis, or both [3,4]. Orthostatic hypotension (OH) is one clinical manifestation of diabetic autonomic neuropathy and is also a significant risk for fall, syncope, cardiovascular disease and all-cause mortality. Mean arterial pressure and pulse pressure are keen to detriment of cerebral perfusion and cognitive performance [5].

Non invasive method for early detection of early vascular aging and cognitive disabilities:

I. **High risk individuals:** They are Young adult offspring's of known Type 2 diabetic and hypertensive parents aged between 18-25 years. Early identification of vascular Dysfunctions by using following non-invasive techniques and simple scoring system.

II. Non-invasive Parameters to be studied:

1. Hypertension Risk Prediction from the Framingham Heart Study [6].
2. Indian Diabetic Risk Score (IDRS [7].

3. Anthropometry :

BMI was calculated by the formula $\text{Weight (kg)} = \{\text{Height (m)}\}^2$.

Waist to hip ratio (WHR).

4. Physiological parameters:

- Resting average heart rate,
- Resting systolic blood pressure,
- Diastolic blood pressure,
- Mean arterial blood pressure,
- Pulse pressure.
- Arterial wave form analysis & augmentation index [1]
- Resting short term 5 min ECG Recording for HRV (Heart rate variability) [8]
- 5. **Neuro-cognitive test battery [9,10]**
 - Trail making test A.
 - Trail making test B.
 - Digit span test – Forward & Reverse.
 - Letter cancellation test.

Discussion, Conclusion & Recommendation:

Future prevention of premature vascular aging will be achieved by early identification of high risk group and life style modification is the first intervention to consider in adults followed by drug therapy to control risk factors such as hypertension, increased blood pressure variability, hyperglycaemia, and hyperlipidaemia. These implementations should start early in life to prevent cardiovascular diseases, maintain good quality of life and thereby increase the life expectancy.

Acknowledgment:

We acknowledge Dr. Madanmohan, Professor and Head, Department of Physiology, Mahatma Gandhi Medical College and Research Institute, Dr. Vivek kumar Sharma, Dr. Bhattacharjee M and Dr. Amudharaj D, Assistant Professors, Department of Physiology, JIPMER, Vardhman Mahavir Medical College and Safdarjung Hospital, Delhi and Aarupadai Veedu Medical College, Pondicherry, who supported us for all our academic events.

References:

1. Nilsson PM, Lurbe E, Laurent S (2008). The early life origins of vascular ageing and cardiovascular risk: the EVA syndrome. *J Hypertens* 26: 1049-1057.
2. Franklin SS (2002) Do diabetes and hypertension interact to accelerate vascular ageing? *J Hypertens* 20: 1693-1696.
3. Dart AM, Kingwell BA (2001)Pulse pressure: a review of mechanisms and clinical relevance. *J Am Coll Cardiol* 37: 975-984.
4. Franklin SS, Gustin W, Wong ND, Larson MG, Weber MA, et al. (1997)Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study *Circulation* 96: 308-315.
5. Wu JS, Yang YC, Lu FH, Wu CH, Wang RH, et al. (2009) Population-based study on the prevalence and risk factors of orthostatic hypotension in subjects with pre-diabetes and diabetes. *Diabetes Care* 32: 69-74.
6. Parikh NI, Pencina MJ, Wang TJ, Benjamin EJ, Lanier KJ (2000) A Risk Score for Predicting Near-Term Incidence of Hypertension: The Framingham Heart Study' *Ann Intern Med* 148: 102-110.
7. Mohan V, Deepa R, Deepa M, et al. (2005) A Simplified Indian Diabetes Risk Score for Screening for Undiagnosed Diabetic Subjects. *J Assoc Physicians India* 53: 759-763.
8. (1996) Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation*. 93: 1043-1065.
9. Sharma VK, Das S, Mondal S, Goswami U, Gandhi A (2006)Effect of sahaj yoga on neuro cognitive functions in patients suffering from major depression. *Indian J Physiol Pharmacol* 50: 375-383.
10. Sharma VK, Rajajeyakumar M, Velkumary S, Subramanian SK, Bhavanani AB, et al. (2014)Effect of Fast and Slow Pranayama Practice on Cognitive Functions in Healthy Volunteers. *J Clin Diagn Res* 8: 10-13.