

Carcinogenesis and Sex Hormones: A Review

Ankita Bohra* and Sumit Bhateja

Department of Oral Medicine Diagnosis & Radiology, Vyas Dental College & Hospital, Jodhpur (Rajasthan), India

Abstract

Carcinogenesis has a multifactorial etiology. The underlying molecular pathogenesis is still poorly understood. Apart from major risk factors like tobacco, alcohol, infections, there have been some recent and less known factors like endocrine hormones. Sex hormones receptors lie outside sex organs in organs like larynx and lungs where they play a role in gene expression involved in lot of physiological and neoplastic processes. The present article is an attempt to highlight role of various sex hormones in development of carcinogenesis, particularly Head and Neck cancers (HNC).

Keywords: Sex hormones; Carcinogenesis; Head and Neck cancer

Introduction

A multicellular organism can thrive only when all its cells function in accordance with the rules that govern cell growth and reproduction. Why does a normal cell suddenly become a “rebel,” breaking the rules, dividing recklessly, invading other tissues, usurping resources, and in some cases eventually killing the body in which it lives?

Cancer results from a series of molecular events that fundamentally alter the normal properties of cells. In cancer cells the normal control systems that prevent cell overgrowth and the invasion of other tissues are disabled. These altered cells divide and grow in the presence of signals that normally inhibit cell growth; therefore, they no longer require special signals to induce cell growth and division. As these cells grow they develop new characteristics, including changes in cell structure, decreased cell adhesion, and production of new enzymes. These heritable changes allow the cell and its progeny to divide and grow, even in the presence of normal cells that typically inhibit the growth of nearby cells. Such changes allow the cancer cells to spread and invade other tissues [1].

Cancer is one of the biggest threats to humans and animals, claiming 7.6 millions of human lives in 2008 and 13.2 million expected cancer deaths by 2030 [2]. Worldwide, HNC affects more than 550,000 individuals' annually [3]. Oral and oropharyngeal cancer is the sixth most common cancer worldwide. The annual estimated incidence is around 275,000 for oral and 130,300 for pharyngeal cancers [4]. Male to female ratio for cancer predilection ranges from 2:1 to 4:1. The incidence rate in males exceeds 20 per 100,000 in different parts of world including India and United States [5]. Treatment for oral cancer is particularly disabling and disfiguring and disrupts the core aspects of daily life.

Oral cancer describes malignancies of the oral cavity, including structures such as the gingiva, buccal mucosa, hard palate, floor of mouth, salivary glands and anterior two-thirds of the tongue. Cancers of the oropharynx and oral cavity share several risk factors, and the term ‘head and neck cancer’ (excluding nasopharyngeal carcinoma) is commonly used to define cancers of the oral cavity and oropharynx. There are multiple causes that lead to cancer development in an individual. Recent literature suggests that apart from the major established risk factors, female sex hormones may contribute to head and neck carcinogenesis and it strongly suggests certain endocrine involvement in its development [6].

Head and neck cancers are associated in most of the individuals who consume tobacco, arecanut, betel quid and alcohol with history of their long term usage. Latest studies revealed that in some cases of squamous cell carcinoma, a predominant viral etiology has been diagnosed, which were non tobacco and non-smoker group females [6]. Head and neck

carcinomas are also associated with exposure of traditional risk factors in addition to genetic and/or environmental factors [7-9].

Endocrinal hormones levels in both males as well as females plays a significant role in the etio-pathogenesis and progression of cancers of various organs that are associated with the sex hormones receptor like endometrial carcinoma, breast and prostate cancer, lung carcinoma [10]. Young females associated with certain disorders with hormonal imbalance like Polycystic Ovarian Disease (PCOD), have more potential risk in development of Ovarian cancer and endometrial carcinoma of uterus [11].

The gender specific risk for Head and Neck cancer (HNC) has different perspectives. Firstly because of some detrimental factors which affect only male patients. Secondly, there are common risk factors which affect both the genders, but females have some inbuilt defense mechanisms owing to their special metabolic and endocrine features [12]. More than 75% of cases of HNSCC are attributable to smoking and alcohol consumption. Smoking increases the risk by 10-fold compared with never smokers, and heavy alcohol intake is an independent risk factor [13]. The combined effect of tobacco and alcohol causes a greater multiplicative risk. Public health measures have been successful in reducing the use of tobacco, and therefore the incidence of HNSCC overall has been decreasing over the past 30 years in developed countries. However, there has been a dramatic increase in the incidence rates of oro-pharyngeal (tonsil and base of tongue) cancers because of infection with high-risk Human Papillomavirus (HPV) [14].

Physiology of Sex Hormones: A Brief Overview

A hormone (from Greek ὁρμή, “impetus”) is a class of signaling molecules produced by glands in multicellular organisms that are transported by the circulatory system to target distant organs to regulate physiology and behavior [13].

Important Sex hormones of human body are estrogen, progesterone and testosterone, belonging to a group of steroid hormones, all of which are derivatives of cholesterol.

***Corresponding author:** Ankita Bohra, Department of Oral Medicine Diagnosis & Radiology, Vyas Dental College & Hospital, Jodhpur (Rajasthan), India, Tel: 9414454413; E-mail: aav1423@hotmail.com

Received November 14, 2014; Accepted January 19, 2015; Published January 26, 2015

Citation: Bohra A, Bhateja S (2015) Carcinogenesis and Sex Hormones: A Review. *Endocrinol Metab Syndr* 4: 156. doi:[10.4172/2161-1017.1000156](https://doi.org/10.4172/2161-1017.1000156)

Copyright: © 2015 Bohra A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Estrogen

Estrogens (ES) are female sex hormones, synthesized in the ovary, placenta, testes and adrenal cortex, contribute to the development and growth of female genital gender and female sex characteristics, and cause endometrial growth. Estrogens are considered to be one of the etiological factors of breast, uterine and ovarian tumors. It is also believed that they have an impact on other disorders, such as thrombo-embolic disease [14].

Types of estrogen:

- 1) Natural
- 2) Synthetic.

There are three basic, biologically active estrogens: Estrone (E1), 17-β Estradiol (E2) and Estriol (E3). Out of them main estrogen with highest biological activity is 17-β estradiol (E2) [15,16].

Testosterone

It is the principal male sex hormone, belong to androgen group and been secreted by testis of males and ovaries in females.

Progesterone

An endogenous steroid hormone involved in the menstrual cycle, pregnancy and embryogenesis of humans. It belongs to a group of steroid hormones called the progestogens and is the major progestogen in the body, plays an important role in brain function as a neurosteroid [17,18].

Effect of sex hormones on cancer development

The mechanism by which sex hormones affect cancer risk is probably largely (Figure 1) through determining the number and mitotic rate of the epithelial cells in the organ concerned [19]. High mitotic rates can increase cancer risk by increasing the chance of mutations occurring and of being replicated before they are repaired, and can also increase the growth of early tumours. In the case of estrogens, it has also been argued that certain metabolites of oestradiol may cause mutations by directly damaging the DNA, but the importance of this possible mechanism has not been established, and we think that the major role of sex hormones in cancer is likely to be as determinants of cell number and mitotic rate [20].

Androgen, estrogen and progesterone, acting through specific receptors, play an important role in the growth and development of several tumors, including breast, endometrium, and prostate carcinomas [21,22]. Additionally, patients with breast carcinomas positive to Estrogen Receptor (ER) and Progesterone Receptor (PgR) have been treated with antagonist hormones and shown decreased recurrences and higher survival rates. Similar results have been reported in patients with prostate cancer positive to Androgen Receptor (AR) [23,24].

Recently studies demonstrated that elevated prolactin levels in HNC can be a marker of poor prognosis [25]. Few other studies have demonstrated increased levels of FSH, LH, prolactin and decreased ratio of testosterone and estradiol in tongue cancer patients. These hormonal fluctuations clearly indicated towards a disruption in the pituitary-adrenal-testicular axis. Thereof, it is suggested that these hormones might play a crucial role in the development and progression of oral cancer [26].

It was also reported that estrogen stimulates the proliferation and maturation of gingival connective tissue, epithelium and salivary glands

[27]. ERs have been identified in a variety of human tumors rather than breast carcinomas using histochemical, immunohistochemical, and molecular biology techniques. Moreover, the expression of sex hormone receptors in certain tumors suggests a role for these receptors in tumor pathogenesis, progression and therapy. There is also direct evidence that the risk for some of these cancers is related to the circulating serum concentration of various sex hormones [28].

Direct influence of estrogen on target cells is caused by combining the steroid receptor complex with DNA. In that way, it can regulate the expression of genes and transform proto-oncogenes into oncogenes. Estrogens have two kinds of receptors – ERα and ERβ. They belong to the nuclear receptor superfamily of transcriptional activators. The affinity of receptors is different to different ligands, e.g. estradiol has a higher affinity for ERα than ERβ. Obesity causes an increase in serum concentrations of bio-available oestradiol, and this factor causes increases in the risk for both endometrial cancer and breast cancer [29].

Effect on head and neck cancers

Some recent reports have shown substantive evidences that certain salivary gland tumours are similar to breast cancer at cellular and molecular level as well [30]. Both the tumors show similar expression of progesterone associated with tumor onset and progression [31].

Estrogen may increase the movement of precancerous cells in the mouth and promote the spread of head and neck cancers. The results may help researchers to understand the risk factors that cause head and neck cancers in addition to the traditional risk factors of tobacco and alcohol exposure [32]. Squamous cell carcinoma of the head and neck (HNSCC), which include cancers of the tongue, mouth and throat, represent the sixth most common type of cancer and is known to be on the increase in some demographic groups, including young women without known risk factors. A recent report showed that 75% of young never-smoker/never drinker HNSCC patients who develop primary oral tongue squamous cell carcinoma are women, suggesting that female hormones may contribute to carcinogenesis [33].

Oestrogen has a strong impact on precancerous and cancerous cells. An earlier study showed that the oestrogen metabolism pathway is altered in lung tissue following tobacco smoke exposure, suggesting that oestrogen metabolism may play a role in the formation of other cancers of the aerodigestive tract [34].

Using cancer cells grown in the laboratory, the team found that oestrogen induced the expression of an enzyme called cytochrome P450 1B1 (CYP1B1) which is responsible for breaking down toxins and metabolising oestrogen. The CYP1B1 induction only occurred in precancerous cells and not in healthy cells or cells that had become cancerous [35,36].

Furthermore, investigators found that depleting the expression of CYP1B1 reduced the ability of precancerous cells to move and divide,




Figure 1: Mechanisms involved in carcinogenesis through sex hormones

in comparison with similar cells containing normal levels of CYP1B1. Oestrogen also reduced cell death in the precancerous cells, irrespective of the amount of CYP1B1 present.

Finally immune-histochemical staining of samples from patients with head and neck cancers showed that 91.9% stained positive for oestrogen receptor β , 99.4% for CYP1B1 and 88.4% for 17 β oestradiol.

“This study is the first to report the detection of estrogen within human head and neck tissue and demonstrate that both estrogen and CYP1B1 may contribute to the progression of HNSCCs,” concluded the authors. They add that CYP1B1 levels may offer an important biomarker of tumorigenesis in head and neck cancers and present a novel target for chemo preventive interventions in patients with premalignant lesions [37,38]. Androgen is known to have an important role in the normal development and differentiation of a variety of cell and tissue types. This function is mediated by its binding to Androgen Receptor (AR), a member of the family of steroid hormone receptors. The presence of AR in a given cell type or organ system indicates a possible role for androgen in its growth and differentiation. AR mediates the effect of androgen by binding to specific DNA sequences and influences the transcription and translation of various genes. Fan et al suggested that AR may have a role in the pathogenesis of salivary duct carcinoma through the mediation of an epidermal growth factor receptor and transforming growth factor- α autocrine pathway similar to that seen in prostatic carcinoma [38].

Experimental studies have shown that androgen influences the expression of proto-oncogenes (like *c-myc*) and apoptotic factors (like the *bcl-2* family) in lacrimal, salivary, and prostatic tissues of both mice and rats, as well as in cell line models. Whether the expression of AR in malignant salivary gland tumors indicates a role for androgen in the pathogenic process or simply represents an epiphenomenon of the malignant transformation remains to be determined.

Although partial remission of a salivary gland carcinoma following goserelin (an antiandrogen) therapy has been reported, this needs to be confirmed in a larger group of patients. It is been clearly demonstrated that there is a strong and consistent expression of AR exclusively in a sharply defined subset of malignant salivary gland tumors [39,40].

Clinical implications of head and neck cancer

Benefits of endocrine therapy in breast cancer by targeting sex steroid hormone receptor, its potential role in HNC are also tried and results of completed clinical trials are eagerly awaited [41]. Recent study conducted on tongue carcinomas demonstrated that ER antagonist interferes with cell adhesion and ultimately results in cell death which further prevents growth and progression of tumor [42]. Treatment with ER antagonists such as tamoxifen is shown to decrease the phosphorylation of focal adhesion kinase (FAK), leading to reduced phosphorylation of extracellular signal-related kinase (Erk) and mitogen-activated protein (MAP) kinase which consequently disrupts tumour growth [39].

Further studies at molecular level and larger clinical trials are need of the hour to determine which and when receptors should be targeted by means of which modulators in management of HNC patients. Since, hormonal therapies may also have some adverse effects on physiological processes, so balance has to be maintained between therapeutic benefit of such interventions and potential adverse effects [43]. Genome sequencing in association with proteomics could be helpful to define and select useful genetic and molecular biomarkers to predict and maintain progression of HNC in nearly future. Identification of specific genetic, epigenetic and metabolic disturbances, in unison with other

conventional techniques in diagnosis and prognostication, are required to make effective treatment strategy [44].

Conclusion

Role of sex hormones is well established in cases of prostate, breast and endometrial carcinomas. Though the role of sex hormones in HNC is controversial and is debatable, so the present article is an attempt to highlight its impending role in HNC so that future studies can be performed on this less known entity.

References

- Venu Gopal Y, Ravindranath A, Kalpana G, Sreenivas SA, Raj Kapoor B (2012) Antitumor and antioxidant activity of *Diospyros peregrina* against Dalton's ascites lymphoma in rodents. *Annals of Biological Research* 3: 4985-4992.
- Muhammed IB, Alfred C, Keshaw PT (2013) A retrospective Study of Canine Tumors in Grenada, West Indies. *Adv Anim Vet Sci* 1: 134-139.
- Parkin DM, Bray F, Ferlay J, Pisani P (2005) Global cancer statistics, 2002. *CA Cancer J Clin* 55: 74.
- Warnakulasuriya S (2009) Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol* 45: 309-316.
- Sankaranarayanan R, Masuyer E, Swaminathan R, Ferlay J, Whelan S (1998) Head and neck cancer: a global perspective on epidemiology and prognosis. *Anticancer Res* 18: 4779.
- Mannarini L, Kratochvil V (2009) Human Papilloma virus in Head & neck region: A review of literature. *Acta Otorhinolaryngol Ital* 29: 119-126.
- Koch WM, Lango M, Sewell D, Zahurak M, Sidransky D (1999) Head and neck cancer in nonsmokers: a distinct clinical and molecular entity. *The Laryngoscope* 109: 1544-1551.
- Divaris K, Olshan AF, Smith J, Bell ME, Weissler MC, et al. (2010) Oral health and risk for head and neck squamous cell carcinoma: the Carolina Head and Neck Cancer Study. *Cancer Causes Control* 21: 567-575.
- Harris SL, Kimple RJ, Hayes DN, Couch ME, Rosenman JG (2010) Never-smokers, never-drinkers: unique clinical subgroup of young patients with head and neck squamous cell cancers. *Head & Neck* 32: 499-503.
- Lukits, Remenár E, Rásó E, Ladányi A, Kásler M, et al. (2007) Molecular identification, expression and prognostic role of estrogen- and progesterone receptors in head and neck cancer. *Int J Oncol* 30: 155-60.
- Badawy A, Elnashar A (2011) Treatment options for polycystic ovary syndrome. *International Journal of Women's Health* 3: 25-35.
- Suba Z (2007) Gender-Related Hormonal Risk Factors for Oral Cancer. *Pathology Oncology Research* 213: 195-202.
- Chaturvedi AK, Anderson WF, Lortet-Tieulent J, Curado MP, Ferlay J, et al. (2013) Worldwide Trends in incidence rates for oral cavity and oropharyngeal cancers. *J Clin Oncol* 31: 4550-4559.
- Chaturvedi AK, Engels EA, Anderson WF, Gillison ML (2008) Incidence trends for human papillomavirus-related and -unrelated oral squamous cell carcinomas in the United States. *J Clin Oncol* 26: 612-619.
- Sanghi DK, Rakesh Tiwle (2014) An elaborate review on hormone & its importance. *International journal of pharmacological screening* 4: 56-68.
- Marta, Halina (2012) The Effect of Using Estrogens in the Light of Scientific Research. *Adv Clin Exp Med* 21: 535-543.
- Sunhee Yu, Minhyung Lee, Seungjin Shin (2011) Apoptosis Induced by Progesterone in Human Ovarian Cancer Cell Line SNU-840. *Journal of Cellular Biochemistry* 82: 445-451.
- Bauernhofer T, Pichler M, Wiecekowsk E, Stanson J, Aigelsreiter A, et al. (2001) Prolactin receptor is a negative prognostic factor in patients with squamous cell carcinoma of the head and neck. *British J Cancer* 104: 1641-1648.
- Bhatavdekar JM, Patel, Vora HH, Balar DB (1994) Circulating hormones in patients with tongue cancer. *Oncology Reports* 1: 463-464.
- Parkar MH, Newman HN, Olsen I (1982) Polymerase chain reaction analysis of oestrogen androgen receptor expression in human gingival and periodontal tissue. *Archs oral Biol* 41: 979-983.

21. Glas AS, Hollema H, Nap RE, Plukker JT (2002) Expression of estrogen receptor, progesterone receptor, and insulin-like growth factor receptor-1 and of MIB-1 in patients with recurrent pleomorphic adenoma of the parotid gland. *Cancer* 94: 2211-2216.
22. Fan CY, Malhelm MF (2001) Expression of androgen receptors, epidermal growth factor receptor & transforming growth factor alpha in salivary duct carcinoma. *Arch Otolaryngol Head Neck Surg* 127: 1095-1097.
23. Linson EM, Peter J (2005) Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet* 365: 1687-1717.
24. Berthelet E, Pickles T, Lee KW, Liu M, Truong PT (2005) Long-term androgen deprivation therapy improves survival in prostate cancer patients presenting with prostate-specific antigen levels > 20 ng/mL. *Int J Radiat Oncol Biol Phys* 63: 781-787.
25. Ciocca DR, Puy LA, Fasoli LC (1989) Study of estrogen receptor, progesterone receptor, and the estrogen-regulated Mr 24,000 protein in patients with carcinomas of the endometrium and cervix. *Cancer Res* 49: 4298-4304.
26. Hye-Rim Lee, Tae-Hee Kim (2012) Functions and physiological roles of two types of estrogen receptors, ER α and ER β , identified by estrogen receptor. *Lab Anim Res* 28: 71-76.
27. Lieghr G (2001) Genotoxicity of the steroidal oestrogens oestrone and oestradiol: possible mechanism of uterine and mammary cancer development. *Hum reported update* 7: 273-281.
28. Samuel MC and Leon BE (1993) Use of Cell Proliferation Data in Modeling Urinary Bladder Carcinogenesis. *Environmental Health Perspectives* 101: 111-114.
29. Murase R, Sumida T, Ishikawa A, Murase R, McAllister SD, et al. (2011) Novel therapeutic strategies for malignant salivary gland tumors: lessons learned from breast cancer. *Int J Otolaryngol* 2011: 187623.
30. Stauber RH, Wunsch D, Knauer SK, Fetz V (2010) An update on the pathobiological relevance of nuclear receptors for cancers of the head and neck. *Histol Histopathol* 25: 1093-1104.
31. Timothy JK, Naomi EA, Pia KV (2001) Macronutrient Metabolism Group Symposium on 'Energy flux and cancer', Energy balance and cancer: the role of sex hormones. *Proceedings of the Nutrition Society* 60: 81-89.
32. Remenár É, Számel I, Buda B (1997) 'Why men?' Hormones and hormone receptors in male head and neck cancer patients. In: *Advances in Laryngology in Europe*. Klainsasser O, Glanz H and Olofsson J (eds). Elsevier, 137-140.
33. Colella G, Izzo G, Carinci F, Campisi G, Lo Muzio L, et al. (2011) Expression of sexual hormones receptors in oral squamous cell carcinoma. *Int J Immunopathol Pharmacol* 24: 129-132.
34. S. Pradhan, Usha Singh (2007) Estrogen and progesterone receptor status in breast cancer: Effect of oral contraceptive pills and hormone replacement therapy. *The Breast* 16: 540-545.
35. Chin D, Boyle GM, Porceddu S (2006) Head and neck cancer: past, present and future. *Expert Rev Anticancer Ther* 6: 1111-1118.
36. Shoker BS, Jarvis C, Sibson DR (1999) Estrogen receptor expression in the normal and pre-cancerous breast. *J Pathol* 188: 237-244.
37. Ekaterina GS, Andres JP, Klein-Szanto, Karthik Devarajan (2011) Estrogen and Cytochrome P450 1B1 Contribute to Both Early- and Late-Stage Head and Neck Carcinogenesis. *Cancer Prev Res* 4: 107-115.
38. McEwan IJ (2004) Molecular mechanisms of androgen receptor-mediated gene regulation: structure-function analysis of the AF-1 domain. *Endocrine-Related Cancer* 11: 281-293.
39. Nasser SM, Faquin WC, Dayal Y (2003) Expression of androgen, estrogen, and progesterone receptors in salivary gland tumors. Frequent expression of androgen receptor in a subset of malignant salivary gland tumors. *Am J Clin Pathol* 119: 801-806.
40. Falk RT, Brinton LA, Dorgan JF, Fuhrman BJ, Veenstra TD, et al. (2013) Relationship of serum estrogens and estrogen metabolites to postmenopausal breast cancer risk: a nested case-control study. *Breast Cancer Res* 15: R34.
41. Hamakawa H, Nakashiro K, Sumida T, Shintani S, Myers JN, et al. (2008) Basic evidence of molecular targeted therapy for oral cancer and salivary gland cancer. *Head Neck* 30: 800-809.
42. Williams MD, Roberts D, Blumenschein GR (2007) Differential expression of hormonal and growth factor receptors in salivary duct carcinomas: biologic significance and potential role in therapeutic stratification of patients. *Am J Surg Pathol* 31: 1645-1652.
43. Leeuw R, Neeffjes J, Michalides R (2011) A Role for Estrogen Receptor Phosphorylation in the Resistance to Tamoxifen. *Int J Breast Cancer* 2011: 232435.
44. Kumar RJ, Barqawi A, Crawford ED (2005) Adverse Events Associated with Hormonal Therapy for Prostate Cancer. *Rev Urol* 5: S37-43.

Citation: Bohra A, Bhateja S (2015) Carcinogenesis and Sex Hormones: A Review. *Endocrinol Metab Synd* 4: 156. doi:10.4172/2161-1017.1000156

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

