

Chemical Approach to Signal Transduction by Inositol Triphosphate

Shoichiro Ozaki*

The Institute of Physical and Chemical Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

Abstract


Berridge discovered that inositol 1,4,5-trisphosphate (IP₃) was generated at the cell surface in response to cell stimulation and functioned as a second messenger to release Ca²⁺ from internal stores. Ozaki et al. succeeded in the first total synthesis of optically active IP₃ by 13 steps. He supported the signal transduction studies by supplying necessary reagents such as IP₃, other IP_x, phosphatidylinositol, new synthetic methods and reagents. He discovered the regulators of Ca²⁺ release and consequent cellular processes.

Keywords: Signal transduction; Inositol triphosphate; IP₃Phosphatidylinositol; Regulator of cellular process

Introduction

The fact that diacyl glycerol is second messenger was found by late Professor Yasutomi Nishizuka [1] and the fact that Inositol triphosphate (IP₃) is a second messenger was discovered by Michael Berridge who showed that it functioned to release Ca²⁺ from internal stores. This bifurcating signaling system is of fundamental importance in regulating a wide range of cellular process.

Signals (first messenger) like light, noise, taste, odor, hormone, neurotransmitter, drug attach to the plasma membrane where they are recognized by cell surface receptors. Upon binding of the ligand to the appropriate receptor, activation of G protein activates in turn phospholipase C. Active phospholipase C hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP₂) giving rise to two products: 1,2-diacylglycerol and inositol 1,4,5-trisphosphate (IP₃). IP₃ stimulates the release of Ca²⁺ from the intracellular stores in the endoplasmic reticulum through IP₃ receptor while regulating a wide range of cellular processes.



Why Plant Biosynthesize Inositol

The rice bran, wheat, corn contain much phytic acid (inositol hexaphosphate) as Ca salt. Plant make glucose by photo synthesis from carbon dioxide and water. Some of glucose is converted to inositol. Inositol is converted to phospholipids (PIP₂) and phytic acid. PIP₂ is converted to IP₃ and diacylglycerol. These two compounds are essential for signal transduction of plant. Plant makes phytic acid as storage of phosphorous. Phosphorous is an essential atom as fertilizer because it is an essential atom to make nucleic acid, DNA. The seed store phosphorous atom as a store so that even when seed germinate at no phosphorous land [1].

Discovery of IP₃

Phospholipid was discovered by Bollow in 1961 [2] from bovine brain. The hypothesis of Michell [3] that the receptor controlled hydrolysis of phosphoinositides could be directly linked to cellular calcium mobilization. The observation by Berridge D-myo-inositol 1,4,5-trisphosphate (IP₃) act as a second messenger, a fundamental cell-signal transduction mechanism has been elucidated. IP₃ stimulates the release of Ca²⁺ from the intracellular stores in the endoplasmic reticulum through IP₃ receptor while regulating a wide range of cellular processes [4-25].

Synthetic Competition of Inositol P

The discovery of inositol phosphate in particular IP₃ led to the dramatic stimulation for the synthesis of inositol phosphates. Many persons challenged the synthesis of inositol phosphate, starting from inositol, glucuro lactone, phytic acid, arenas, quinic acid and L-quebrachitol.

A symposium; Inositol phosphates and Derivatives. Synthesis, biochemistry, and therapeutic potential was held by the division of carbohydrate Chemistry at the 200th National meeting of the American Chemical Society, Washington DC, August 26-31.1990. ACS Symposium Series. 463 Edited by Allen B.Reitz was published.

The key problems in the synthesis of inositol phosphates are (1) synthesis and optical resolution of suitably protected inositol derivatives, (2) efficient phosphorylation of vicinal hydroxy groups.

In 1986, Ozaki et al succeeded in the first total synthesis of optically active myo-inositol tris (1,4,5) phosphate from myo-inositol by 13 steps [26]. At this report, phosphorylation yield of 2,3,6-tribenzyl myo-inositol by dianilidephophotyl chloride isolation yield was only 10%. Then we have studied phosphorylation reagents and discovered new phosphorylation method Then we could get IP₃ by best method in good

*Corresponding author: Shoichiro Ozaki, The Institute of Physical and Chemical Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan, Tel/ Fax: 81-0467670991; E-mail: ozaki-0991@m.jcnnet.jp

Received September 15, 2014; Accepted September 24, 2014; Published October 04, 2014

Citation: Ozaki S (2014) Chemical Approach to Signal Transduction by Inositol Triphosphate. J Bioengineer & Biomedical Sci 4: 133. doi: 10.4172/2155-9538.1000133

Copyright: © 2014 Ozaki S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

yield as shown in Figure 1 [26,27].

This IP₃ is produced by this method at DOJINDO (Kumamoto, Japan) and is distributed all over the world by the name of synthetic IP₃.

The synthesis of I(1,4,5)P₃ are reported by many investigators Billington and Vacca (from myo-inositol orthoformate [28], Ballou, from myo-inositol [29], Gigg [30], Ley from arenes using Pseudomonasoxidation [31], Falch from Quinic acid [32], Stepanov and Shvets [33], Prestwich prepared D-myo-(³H)I(1,4,5)P₃, essential and most used reagent for the study of signal transduction [34,35] Phosphothioate analog of IP₃ by Potter [36].

Different Source and Methods

IP₃ was obtained through 6 different sources:


- Starting from myo -inositol [26-32,35-38]
- Starting from L-quebrachitol (natural rubber industry by-product) [39-43]
- Starting from arenes [33]
- Starting from Quinic acid [34]
- Starting from D-glucuronolactone [42]
- Chemoenzymtic synthesis of D-myo-inositol 1,4,5—triphosphate [43-49]

Methods to Get Optically Pure Compound by 5 Different Methods

- › Separation of diastereomers
 - L- mentoxy acetyl chloride gave best result, because desired product was crystal [26]
- › Starting from optically pure natural product
 - From D-glucuronolactone [42], From Quinic acid [34]
 - From L-quebrachitol [39-43]
- › Use of tartaric acid ester [50]
- › Use of enzyme (like Phosphorylase). Enzyme aided synthesis of D-myo-inositol 1,4,5-triphosphate [43-48]
- › Enzymic resolution of racemic 1,2:5,6-di-O-cyclohexylidene and 1,2:3,4-di-O-cyclohexylidene-myo-inositol [45]
- › Enzymic resolution of sterically hindered myo-inositol derivatives [48]
- › Enzyme aided regioselective acylation of nucleosides [49]

Phosphorylation Reagents [51]


- Tetrabenzyl pyrophosphate (TBPP) and n-BuLi [52].




TBPP
n-BuLi

- New phosphorylating reagent called OXDEP (o-xylylene N,N-diethylphosphoramidite) [53,54]. By using this reagent, IP₃ and PIPx

were obtained in good yield [27].



- **DBPF** Dibenzyl phosphorofluoridate (BnO)₂P(O)F [55]. This reagent was used for the synthesis of phosphofluoridate analogues. Obtained phosphofluoridates showed very interesting biological activity [56].




- Step wise phosphorylation using PCl₃, BnOH, C₆H₅COOH [57]
- Phosphorothioate synthesis based on the redox reaction of phosphite with tellurium (IV) chloride [58].


Discovery of Phosphonium Salt Methodology

This phosphonium salt methodology [59,60] provide a regioselective phosphorylation. 1,2-Diol were phosphorylated regioselectively at C-1 with tribenzyl phosphite to give 1-dibenzyl phosphate 2-hydroxy free compound as shown in Figure 2. Other phosphorylating reagents do not have such selectivity. By using this free hydroxy group, we could get 2-acyl analog and IPx and PIPx. Three kind of combined reagents are possible.


Trialkyl phosphite and pyridinium bromide perbromide method [59]



1H-Tetrazol catalyzed the reaction of trialkyl phosphite [61]



Utilization of oxidizing character of TeCl₄ [62]



The reaction of an alcohol with a trialkylphosphite in the presence of pyridinium bromide per bromide proceed via the phosphonium salt (RO)₃P⁺Br⁻ to afford the triester R₁-OP(O)(OR)₂, which can be converted to the phosphoric monoester by deprotection.

On the other hand, starting from dialkyl phosphoramidite (RO)₂PNR, the corresponding triester product (R₁O-R₂O-P(O)OR³) gave phosphoric mixed diesters.

The reactivity of phosphonium salt toward an alcohol seems to be between P^{III} and P^V, therefore we expected that the phosphonium salt methodology would provide a regioselective phosphorylation method. 1,2-Diol was phosphorylated regioselectively.

Applying the phosphonium salt approach to the synthesis of phosphoinositides, the use of glyceryl phosphite, which was derived by the reaction of the glycerol derivatives with dimethylphosphoramidite in the presence of tetrazol, gave the protected PI(4,5)P₂.

We synthesized Phosphatidylinositol 3,4,5-triphosphate [63,64], Unsaturated phosphatidyl inositol-3,4,5-triphosphate [65], myo-inositol 1,2,5,6-tetrakisphosphate [66], 4-α-D-glucopyranosyl-myo-inositol, enzymic transglycosylation product. [67], 2,6-Di-O-(D-mannopyranosyl)phosphatidyl-D-myo-inositol [68], Phosphofluoridate analogs of myo-inositol 1,4,5-tris(phosphate) [56].




Figure 1: Synthesis of D—myo-inositol 1,4,5-trisphosphate

Finding of New Reaction, New Methods and New Reagents

Finding of new protection methods

- Protection by tetraisopropylidisiloxane-1,3-diyl group [69]
- Proximally assisted and chemoselectively cleavable protecting groups for alcohols, 2-[2-(arylmethoxy)ethyl]benzoic esters [70].

Finding of new deprotection methods

- Deprotection of methyl group by AlCl₃-NaI, AlCl₃-Bu₄NI [71]
- Deprotection of benzyl and allyl group by AlCl₃-dimethyl aniline [72]
- Deprotection p-methoxybenzyl by trimethylsilylchloride-tin(II) chloride—anisole [73].

Finding of diastereoselective addition methods

Diastereoselective addition of organometallics to α-keto esters [74].

Finding of diastereoselective reduction methods

Diastereoselective reduction of ketoester bearing chiro-inositol as chiral auxiliaries [75].

Finding of novel deacylation Methods

A Grignard reagent was used for deacylation without affecting the neighboring base-sensitive functional groups [76].

Finding of novel enantioselective acylation and deacylation Method

Enantioselective acylation and deacylation method using enzyme [38,77].

Finding of glycosidation method

Glycosidation based on phosphite chemistry [78,79]

Phosphorylation of inositol 1,4,5-trisphosphate analogs by 3-kinase and dephosphorylation of inositol 1,3,4,5-tetrakisphosphate analogs by 5-phosphatase [80].

Use of Inositol Derivatives as Chiral Auxiliaries

- Diastereoselective addition of organometallics to keto esters [74].
- Asymmetric synthesis of tetrahydrofurans by diastereoselective (3+2) cycloaddition of allylsilanes with ketoesters bearing optically active cyclitol as a chiral auxiliary [81].
- Preparation of optically active D 2-isooxazolines via addition




Figure 2: Phosphosponium salt methodology for the synthesis of IP_x and PIP_x.




Figure 3: Preparation of ins (1,4,5)P₃ analogues

of nitril oxides to chiral acryloxy esters bearing cyclitols as auxiliaries [82].

- Asymmetric synthesis of functionalized tertiary alcohols by the diastereoselective aldol reaction [83].

Preparation of IP_x, IP₃ derivatives and IP₃ Analog, and Assessment of their Activities

Synthesized inositol poly phosphate [66,84-86]

Myo-inositol 1-phosphate [87,88], myo-inositol

1,3,4-trisphosphate [89], myo-inositol 1,4,6-trisphosphate [90], myo-inositol 1,3,4,5-tetrakisphosphate [91-93], myo-inositol 1,4,5,6-tetrakisphosphate [94], myo-inositol 2,4,5-trisphosphate [57], 1,2-cyclic-4,5-, 1,4,5- and 2,4,5-triphosphate [95], myo-inositol 1,2,5,6-tetrakisphosphate [65], 2,6-Di-O-(D-mannopyranosyl) phosphatidyl-D-myo-inositol [68], Phosphofluoridate analogs of myo-inositol 1,4,5-tris(phosphate) [55]. 4-a-D-glucopyranosyl-myo-inositol [50].

2-substituted IP₃ analogs

- These were synthesized as shown in Figure 3. These analogs were used for the preparation of affinity columns [96].
- Many IP_n and derivatives were prepared and their activities were measured by Prof. Hirata, Masato [96-113].
- Synthesis of IP₃ having biotinyl and azidobenzoyl groups [100].
- Synthesis of 2-substituted myo-inositol 1,3,4,5-tetrakis(phosphate) and 1,3,4,5,6-pentakis(phosphate) analogues [101].

Phosphofluoridate analogues

Phosphofluoridate analogs of myo-inositol 1,4,5-tris(phosphate) were prepared as shown in Figure 4 [56].

The three phosphofluoridates thus prepared had potencies for inhibiting (³H) InsP₃ binding to purified InsP₃ receptor that were less than for InsP₃. Two analogues 44 and 40 were found to inhibit the dephosphorylation of (³H) InsP₃ by the 5-phosphatase with potencies similar to that for InsP₃. Surprisingly, the inhibitory potency of 5-phosphofluoridate 44 toward 5-phosphatase was higher (about 20 fold) than those of InsP₃ and the another fluoridates 40 and 45.

Preparation of Affinity Column

Inositol 1,4,5-trisphosphate affinity columns 24, 25 were prepared from 20, 21 as shown in Figure 5 to fish out IP₃-binding proteins [98,113,114].

Isolation and Characterization of Many IP₃-Binding Proteins

The following many proteins were isolated by affinity column and characterizations were carried out [103,111].

- IP₃ binding protein [102,103,113,115] co-work with Hirata Masato
- Phospholipase C-d1 [110,116] co-work with Hirata Masato.
- Porcine tracheal smooth muscle aldase [109], collaboration with Carl Baron and Masato Hirata.
- 3-Kinase, 5-phosphatase [111,117] collaboration with Van Dijken
- Growth factor activating protein [112,118] collaboration with Moriya Shigeharu
- IP₃ 3-kinase from porcine smooth muscle [119-121] co-work with Denborough.
- RAC-protein kinase (PKB/Akt [120] collaboration with Matsuzaki
- Expression and characterization of IP₃-binding domain of phosphatidyl inositol-specific phospholipase C [122] collaboration with Yagisawa, Hitoshi.
- IP₃ 3-kinase from chicken erythrocytes [123] collaboration with George Myer.






Figure 4: Synthesis of Phosphofluoridate analogues.



- Homogeneous Ca²⁺ Stores [124] collaboration with Inoue, Masumi
- Na⁺,K⁺-ATPase [125] collaboration with Lin, Hai
- Plasma membrane PtdIns-4,5-P₂ [126] collaboration with Ronald Holz

Observation of Behaviors of IP_x, PIP_x and Ca²⁺ Flux in the cells

IP₃ and PIP_x are charged compounds. Therefore they do not readily penetrate through cell membranes. But it becomes permeable following salt formation with amine and this is a new method to put IP_n or PIP_n into the cell [127-130].

Ozaki synthesized about 20 fluorescent IP₃, IP₄, PIP, PIP₂ containing fluorescent amines with green and red colors (Figure 6), and introduced these tagged molecules into cells. He then used fluorescent microscopy.

He observed how and how fast the IP_n or PIP_n entered into the cell and how moved and how changed, metabolized and also observed a calcium flux (time, location, concentration) in NIH 3T3 Fibroblasts, when complexes of carrier and Ptd Ins (4,5) or Inos (1,4,5) P₃ were added extra cellular. He took more than a thousand pictures and movies.

Detection of Ca²⁺ Flux

Fluoro-3 was used to measure intracellular calcium concentration. In case of IP₃ complex, Ca²⁺ maximum peak (2.5×10⁻⁷M) was observed after 3.5 min. after addition of IP₃. In case of PIP_x complex, Ca²⁺ maximum peak (4.2×10⁻⁷M) was observed after 6.5 min.


Discovery of DAB; Regulators of Ca²⁺ Release and Cellular Response

In 1997, we identified 2-aminoethyl diphenylborinate (2-APB) as being an IP₃ receptor inhibitor and regulate IP₃ induced calcium release [131,132]. This discovery rose a substantial interest and had a great impact as it gained more than 600 citations and more than 1000 studies on 2-APB have been published so far. This was supported by increasing sales of 2-APB by Sigma-Aldrich as membrane-permeable modulator of calcium release. We aimed at generate better modulator of calcium release than 2-APB.

We synthesizes several 2APB analogues and measured their inhibitory activities on Store Operated Calcium Entry (SOCE) and IP₃ Induced Calcium Release (IICR).

We found that bis boron compound DBP 161 and DBP 163 were 10 times more effective than 2-APB [133-138] We extended these studies and synthesized 493 analogues [139,140] increasing the number of borons, changing diphenyl to diaryl, monoaryl, mono-aliphatic dialiphatic compounds, substitution of aminoethyl to amino acid derivative as well as aminoethanol to aminoethylthiol and studied the structure/activities correlation.

We found that Diphenyl (amino acidonate O,N) borane DAB are best compounds



We found [139,140] that compounds DAB Diphenyl (amino acidonate N,O)borane could regulate IP₃- induced Ca²⁺ release (IICR), Store-Operated Ca²⁺ entry (SOCE)) and could regulate cellular responses. We found that the adduct of amino acid (especially basic amino acid) and diphenyl borinic acid have strong inhibitory activity to SOCE. And some of them 919 Diphenyl (2,3-diaminopropionate O,N) borane, 911 Diphenyl (L- lysinate O,N) borane showed 10 times strong activity than 2-APB. 2APB is said to be a excellent lead compound for heat disease and Alzheimer`s diseases as Berridge predicts [141-147].

2APB analogues presented in this study could be proven to be excellent lead compounds for many human diseases including heart disease [143,144], Alzheimer`s [145-146] and Huntington disease [148,149].

We found that boron compounds also can inhibit transglutaminase (Ca²⁺-dependent enzyme) [130]. There are many neurodegenerative disease, including Alzheimer`s disease, Huntington`s disease [136,149]. The boron compounds were found to be effective as inhibitor of acyl protein thioesterase [150].

We looked for more effective transglutaminase inhibitors. We synthesized 250 β- aminoethyl ketones and found that these compounds had strong transglutaminase inhibitory activities [151,152]. A typical compound is 5-bromo-2-thienyl-(N-t-butyl-N-benzyl)-aminoethyl ketone.

Acknowledgement

I would like to thank Dr. M. J. Berridge for valuable suggestions and advices.

References

- Nishizuka Y (1984) The role of protein kinase C in cell surface signal transduction and tumour promotion. *Nature* 308: 693-698.
- Frezza C (2014) The role of mitochondria in the oncogenic signal transduction. *Int J Biochem Cell Biol* 48: 11-17.
- Michell RH (1975) Inositol phospholipids and cell surface receptor function. *Biochim Biophys Acta* 415: 81-47.
- Fain JN, Berridge MJ (1979) Relationship between hormonal activation of phosphatidylinositol hydrolysis, fluid secretion and calcium flux in the blowfly salivary gland. *Biochem J* 178: 45-58.
- Fain JN, Berridge MJ (1979) Relationship between phosphatidylinositol synthesis and recovery of 5-hydroxytryptamine-responsive Ca²⁺ flux in blowfly salivary glands. *Biochem J* 180: 655-661.
- Berridge MJ, Downes CP, Hanley MR (1982) Lithium amplifies agonist-dependent phosphatidylinositol responses in brain and salivary glands. *Biochem J* 206: 587-595.
- Berridge MJ, Dawson RMC, Downes CP, Heslop JP, Irvine RF (1983) Changes in the levels of inositol phosphates after agonist-dependent hydrolysis of membrane phosphoinositides. *Biochem J* 212: 473-482.
- Berridge MJ (1983) Rapid accumulation of inositol trisphosphate reveals that agonists hydrolyse polyphosphoinositides instead of phosphatidylinositol. *Biochem J* 212: 849-858.
- Streb H, Irvine RF, Berridge MJ, Schulz I (1983) Release of Ca²⁺ from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* 306: 67-69.
- Berridge MJ, Heslop JP, Irvine RF, Brown KD (1984) Inositol trisphosphate formation and calcium mobilization in Swiss 3T3 cells in response to platelet-derived growth factor. *Biochem J* 222: 195-201.
- Fein A, Payne R, Corson DW, Berridge MJ, Irvine RF (1984) Photoreceptor excitation and adaptation by inositol 1,4,5-trisphosphate. *Nature* 311: 157-160.
- Brown JE, Rubin LJ, Ghalayini AJ, Tarver AP, Irvine RF, et al. (1984) A biochemical and electrophysiological examination of myo-inositol polyphosphate as a putative messenger for excitation in *Limulus* ventral photoreceptor cells. *Nature* 311: 160-163.
- Burgess GM, Godfrey PP, McKinney JS, Berridge MJ, Irvine RF, et al. (1984) The second messenger linking receptor activation to internal Ca release in liver. *Nature* 309: 63-66.
- Prentki M, Biden TJ, Janjic D, Irvine RF, Berridge MJ, et al. (1984) Rapid mobilization of Ca²⁺ from rat insulinoma microsomes by inositol-1,4,5-trisphosphate. *Nature* 309: 562-564.
- Irvine RF, Brown KD, Berridge MJ (1984) Specificity of inositol trisphosphate-induced calcium release from permeabilized Swiss-mouse 3T3 cells. *Biochem J* 222: 269-272.
- Irvine RF, Letcher AJ, Heslop JP, Berridge MJ (1986) The inositol tris/tetrakisphosphate pathway--demonstration of Ins(1,4,5)P₃ 3-kinase activity in animal tissues. *Nature* 320: 631-634.
- Rapp PE, Berridge MJ (1981) The control of transepithelial potential oscillations in the salivary gland of *Calliphora erythrocephala*. *J Exp Biol* 93: 119-132.
- Missiaen L, Taylor CW, Berridge MJ (1991) Spontaneous calcium release from inositol trisphosphate-sensitive calcium stores. *Nature* 352: 241-244.
- Berridge MJ, Irvine RF (1984) Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* 312: 315-321.
- Berridge MJ (1987) Inositol trisphosphate and diacylglycerol: two interacting second messengers. *Annu Rev Biochem* 56: 159-193.
- Berridge MJ, Irvine RF (1989) Inositol phosphates and cell signalling. *Nature* 341: 197-205.
- Berridge MJ, Downes CP, Hanley MR (1989) Neural and developmental actions of lithium: a unifying hypothesis. *Cell* 59: 411-419.
- Berridge MJ (1993) Inositol trisphosphate and calcium signalling. *Nature* 361: 315-325.
- Bootman MD, Berridge MJ (1995) The elemental principles of calcium signaling. *Cell* 83: 675-678.
- Decrock E, De Bock M, Wang N, Gadicherla AK, Bol M, et al. (2013) IP₃, a small molecule with a powerful message. *Biochim Biophys Acta* 1833: 1772-1786.
- Shoichiro O, Yutaka W, Tomio O, Yoshihasa K, Naokazu S, et al. (1986) Total synthesis of optically active myo-inositol 1,4,5-tris(phosphate). *Tetrahedron Letters* 27: 3157-3160.
- Shoichiro O, Yoshihasa K, Naokazu S, Tomio O, Yutaka W (1992) Synthesis and some properties of D-myo-inositol 1,4,5-tris(dihydrogen phosphate). *Journal of the Chemical Society, Perkin Transactions 1*: 729-737.
- Falck JR, Abdali A, Wittenberger SJ (1990) Total synthesis of the 5-methylenephosphonate analogue of D-myo-inositol 1,4,5-trisphosphate. *J Chem Soc, Chem Commun*: 953-955.
- <http://www.sciencedirect.com/science/journal/00086215/140/1>
- Vacca JP, deSolms JS, Huff JR, Billington DC, Baker R, et al. (1989) The total synthesis of myo-inositol polyphosphates. *Tetrahedron* 45: 5679-5702.
- Ley SV, Parra M, Redgrave AJ, Sternfeld F (1990) Microbial oxidation in synthesis: preparation of myo-inositol phosphates and related cyclitol derivatives from benzene. *Tetrahedron* 46: 4995-5026.
- Falck JR, Yadagiri PJ (1989) Enantiospecific synthesis of D-myo-inositol 1,4,5-trisphosphate from (-)-quinic acid. *J Org Chem* 54: 5851-5852.
- <http://www.sciencedirect.com/science/journal/00404039/10/59>
- Maracek JP, Prestwich GD (1989) Synthesis of D-myo-(-3 H)(1,4,5)P₃. *J Labelled Comp* 27: 917.
- Prestwich GD, Marecek JF (1991) Chemical modification of inositol trisphosphate: Tritiated, fluorinated and phosphate-tethered analogues. *ACS Symposium Series 463 Inositolpolyphosphate and derivatives* Edit Allen B. Reitz: 3322-131.
- Potter BVL (1991) Phosphothioate analogues of D-myo-inositol 1,4,5-Trisphosphate. *ACS Symposium Series 463 Inositolpolyphosphate and derivatives* Edit Allen B. Reitz: 186-201.
- Takahiko A, Naoto T, Shoichiro O (1990) Chiral synthesis of D-myo-inositol 1-phosphate starting from L-quebrachitol. *Tetrahedron Letters* 31: 1433-1434.
- Takahiko A, Hiroaki S, Shoichiro O (1991) A concise synthesis of (-)-conduiritol F from L-quebrachitol via AlCl₃-n-Bu₄Nl mediated demethylation. *Tetrahedron Letters* 32: 5593-5696.
- Takahiko A, Masatoshi O, Hiroaki S, Shoichiro O (1991) Total synthesis of cyclophellitol from L-quebrachitol. *Synlett* 11: 831-832.
- Takahiko A, Hiroaki S, Masatoshi O, Tadashi O, Shoichiro O (1993) Synthesis of (-)-conduiritol F, (+)-conduiritol B, cyclophellitol from L-quebrachitol. *Bulletin of the Chemical Society of Japan* 66: 3760-3767.
- Takahiko A, Hiroyuki N, Takaaki K, Shoichiro O (1994) Stereodivergent synthesis of optically active α-hydroxy acids via diastereoselective reduction of α-keto esters derived from L-quebrachitol. *Bulletin of the Chemical Society of Japan* 67: 180-188.
- Yutaka W, Motohiro M, Shoichiro O (1987) Synthesis of optically active inositol derivatives starting from D-glucurono-6,3-lactone. *Chemistry Letters* 1: 123-126.
- Lei L, Yutaka W, Takahiko A, Shoichiro O (1992) A new efficient method for resolution of myo-inositol derivatives by enzyme catalyzed regio- and enantioselective esterification in organic solvent. *Tetrahedron Letters* 33: 1911-1914.
- Ling L, Li X, Watanabe Y, Akiyama T, Ozaki S (1993) Enzymatic resolution of racemic 1,2:5,6-di-O-cyclohexylidene and 1,2:3,4-di-O-cyclohexylidene-myo-inositol. *Bioorg Med Chem* 1: 155-159.
- Lei L, Shoichiro O (1993) Enzyme aided synthesis of D-myo-inositol 1,4,5-trisphosphate. *Tetrahedron Letter* 34: 2501-2504.
- Ling L, Ozaki S (1994) A chemoenzymatic synthesis of D-myo-inositol 1,4,5-trisphosphate. *Carbohydr Res* 256: 49-58.

47. Lei L, Shoichiro O (1995) Enzymic resolution of the sterically hindered myo-inositol derivative. *Bulletin of the Chemical Society of Japan* 68: 1200-1205.
48. Shoichiro O, Lei L (1997) Chemoenzymatic Synthesis of Optically Active myo-inositol Polyphosphate. *Carbohydrates in Drug Design*.
49. Shoichiro O, Kayoko Y, Takako K, Takahiro M, Mitsuaki E, et al. (1995) Enzyme aided regioselective acylation of nucleosides. *Nucleosides & Nucleotides* 14: 401-104.
50. Yutaka W, Akinori O, Yasushi S, Shoichiro O (1990) Easy access of optically active myo-inositol derivatives by enantioselective acylation using a tartaric acid monoester. *Tetrahedron Letters* 31: 2613-2616.
51. Yutaka W, Tadashi O, Takashi Y, Shoichiro O (1996) Synthetic strategies based on phosphite chemistry for inositol phosphates and phospholipids. *Phosphorus, Sulfur and Silicon and the Related Elements*: 109-110.
52. Yutaka W, Hiroyuki N, Motonobu B, Shoichiro O (1987) An efficient method for polyphosphorylation of inositol derivatives. *Tetrahedron Letters* 28: 4179-4180.
53. Yutaka W, Yasunobu K, Katsumi E, Shoichiro O (1990) An efficient phosphorylation method using a new phosphitylating agent, 2-(diethylamino)-1,3,2-benzodioxaphosphane. *Tetrahedron Letters* 31: 255-256.
54. Yutaka W, Yasunobu K, Shoichiro O (1992) Utilization *o*-xylylene *N,N*-diethylphosphoramidite for the synthesis of phosphoric diesters. *Tetrahedron Letters* 33: 133-136.
55. Yutaka W, Nobuyuki H, Shoichiro O (1988) Dibenzyl phosphorofluoridate, a new phosphorylating agent. *Tetrahedron Letters* 29: 5763-5764.
56. Yutaka W, Shinsuke S, Shoichiro O, Masato H (1996) Synthesis of phosphorofluoridate analogs of myo-inositol 1,4,5-tris(phosphate) and their biological activity. *Chemical Communications (Cambridge)* 15: 1815-1816.
57. Yutaka W, Tomio O, Naokazu S, Shoichiro O (1987) Stepwise phosphorylation of vicinal diol and sterically hindered alcohol directed toward D-myo-inositol 2,4,5-trisphosphate. *Tetrahedron Letters* 28: 2607-2610.
58. Yutaka W, Shinji I, Takashi Y, Shoichiro O (1995) Phosphorothioate synthesis based on the redox reaction of phosphite with tellurium(IV) chloride. *Synthesis* 10: 1243-1244.
59. Yutaka W, Eiji I, Masanao J, Shoichiro O (1993) Phosphonium salt methodology for the synthesis of phosphoric monoesters and diesters and its application to selective phosphorylation. *Tetrahedron Letters* 34: 497-500.
60. Shoichiro O, Yutaka W, Takehiro M, Yuichi H, Tomio O, et al. (1998) Synthesis of phosphatidyl-myo-inositol polyphosphates and derivatives. *ACS Symposium Series 718 Phosphoinositides*, Edit Karol Bruzik: 212-221.
61. Yutaka W, Shinichi M, Shoichiro O (1992) 1H-Tetrazole catalyzed reactions of trialkyl phosphites: observation of a five-membered cyclic hydrophosphorane intermediate during the transesterification. *Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry* 15: 1879-1880.
62. Yutaka W, Takashi Y, Taroh I, Shoichiro O (1994) Utilization of oxidizing character of TeCl₄ in the reaction with trialkyl phosphite. A novel redox phosphorylation. *Chemistry Letters* 10: 1881-1884.
63. Yutaka W, Hajimu H, Shoichiro O (1994) Synthesis of a phosphatidylinositol 3,4,5-trisphosphate. *Tetrahedron Letters* 35: 123-124.
64. Yutaka W, Masaya T, Shoichiro O (1995) Synthesis of 1D-distearoylphosphatidyl-myo-inositol 3,4,5-tris(dihydrogen phosphate). *Tetrahedron* 51: 8969-8976.
65. Shoichiro O, Xiang-Zheng K, Yutaka W, Tomio O (1998) Synthesis of unsaturated phosphatidyl inositol-3,4,5-trisphosphate. *Chinese Journal of Chemistry* 16: 51-57.
66. Shoichiro O, Watanabe Y (1991) Synthesis of Inositol Polyphosphate and Their Derivatives. *ACS Symposium Series 463 Inositolpolyphosphate and derivatives* Edit Allen B Reitz: 41-64.
67. Yutaka W, Chikara N, Shoichiro O, Michikatsu S, Kyoko K (1993) Absolute configuration of 4-*a*-D-glucopyranosyl-myo-inositol, enzymic transglycosylation product. *Journal of Carbohydrate Chemistry* 12: 685-692.
68. Yutaka W, Takashi Y, Shoichiro O (1996) Synthesis of 2,6-Di-O-(*a*-D-mannopyranosyl)phosphatidyl-D-myo-inositol. *Tetrahedron* 53: 903-918.
69. Yutaka W, Motohiro M, Takao M, Shoichiro O (1989) Highly efficient protection by the tetraisopropylidisiloxane-1,3-diyl group in the synthesis of myo-inositol phosphates as inositol 1,3,4,6-tetrakisphosphate. *Journal of the Chemical Society, Chemical Communications* 8: 482-483.
70. Yutaka W, Masanori I, Shoichiro O (1994) Proximally assisted and chemoselectively cleavable protecting groups for alcohols, 2-[2-(arylmethoxy)ethyl]benzoic esters. *Chemistry Letters* 11: 2163-2166.
71. Takahiko A, Naoto T, Hiroaki S, Shoichiro O (1990) Anchimerically assisted demethylation of methyl ethers in inositol derivatives with an aluminum chloride-sodium iodide system. *Chemistry Letters* 10: 1881-1884.
72. Takahiko A, Hajimu H, Shoichiro O (1992) Aluminium chloride-*N,N*-dimethylaniline: a novel benzyl and allyl ether cleavage reagent. *Bulletin of the Chemical Society of Japan* 65: 1932-1938.
73. Takahiko A, Hiroaki S, Shoichiro O (1992) Trimethylsilyl chloride-tin(II)chloride-anisole: a novel *p*-methoxybenzyl ether cleavage reagent. *Synlett* 5: 415-416.
74. Takahiko A, Hiroyuki N, Keiichi I, Shoichiro O (1992) Diastereoselective addition of organometallics to *a*-keto esters bearing chiro-inositol derivatives as chiral auxiliaries. *Chemistry Letters* 3: 447-450.
75. Tadahiaki A, Hiruyuki N, Shoichiro O (1991) Diastereoselective reduction of *a*-ketoesters bearing chiro-inositol derivatives as chiral auxiliaries. *Tetrahedron Letters* 32: 1335-1338.
76. Yutaka W, Takahiro F, Shoichiro O (1992) A novel deacylation method using Grignard reagent without affecting the neighboring base-sensitive functional groups. *Journal of the Chemical Society, Chemical Communications* 9: 681-683.
77. Ozaki S, Uemura A, Konishi T, Yamashita K, Maekawa T, et al. (1993) Enzyme aided regio-selective acylation and deacylation of nucleosides. *Nucleic Acids Symp Ser*: 53-54.
78. Yutaka W, Chikara N, Shoichiro O (1993) Glycosidation based on phosphite chemistry. *Synlett* 2: 115-116.
79. Yutaka W, Chikara N, Takashi Y, Shoichiro O (1994) Glycosylation using glycosyl phosphite as a glycosyl donor. *Tetrahedron* 50: 6523-6536.
80. Van Dijken P, Lammers AA, Ozaki S, Potter BV, Erneux C, et al. (1994) Phosphorylation of inositol 1,4,5-trisphosphate analogues by 3-kinase and dephosphorylation of inositol 1,3,4,5-tetrakisphosphate analogues by 5-phosphatase. *Eur J Biochem* 226: 561-566.
81. Akiyama T, Yasusa T, Ishikawa K, Ozaki S (1994) Asymmetric synthesis of tetrahydrofurans by diastereoselective [3+2]-cycloaddition of allylsilanes with *a*-keto esters bearing an optically active cyclitol as a chiral auxiliary. *Tetrahedron Letters* 35: 8401-8404.
82. Akiyama T, Okada K, Ozaki S (1992) The preparation of optically active D 2-isoxazolines via addition of nitrile oxides to chiral acryloyl esters bearing cyclitols as auxiliaries. *Tetrahedron Letters* 33: 5763-5766.
83. Akiyama T, Ishikawa K, Ozaki S (1994) Asymmetric synthesis of functionalized tertiary alcohols by diastereoselective aldol reaction of silyl enol ether and ketene silyl acetal with *a*-keto esters bearing an optically active cyclitol as a chiral auxiliary. *Synlett* 4: 275-276.
84. Ozaki S, Watanabe Y, Hirata M, Awaya A (1991) Preparation of inositol polyphosphate derivatives for control of the calcium ion-participating metabolic steps. *PCT Int. Appl*: 139.
85. Ozaki S, Watanabe Y (1989) Synthesis of inositol polyphosphates. *Yuki Gosei Kagaku Kyokaiishi* 47: 363-373.
86. Watanabe Y, Ozaki S (1992) Organic chemical approach to inositol phospholipid-associated signal transduction. *Farumashia* 28: 598-602.
87. Ozaki S, Akiyama T, Takechi N, Kageyama K, Machida M (1991) Preparation of d-myo-inositol 1-phosphate. *Ger Offen* 8.
88. Akiyama T, Takechi N, Ozaki S, Shiota K (1992) A chiral synthesis of D-myo-inositol 1-phosphate starting from L-quebrachitol. *Bulletin of the Chemical Society of Japan* 65: 366-372.
89. Ozaki S, Kohno M, Nakahira H, Bunya M, Watanabe Y (1988) Synthesis of optically active myo-inositol 1,3,4-trisphosphate. *Chemistry Letters*: 77-80.
90. Watanabe Y, Ogasawara T, Ozaki S, Hirata M (1994) Synthesis of myo-inositol 1,4,6-trisphosphate, an analog of myo-inositol 1,4,5-trisphosphate. *Carbohydrate Research* 258: 87-92.
91. Ozaki S, Kondo Y, Nakahira H, Yamaoka S, Watanabe Y (1987) Synthesis of D-myo-inositol 1,3,4,5-tetrakisphosphate. *Tetrahedron Letters* 28: 4691-4694.

92. Watanabe Y, Shinohara T, Fujimoto T, Ozaki S (1990) A short step and practical synthesis of myo-inositol 1,3,4,5-tetrakisphosphate. *Chemical & Pharmaceutical Bulletin* 38: 562-563.
93. Watanabe Y, Fujimoto T, Shinohara T, Ozaki S (1991) A short step synthesis of optically active myo-inositol 1,3,4,5-tetrakis(phosphate) and myo-inositol 1,4,5-tris(phosphate) from 1,3,5-tri-O-benzoyl-myo-inositol. *Journal of the Chemical Society, Chemical Communications* 6: 428-429.
94. Ozaki S, Ling L, Ogasawara T, Watanabe Y, Hirata M (1994) A convenient chemoenzymatic synthesis of D-and L-myo-inositol 1,4,5,6-tetrakisphosphate. *Carbohydr Res* 259: 307-310.
95. Watanabe Y, Ogasawara T, Nakahira H, Matsuki T, Ozaki S (1988) A versatile intermediate, D-4,5-bis(dibenzyl phosphoryl)-myo-inositol derivative, for synthesis of inositol phosphates. Synthesis of 1,2-cyclic-4,5-, 1,4,5-, and 2,4,5-trisphosphate. *Tetrahedron Letters* 29: 5259-5262.
96. Ozaki S, Watanabe Y, Ogasawara T, Hirata M, Kanematsu T (1992) Synthesis and biological properties of 2-substituted myo-inositol 1,4,5-trisphosphate analogues directed toward affinity chromatography and photoaffinity labeling. *Carbohydr Res* 234: 189-206.
97. Hirata M, Watanabe Y, Ishimatsu T, Ikebe T, Kimura Y, et al. (1989) Synthetic inositol trisphosphate analogs and their effects on phosphatase, kinase, and the release of Ca²⁺. *J Biol Chem* 264: 20303-20308.
98. Hirata M, Watanabe Y, Ishimatsu T, Yanaga F, Koga T, et al. (1990) Inositol 1,4,5-trisphosphate affinity chromatography. *Biochem Biophys Res Commun* 168: 379-386.
99. Hirata M, Yanaga F, Koga T, Ogasawara T, Watanabe Y, et al. (1990) Stereospecific recognition of inositol 1,4,5-trisphosphate analogs by the phosphatase, kinase, and binding proteins. *J Biol Chem* 265: 8404-8407.
100. Hirata M, Kimura Y, Ishimatsu T, Yanaga F, Shuto T, et al. (1991) Synthetic inositol 1,3,4,5-tetrakisphosphate analogues. *Biochem J* 276: 333-336.
101. Kimura Y, Kanematsu T, Watanabe Y, Ozaki S, Koga T, et al. (1991) Synthetic inositol 1,3,4,5-tetrakisphosphate analogs and their effect on the binding to microsomal fraction of rat cerebellum. *Biochimica et Biophysica Acta, Biomembranes* 1069: 218-222.
102. Kanematsu T, Takeya H, Watanabe Y, Ozaki S, Yoshida M, et al. (1992) Putative inositol 1,4,5-trisphosphate binding proteins in rat brain cytosol. *J Biol Chem* 267: 6518-6525.
103. Kamata H, Hirata M, Ozaki S, Kusaka I, Kagawa Y, et al. (1992) Partial purification and reconstitution of inositol 1,4,5-trisphosphate receptor/Ca²⁺ channel of bovine liver microsomes. *J Biochem* 111: 546-552.
104. Hirata M, Watanabe Y, Yoshida M, Koga T, Ozaki S (1993) Roles for hydroxyl groups of D-myo-inositol 1,4,5-trisphosphate in the recognition by its receptor and metabolic enzymes. *J Biol Chem* 268: 19260-19266.
105. Hashii M, Hirata M, Ozaki S, Nozawa Y, Higashida H (1994) Ca²⁺ influx evoked by inositol-3,4,5,6-tetrakisphosphate in ras-transformed NIH/3T3 fibroblasts. *FEBS Lett* 340: 276-280.
106. Hirata M, Narumoto N, Watanabe Y, Kanematsu T, Koga T, et al. (1994) DL-myo-inositol 1,2,4,5-tetrakisphosphate, a potent analog of D-myo-inositol 1,4,5-trisphosphate. *Mol Pharmacol* 45: 271-276.
107. Hashii M, Hirata M, Ozaki S, Nozawa Y, Higashida H (1994) Ca²⁺ influx gated by inositol-3,4,5,6-tetrakisphosphate in NIH/3T3 fibroblasts. *Biochem Biophys Res Commun* 200: 1300-1306.
108. Hirata M, Watanabe Y, Kanematsu T, Ozaki S, Koga T (1995) D-myo-inositol 1,4,5-trisphosphate analogues substituted at the 3-hydroxyl group. *Biochim Biophys Acta* 1244: 404-410.
109. Baron CB, Ozaki S, Watanabe Y, Hirata M, LaBelle EF, et al. (1995) Inositol 1,4,5-trisphosphate binding to porcine tracheal smooth muscle aldolase. *J Biol Chem* 270: 20459-20465.
110. Kanematsu T, Misumi Y, Watanabe Y, Ozaki S, et al. (1996) A new inositol 1,4,5-trisphosphate binding protein similar to phospholipase C-d 1. *Biochemical Journal* 313: 319-325.
111. Watanabe Y, Hirata M, Ogasawara T, Koga T, Ozaki S (1991) Synthesis and characterization of a photoaffinity probe possessing biotinyl and azidobenzoyl moieties for IP₃-affiliated protein. *Bioorganic & Medicinal Chemistry Letters* 1: 399-402.
112. Ozaki S, Koga Y, Ling L, Watanabe Y, Kimura Y, et al. (1994) Synthesis of 2-substituted myo-inositol 1,3,4,5-tetrakis(phosphate) and 1,3,4,5,6-pentakis(phosphate) analogs. *Bulletin of the Chemical Society of Japan* 67: 1058-1063.
113. Ozaki S, Watanabe Y, Hirata M, Ogasawara Y, Kanematsu T, et al. (1993) Inositol 1,4,5-trisphosphate Affinity Chromatography. Fishing out Ins (1,4,5) P₃-recognizable Protein. *Drug Design for neuroscience* Edit Alan Kozikowski Raven Press: 417-434.
114. Ozaki S, Kong X, Watanabe Y, Ogasawara T (1997) Synthesis of affinity column of phosphatidylinositol-3,4-diphosphate. *Chinese Journal of Chemistry* 15: 556-561.
115. Yagisawa H, Hirata M, Kanematsu T, Watanabe Y, Ozaki S, et al. (1994) Expression and characterization of an inositol 1,4,5-trisphosphate binding domain of phosphatidylinositol-specific phospholipase C-delta 1. *J Biol Chem* 269: 20179-20188.
116. Yoshida M, Kanematsu T, Watanabe Y, Koga T, Ozaki S, et al. (1994) D-myo-inositol 1,4,5-trisphosphate-binding proteins in rat brain membranes. *J Biochem* 115: 973-980.
117. Hirata M, Kanematsu T, Sakuma K, Koga T, Watanabe Y, et al. (1994) D-myo-Inositol 1,4,5-trisphosphate binding domain of phospholipase C-d 1. *Biochemical and Biophysical Research Communications* 205: 1563-1571.
118. Moriya S, Kazlauskas A, Akimoto K, Hirai S, Mizuno K, et al. (1996) Platelet-derived growth factor activates protein kinase C epsilon through redundant and independent signaling pathways involving phospholipase C gamma or phosphatidylinositol 3-kinase. *Proc Natl Acad Sci U S A* 93: 151-155.
119. Hogan SP, Foster PS, Hansbro PM, Ozaki S, Denborough MA (1994) Detection and partial purification of inositol 1,4,5-trisphosphate 3-kinase from porcine skeletal muscle. *Cell Signal* 6: 233-243.
120. Hidenori M, Hiroaki K, Motonari T, Yoshitaka O, Tadaomi T, et al. (1996) Isolation of the active form of RAC-protein kinase (PKB/Akt) from transfected COS-7 cells treated with heat shock stress and effects of phosphatidylinositol 3,4,5-trisphosphate and phosphatidylinositol 4,5-bisphosphate on its enzyme activity. *FEBS Letters* 396: 305-308.
121. Foster PS, Hogan SP, Hansbro PM, O'Brien R, Potter BV, et al. (1994) The metabolism of D-myo-inositol 1,4,5-trisphosphate and D-myo-inositol 1,3,4,5-tetrakisphosphate by porcine skeletal muscle. *Eur J Biochem* 222: 955-964.
122. Yagisawa H, Hirata M, Kanematsu T, Watanabe Y, Ozaki S, et al. (1994) Expression and characterization of an inositol 1,4,5-trisphosphate binding domain of phosphatidylinositol-specific phospholipase C-d 1. *Journal of Biological Chemistry* 269: 79-88.
123. Bertsch U, Haefs M, Moller M, Deschermeier C, Fanick W, et al. (1999) A novel A-isoform-like inositol 1,4,5-trisphosphate 3-kinase from chicken erythrocytes exhibits alternative splicing and conservation of intron positions between vertebrates and invertebrates. *Gene* 228: 61-71.
124. Inoue M, Sakamoto Y, Fujishiro N, Imanaga I, Ozaki S, et al. (2003) Homogeneous Ca²⁺ stores in rat adrenal chromaffin cells. *Cell Calcium* 33: 19-26.
125. Lin H, Ozaki S, Fujishiro N, Takeda K, Imanaga I, et al. (2005) Subunit composition and role of Na⁺,K⁺-ATPases in adrenal chromaffin cells. *J Physiol* 564: 161-172.
126. Ronald WH, Michael DH, Scott DS, Stephen KF, Balla T, et al. (2000) A pleckstrin homology domain specific for phosphatidylinositol 4,5-bisphosphate (PtdIns-4,5-P₂) and fused to green fluorescent protein identifies plasma membrane PtdIns-4,5-P₂ as being important in exocytosis. *Journal of Biological Chemistry* 275: 17878-17885.
127. Ozaki S, Wald D, Shope BD, Chen CJ, Jian P, et al. (2000) Intracellular delivery of phosphoinositides and inositol phosphates using polyamine carriers. *Proceedings of the National Academy of Sciences of the United States of America* 97: 11286-11291.
128. Prestwich GD, Chen R, Feng L, Ozaki S, Ferguson CG, et al. (2002) In situ detection of phospholipid and phosphoinositide metabolism. *Adv Enzyme Regul* 42: 19-38.
129. Dewald DB, Ozaki S, Malaviya S, Shope JC, Manabe K, et al. (2005) Cellular calcium mobilization in response to phosphoinositide delivery. *Cell Calcium* 38: 59-72.

130. Ozaki S, DeWald DB, Shope JC, Chen J, Prestwich GD (2000) Intracellular delivery of phosphoinositides and inositol phosphates using polyamine carriers. *Proc Natl Acad Sci U S A* 97: 11286-11291.
131. Maruyama T, Kanaji T, Nakade S, Kanno T, Mikoshiba K (1997) 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P₃-induced Ca²⁺ release. *J Biochem* 122: 498-505.
132. Iwasaki H, Mori Y, Hara Y, Uchida K, Zhou H, et al. (2001) 2-Aminoethoxydiphenyl borate (2-APB) inhibits capacitative calcium entry independently of the function of inositol 1,4,5-trisphosphate receptors. *Receptors Channels* 7: 429-439.
133. Zhou H, Iwasaki H, Nakamura T, Nakamura K, Maruyama T, et al. (2007) 2-Aminoethyl diphenylborinate analogues: selective inhibition for store-operated Ca²⁺ entry. *Biochem Biophys Res Commun* 352: 277-282.
134. Mikoshiba K, Ozaki S, Suzuki A, Nakamura T (2007) Preparation of bisboron compounds controlling calcium concentration in cells. *PCT Int Appl* 118.
135. Mikoshiba K, Ozaki S, Ebii E (2009) Preparation of phenylborinic acid, poly(arylhydroxyborane), and their esters as intracellular calcium concentration regulators. *Jpn Kokai Tokkyo Koho* 138.
136. Mikoshiba K, Nukina N, Ozaki S, Hamada K, Goto J, et al. (2010) Preparation of phenylboron compounds as polyglutamine aggregation inhibitor.
137. Goto J, Suzuki AZ, Ozaki S, Matsumoto N, Nakamura T, et al. (2010) Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca(2+) entry via STIM proteins. *Cell Calcium* 47: 1-10.
138. Suzuki AZ, Ozaki S, Goto J, Mikoshiba K (2010) Synthesis of bisboron compounds and their strong inhibitory activity on store-operated calcium entry. *Bioorganic & Medicinal Chemistry Letters* 20: 1395-1398.
139. Ozaki S, Suzuki AZ, Bauer PO, Ebisui E, Mikoshiba K (2013) 2-Aminoethyl diphenylborinate (2-APB) analogues: regulation of Ca²⁺ signaling. *Biochem Biophys Res Commun* 441: 286-290.
140. Ozaki S (2014) 2-Aminoethyl diphenylborinate (2APB) analogues: part 2. regulators of Ca²⁺ release and consequent cellular processes. *Archives of Physiology* 1: 1-6.
141. Mackenzie L, Bootman MD, Berridge MJ, Lipp P (2001) Predetermined recruitment of calcium release sites underlies excitation-contraction coupling in rat atrial myocytes. *J Physiol* 530: 417-429.
142. Lipp P, Laine M, Tovey SC, Burrell KM, Berridge MJ, et al. (2000) Functional InsP₃ receptors that may modulate excitation-contraction coupling in the heart. *Curr Biol* 10: 939-942.
143. Mackenzie L, Bootman MD, Laine M, Berridge MJ, Thuring J, et al. (2002) The role of inositol 1,4,5-trisphosphate receptors in Ca(2+) signalling and the generation of arrhythmias in rat atrial myocytes. *J Physiol* 541: 395-409.
144. Proven A, Roderick HL, Conway SJ, Berridge MJ, Horton JK, et al. (2006) Inositol 1,4,5-trisphosphate supports the arrhythmogenic action of endothelin-1 on ventricular cardiac myocytes. *J Cell Sci* 119: 3363-3375.
145. Berridge MJ (2010) Calcium hypothesis of Alzheimer's disease. *Pflugers Arch* 459: 441-449.
146. Berridge MJ (2011) Calcium signalling and Alzheimer's disease. *Neurochem Res* 36: 1149-1156.
147. Chen R, Valencia I, Zhong F, McColl KS, Roderick HL, et al. (2004) Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate receptors to regulate calcium release from the ER in response to inositol 1,4,5-trisphosphate. *J Cell Biol* 166: 193-203.
148. Berridge MJ (2013) Dysregulation of neural calcium signaling in Alzheimer disease, bipolar disorder and schizophrenia. *Prion* 7: 2-13.
149. Bauer OP, Hudec R, Ozaki S, Okuno M, Ebisui E, et al. (2011) Genetic ablation and chemical inhibition of IP3R1 reduce mutant Huntington aggregation. *Biochem Biophys Res Commun* 416: 13-17.
150. Zimmermann TJ, Bürger M, Tashiro E, Kondoh Y, Martinez NE, et al. (2013) Boron-based inhibitors of acyl protein thioesterases 1 and 2. *Chembiochem* 14: 115-122.
151. Ozaki S, Ebisui E, Hamada K, Goto J, Suzuki AZ, et al. (2010) Potent transglutaminase inhibitors, aryl beta-aminoethyl ketones. *Bioorg Med Chem Lett* 20: 1141-1144.
152. Ozaki S, Ebisui E, Hamada K, Suzuki AZ, Terauchi A, et al. (2011) Potent transglutaminase inhibitors, Dithioβ-aminoethyl ketones. *Bioorganic & Medicinal Chemistry Letters* 21: 377-379.