

Chromatin Remodeling Directs the Fate of Epithelial Cells on Soft Matrix

Dominique Vautier*

Inserm UMR 1121, 11 rue Humann, 67085 Strasbourg, France. Université de Strasbourg, Faculté de Chirurgie Dentaire, 8 rue Sainte-Elisabeth, 67000 Strasbourg, France. Fédération de Médecine Translationnelle, Strasbourg

*Corresponding author: Dominique Vautier, Inserm UMR 1121, 11 rue Humann, 67085 Strasbourg, France. Université de Strasbourg, Faculté de Chirurgie Dentaire, 8 rue Sainte-Elisabeth, 67000 Strasbourg, France. Fédération de Médecine Translationnelle, Strasbourg, Tel: 330368853374; E-mail: vautier@unistra.fr

Received date: May 23, 2016; Accepted date: May 30, 2016; Published date: June 02, 2016

Copyright: © 2016 Vautier D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: Hydrogels; Substrate stiffness; Chromatin; Histone deacetylase inhibition

Introduction

Cell therapy is an emerging field in the development of regenerative medicine, tissue engineering and cancer immunotherapy. Its efficiency depends on the release of healthy cells to injured sites. These cells can exert therapeutic effects, repair tissue and damaged organs or eliminate malignant tumors. On the other hand, hydrogels, due to their biochemical and physical properties, are materials of choice to mimic the extracellular matrix. Devising strategies to improve cell survival in hydrogels is a major challenge still in progress. To successfully design hydrogels, it is important to know the cellular responses like adhesion, migration, contraction and protrusion to environmental signals. Mechanotransduction events by which cells sense their environment were extensively investigated on mechanosensitive proteins at focal adhesions and inside the cytoskeleton [1-4]. Extra- and intracellular forces are transmitted across the cytoskeleton to the nucleus. These forces can activate integrins at focal adhesions linked to

actin filaments, themselves connected to microtubules and to intermediate filaments. The linker of nucleoskeleton and cytoskeleton complex (LINC complex), which enables force transmission across the nuclear envelope, connects cytoskeletal filaments to the nucleus where lamins form an extended part of the LINC complex [5-10]. These forces ultimately propagate to chromatin that represents a site of signal integration and interpretation for activation or gene silencing [11-12].

Interestingly, for the first time, Rabineau et al. [13] focused on how chromatin plasticity is influenced by a change of substrate rigidity. In this context, several works revealed that the nucleus itself acts as a cellular mechanosensor bypassing propagation of mechano-signaling processes through the cytoplasm [8,14]. Rabineau et al. [13] developed a hydrogel based on polyelectrolyte multilayers, made of poly(L-lysine) and hyaluronic acid, (PLL/HA)₂₄, capped with a poly(sodium styrene sulfonate)/poly(allylamine hydrochloride), (PSS/PAH)_n, multilayer film (Figure 1) as matrix model mimicking the extracellular matrix rigidity of biological tissues [15-17], on which epithelial PtK2 cells were grown. In their model, the rigidity of the hydrogel increases with the number n of PSS/PAH layer pairs [18].




Figure 1: Vertical section image of a (PLL/HA)₂₃-PLL^{FITC}-HA capped with a (PSS/PAH)_n-PSS^{Rho}-PAH multilayer film observed by CLSM.

Two well-defined cytological compartments are structured in the nucleus: the condensed, inactive heterochromatin and the extended, active euchromatin. Heterochromatin is restricted to an irregular rim located at the nuclear periphery and around the nucleolus as well as in patches throughout the nucleoplasm, whereas euchromatin fills up the majority of the nucleus. Rabineau et al. revealed that on stiff matrices, chromatin is in its euchromatin form, whereas a soft matrix partially induces remodeling in its heterochromatin form. On a very soft matrix cells die by necrosis. The original idea of the authors is to influence the cell fate by modifying the organization of the chromatin. The opposing actions of histone acetyl transferases and histone deacetylases (HDACs) dynamically control the acetylation status of chromatin and hence chromatin compaction, by respectively loosening (euchromatin) or condensing (heterochromatin) chromatin structures [19]. More

precisely, concerning the formation of heterochromatin, it has been shown that both of H3K9me_{2/3} and H3K27me₃ are associated with this process in mammals [19]. H3K9me_{2/3} is able to recruit HP1 α , which plays a key role in the stability of the higher-order structure of pericentric heterochromatin [20]. Interestingly, as both of the G9a/GLP complex and Suv39h1/2 enzymes interact with DNA methyltransferases and play a role in the maintenance of DNA methylation [21,22], the writers of H3K9me_{2/3} are also required for the formation of heterochromatin independently of HP1 α at specific chromatin loci. The writer for the establishment of H3K27me₃ is the PRC2 complex [23]. Recently, it has been shown that PRC2 and H3K27me₃ cooperate with H3K9 methylation to maintain HP1 α at chromatin [24].

Rabineau et al. [13] use a drug that inhibits histone deacetylases, namely trichostatin A (TSA), to maintain chromatin in euchromatin. On the very soft matrices, they find that PtK2 cells treated by TSA preserve acetylated forms of histones H3, maintain their chromatin in euchromatin and a uniform nuclear distribution of HP1 β . These treated cells maintain also their nuclear envelopes intact as well as a residual intermediate filament network around their nuclei. This allows cells to survive in a non-adherent state without undergoing proliferation suggesting that cells might enter in a quiescence state. Importantly, when transferred on a stiff matrix these cells retain their capacity to adhere, to spread and to enter a novel mitotic cycle in a way that depends on their transcriptional competence. These findings might be relevant to maintain cells in the best settings within synthetic scaffolds and in tissue-derived matrices used in tissue regeneration strategies.

The work of Rabineau et al. [13] represent the rational for detailed molecular studies aimed at developing drugs to preserve important nuclear structure, in particular euchromatin and the nuclear envelope, under unfavorable mechanical environment. Chromatin organization has a strong influence on the expression of the genome and chromatin remodeling contributes to many cellular properties as for instance the deformation of the nucleus as well as the cell pluripotency and the cell differentiation [25]. Hydrogels with variable elastic moduli might control the organisation of chromatin and therefore guide the fate of cells of interest for tissue engineering. For example, the behavior of several types of stem cells will be tested according to their level of differentiation. An important aspect is to determine if the elastic modulus durably remodels chromatin after mechanical pre conditioning on a soft substrate. In recent works, the elastic modulus can epigenetically reprogram the cells which maintain a "nuclear mechanical memory" [26-28].

References

- Riveline D, Zamir E, Balaban NQ, Schwartz US, Ishizaki T, et al. (2001) Focal contacts as mechanosensors: Externally applied local mechanical force induces growth of focal contacts by an mDia1-dependent and ROCK-independent mechanism. *J Cell Biol* 153:1175-1186.
- Vogel V, Sheetz M (2006) Local force and geometry sensing regulate cell functions. *Nat Rev Mol Cell Biol* 7: 265-275.
- Roca-Cusachs P, del Rio A, Puklin-Faucher E, Gauthier NC, Biais N, et al. (2013) Integrin-dependent force transmission to the extracellular matrix by α -actinin triggers adhesion maturation. *Proc Natl Acad Sci USA* 110:1361-1370.
- Dumbauld DW, Lee TT, Singh A, Scrimgeour J, Gersbach CA, et al. (2013) How vinculin regulates force transmission. *Proc Natl Acad Sci U S A* 110: 9788-9793.
- Crisp M, Liu Q, Roux K, Rattner JB, Shanahan C, et al. (2006) Coupling of the nucleus and cytoplasm: role of the LINC complex. *J Cell Biol* 172: 41-53.
- Lombardi ML, Jaalouk DE, Shanahan CM, Burke B, Roux KJ, et al. (2011) The interaction between nesprins and sun proteins at the nuclear envelope is critical for force transmission between the nucleus and cytoskeleton. *J Biol Chem* 286: 26743-26753.
- Iyer KV, Pulford S, Mogilner A, Shivashankar GV (2012) Mechanical activation of cells induces chromatin remodeling preceding MKL nuclear transport. *Biophys J* 103: 1416-1428.
- Wang N, Tytell JD, Ingber DE (2009) Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol* 10: 75-82.
- Mellad JA, Warren DT, Shanahan CM (2011) Nesprins LINC the nucleus and cytoskeleton. *Curr Opin Cell Biol* 23: 47-54.
- Gruenbaum Y, Margalit A, Goldman RD, Shumaker DK, Wilson KL (2005) The nuclear lamina comes of age. *Nat Rev Mol Cell Biol* 6: 21-31.
- Schneider R, Grosschedl R (2007) Dynamics and interplay of nuclear architecture, genome organization, and gene expression. *Genes Dev* 21: 3027-3043.
- Shimi T, Pfliegerhaer K, Kojima S, Pack CG, Solovei I, et al. (2008) The A- and B- type nuclear lamin networks microdomains in chromatin organization and transcription. *Genes Dev* 22: 3409-21.
- Rabineau M, Flick F, Mathieu E, Tu A, Senger B, et al. (2015) Cell guidance into quiescent state through chromatin remodeling induced by elastic modulus of substrate. *Biomaterials* 37: 144-155.
- Isermann P, Lammerding J (2013) Nuclear mechanics and mechanotransduction in health and disease. *Curr Biol* 23: R1113-1121.
- Mertz D, Vogt C, Hemmerlé J, Mutterer J, Ball V, et al. (2009) Mechanotransductive surfaces for reversible biocatalysis activation. *Nat Mater* 8: 731-735.
- Kocgozlu L, Lavalle P, Koenig G, Senger B, Haikel Y, et al. (2010) Selective and uncoupled role of substrate elasticity in the regulation of replication and transcription in epithelial cells. *J Cell Sci* 123: 29-39.
- Levental I, Georges PC, Janmey PA (2007) Soft biological materials and their impact on cell function. *Soft Matter* 3: 299-306.
- Francius G, Hemmerlé J, Ball V, Lavalle P, Picart C, et al. (2007) Stiffening of soft polyelectrolyte architectures by multilayer capping evidenced by viscoelastic analysis of AFM indentation measurements. *J Phys Chem C* 111: 8299-8306.
- Kouzarides T (2007) Chromatin modifications and their function. *Cell* 128: 693-705.
- Bannister AJ, Zegerman P, Partridge JF, Miska EA, Thomas JO, et al. (2001) Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. *Nature* 410: 120-124.
- Fuks F, Hurd PJ, Deplus R, Kouzarides T (2003) The DNA methyltransferases associate with HP1 and the SUV39H1 histone methyltransferase. *Nucleic Acids Res* 31: 2305-2312.
- Zhang T, Termanis A, Özkan B, Culley J, de Lima Alves F, et al. (2016) G9a/GLP complex maintains imprinted DNA methylation in embryonic stem cells. *Cell Rep* 15: 77-85.
- Margueron R, Reinberg D (2011) The Polycomb complex PRC2 and its mark in life. *Nature* 469: 343-349.
- Boros J, Arnoult N, Stroobant V, Collet JF, Decottignies A (2014) Polycomb repressive complex 2 and H3K27me3 cooperate with H3K9 methylation to maintain heterochromatin protein 1a at chromatin. *Mol Cell Biol* 34: 3662-3674.
- Meshorer E, Misteli T (2006) Chromatin in pluripotent embryonic stem cells and differentiation. *Nat Rev Mol Cell Biol* 7: 540-546.
- Yang C, Tibbitt MW, Basta L, Anseth KS (2014) Mechanical memory and dosing influence stem cell fate. *Nat Mater* 13: 645-652.
- Lee J, Abdeen AA, Kilian KA (2014) Rewiring mesenchymal stem cell lineage specification by switching the biophysical microenvironment. *Sci Rep* 4:5188.
- Heo SJ, Thorpe S, Driscoll TP, Duncan RL, Lee DA, et al. (2015) Biophysical regulation of chromatin architecture instills a mechanical memory in mesenchymal stem cells. *Sci Rep* 5:16895.