

# Classification of 2-Dimensional Subalgebras and Corresponding Reductive Pairs of Lie Algebra of All Real $2 \times 2$ Matrices

Shtukar U\*

Associate Professor, Math/Physics Department, North Carolina Central University, Durham

## Abstract

The purpose of the article is to describe all 2-dimensional subalgebras and all corresponding reductive pairs of Lie algebra  $g$  of all  $2 \times 2$  real matrices. This Lie algebra is 4-dimensional as a vector space, it's not simple, and it's not solvable. The evaluation procedure utilizes canonical bases for subspaces that were introduced. Part I of the article contains necessary basic information. In Part II, all 2-dimensional subalgebras of the given Lie algebra  $g$  are classified. All reductive pairs  $\{h, m\}$  with 2-dimensional subalgebras  $h$  are found in Part III. The separate article contributes classification of all 3-dimensional subalgebras and its reductive pairs. Together, both articles give the total classification of all subalgebras and all reductive pairs of Lie algebra  $g$ .

**Keywords:** Lie algebra; Subalgebras; Reductive pairs

## Introduction

Katsumi Nomizu introduced reductive homogeneous spaces at his fundamental manuscript [1,2] where the author investigated invariant affine connections and Riemannian metrics on them. Sagle and Winter at their article [3] analyzed algebraic structures generated by reductive pairs of simple Lie algebras. One more problem that concerns to this article is classification of subalgebras of low dimensional Lie algebras. For example, Patera and Winternitz classified all subalgebras of real Lie algebras of dimensions  $d=3$  and  $d=4$  at the manuscript [4]. Their classification of subalgebras of real Lie algebras was done by a representative of each conjugacy class where the conjugacy was considered under the group of inner automorphisms of Lie algebras. All the articles mentioned above have stimulated this research of all 2-dimensional subalgebras and their reductive pairs of Lie algebra  $g$  of all real  $2 \times 2$  matrices. In contrast to the article [5], the current research is utilized a different method. Our method involves canonical bases for subspaces [1] that allow us to find all 2-dimensional subalgebras and the corresponding reductive pairs of the given Lie algebra  $g$ . This article finalizes the total classification of reductive pairs of the considering Lie algebra. New knowledge concerning the structure of Lie algebra  $g$  is important for Algebra, Geometry, and Physics.

## Part I. Basic information and necessary statement

We remind some information for the readers' convenience that includes the basic statement about canonical bases from the article [1].

**Definition 1:** Let  $g$  be Lie algebra,  $h$  be Lie subalgebra of  $g$ . If there exists a subspace  $m$  of  $g$  such that  $g=h \oplus m$  and  $[h, m] \subset m$ , then  $\{h, m\}$  is called a reductive pair of  $g$ , and  $\{g, h, m\}$  is called a reductive triple. We say also that subspace  $m$  is a reductive complement for  $h$  at  $g$ .

## Lie algebra $g$ and its standard basis

This Lie algebra contains all  $2 \times 2$  matrices over the field of all real numbers. The standard basis of this algebra consists of the next four matrices

$$\bar{e}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \bar{e}_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \bar{e}_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \bar{e}_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

It is well known that the Lie multiplication operation  $[A, B]$  for any two square matrices  $A$  and  $B$  of the same size is defined to be  $[A, B]=AB-BA$ . According this rule, the fundamental nonzero products of the basic vectors (matrices)  $\bar{e}_1, \bar{e}_2, \bar{e}_3, \bar{e}_4$  can be computed:

$$[\bar{e}_1, \bar{e}_2] = \bar{e}_2, [\bar{e}_1, \bar{e}_3] = -\bar{e}_3, [\bar{e}_2, \bar{e}_3] = \bar{e}_1 - \bar{e}_4, [\bar{e}_2, \bar{e}_4] = \bar{e}_2, [\bar{e}_3, \bar{e}_4] = -\bar{e}_3. (*)$$

All other products of basic vectors are zeros.

## Canonical bases for 2-dimensional subspaces of 4-dimensional vector space

Let  $h = Span\{\bar{a}, \bar{b}\}$  be any 2-dimensional subspace of a 4-dimensional vector space  $g$  generated by linearly independent vectors  $\bar{a} = a_1\bar{e}_1 + a_2\bar{e}_2 + a_3\bar{e}_3 + a_4\bar{e}_4$ , and  $\bar{b} = b_1\bar{e}_1 + b_2\bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4$ . According the article [1], all canonical bases for 2-dimensional subspaces  $h$  are:

- (1)  $\bar{a} = \bar{e}_1 + a_3\bar{e}_3 + a_4\bar{e}_4, \bar{b} = \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4$ ;
- (2)  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2 + a_4\bar{e}_4, \bar{b} = \bar{e}_3 + b_4\bar{e}_4$ ;
- (3)  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2 + a_3\bar{e}_3, \bar{b} = \bar{e}_4$ ; (4)  $\bar{a} = \bar{e}_2 + a_4\bar{e}_4, \bar{b} = \bar{e}_3 + b_4\bar{e}_4$ ;
- (5)  $\bar{a} = \bar{e}_2 + a_3\bar{e}_3, \bar{b} = \bar{e}_4$ ; (6)  $\bar{a} = \bar{e}_3, \bar{b} = \bar{e}_4$ .

Bases of the type (1) form 4-dimensional manifold, bases of the type (2) form 3-dimensional manifold, bases of the type (3) form 2-dimensional manifold, bases of the type (4) form another 2-dimensional manifold, bases of the type (5) form 1-dimensional manifold, and the basis (6) is a unique one. These manifolds are interested to be studied but the main goal of this article is different. About terminology: we will say just a basis (1)-(6) instead of a manifold of bases.

## Part II. 2-dimensional subalgebras of Lie algebra $g$

Now we start to determine when a 2-dimensional subspace

\*Corresponding author: Uladzimir Shtukar, Associate Professor, Math/Physics Department, North Carolina Central University, 1801 Fayetteville Street, Durham, NC 27707, USA, Tel: 919-597-0375, E-mail: vshtukar@yahoo.com

Received February 28, 2017; Accepted March 22, 2017; Published March 27, 2017

Citation: Shtukar U (2017) Classification of 2-Dimensional Subalgebras and Corresponding Reductive Pairs of Lie Algebra of All Real  $2 \times 2$  Matrices. J Generalized Lie Theory Appl 11: 260. doi:10.4172/1736-4337.1000260

Copyright: © 2017 Shtukar U. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$h = \text{Span}\{\bar{a}, \bar{b}\}$  is subalgebra of Lie algebra  $g$ . Obviously, the 6 nonequivalent canonical bases listed above should be used to analyze any subspace  $h$ . The condition  $[h, h] \subset h$  will be checked for each of these 6 canonical bases. The table (\*) of products from Part I will be used when a product  $[\bar{a}, \bar{b}]$  is computed.

Let  $\bar{a} = \bar{e}_1 + a_3\bar{e}_3 + a_4\bar{e}_4, \bar{b} = \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4$  be the basis (1) for  $h$ . Evaluate the product  $[\bar{a}, \bar{b}]$ . We have

$$[\bar{a}, \bar{b}] = [\bar{e}_1 + a_3\bar{e}_3 + a_4\bar{e}_4, \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4] = \bar{e}_2 - b_3\bar{e}_3 - a_3(\bar{e}_1 - \bar{e}_4) - a_3b_3\bar{e}_3 - a_4\bar{e}_2 + a_4b_3\bar{e}_3.$$

This product  $[\bar{a}, \bar{b}]$  should be located into the subspace  $h$ , i.e.  $[\bar{a}, \bar{b}] = x\bar{a} + y\bar{b}$ . So, we have the following conditions for  $x, y$  and for the components of  $\bar{a}$  and  $\bar{b}$ :

$$x = -a_3, y = 1 - a_4, xa_3 + yb_3 = -b_3 - a_3b_4 + a_4b_3, xa_4 + yb_4 = a_3.$$

The last 2 conditions generate the next system of 2 equations for  $a_3, a_4, b_3, b_4$ :

$$-a_3^2 + (1 - a_4)b_3 = -b_3 - a_3b_4 + a_4b_3, -a_3a_4 + (1 - a_4)b_4 = a_3, \\ \text{or } 2(1 - a_4)b_3 = a_3^2 - a_3b_4, (1 - a_4)b_4 = (1 + a_4)a_3.$$

To solve this system of equations, consider two cases:  $a_4 = 1$ , and  $a_4 \neq 1$ . If  $a_4 = 1$ , then  $a_3(a_3 - b_3) = 0, 2a_3 = 0$ . So,  $a_3 = 0, a_4 = 1$ , and  $b_3, b_4$  are any components. The first set of 2-dimensional subalgebras is:

$$h_1 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4\}.$$

If  $a_4 \neq 1$ , then  $b_4 = \frac{1 + a_4}{1 - a_4}a_3, b_3 = \frac{-a_4a_3^2}{(1 - a_4)^2}$ , and a new set of 2-dimensional subalgebras is:

$$h_2 = \text{Span}\{\bar{e}_1 + a_3\bar{e}_3 + a_4\bar{e}_4, \bar{e}_2 - \frac{a_4a_3^2}{(1 - a_4)^2}\bar{e}_3 + \frac{1 + a_4}{1 - a_4}a_3\bar{e}_4\}, a_4 \neq 1.$$

Let  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2 + a_4\bar{e}_4, \bar{b} = \bar{e}_3 + b_4\bar{e}_4$  be the basis (2) for  $h$ . Evaluate the product  $[\bar{a}, \bar{b}]$ :

$$[\bar{a}, \bar{b}] = [\bar{e}_1 + a_2\bar{e}_2 + a_4\bar{e}_4, \bar{e}_3 + b_4\bar{e}_4] = -\bar{e}_3 + a_2(\bar{e}_1 - \bar{e}_4) + a_2b_4\bar{e}_2 + a_4\bar{e}_3 = x\bar{a} + y\bar{b}.$$

This vector equality generates the next system of 2 equations for  $a_2, a_4, b_4$ :

$$a_2^2 = a_2b_4, a_2a_4 + (a_4 - 1)b_4 = -a_2.$$

The 2<sup>nd</sup> equation gives the result  $b_4 = \frac{1 + a_4}{1 - a_4}a_2 (a_4 \neq 1)$ , and the 1<sup>st</sup> equation produces  $b_4 = a_2$  or  $a_2 = 0$ . If  $a_2 = 0$  then  $b_4 = 0$ . If  $b_4 = a_2$  then  $a_4 = 0$ . If  $a_4 = 1$  then  $a_2 = 0$ . So, three new sets of 2-dimensional subalgebras are obtained:

$$h_3 = \text{Span}\{\bar{e}_1 + a_4\bar{e}_4, \bar{e}_3\} \quad (a_4 \neq 1), \quad h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\}, \\ h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}.$$

Let  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2 + a_3\bar{e}_3, \bar{b} = \bar{e}_4$  be the basis (3) for  $h$ . Compute the product  $[\bar{a}, \bar{b}]$ :

$$[\bar{a}, \bar{b}] = [\bar{e}_1 + a_2\bar{e}_2 + a_3\bar{e}_3, \bar{e}_4] = a_2\bar{e}_2 - a_3\bar{e}_3 = x\bar{a} + y\bar{b}.$$

The last vector equality generates the values  $x = 0, y = 0$ , and we have immediately  $a_2 = 0, a_3 = 0$ . This means that the following abelian subalgebra is obtained:

$$h_6 = \text{Span}\{\bar{e}_1, \bar{e}_4\}.$$

Let  $\bar{a} = \bar{e}_2 + a_4\bar{e}_4, \bar{b} = \bar{e}_3 + b_4\bar{e}_4$  be the basis (4) for  $h$ . Evaluate the product  $[\bar{a}, \bar{b}]$ , and determine when it belongs to  $h$ . We have:

$$[\bar{a}, \bar{b}] = [\bar{e}_2 + a_4\bar{e}_4, \bar{e}_3 + b_4\bar{e}_4] = (\bar{e}_1 - \bar{e}_4) + b_4\bar{e}_2 + a_4\bar{e}_3 = x\bar{a} + y\bar{b}.$$

The last vector equality is impossible because basic vector  $\bar{e}_1$  can't be generated by vectors  $\bar{b}$  and  $\bar{b}$ . This means that no subalgebra exists with the basis (4).

Let  $\bar{a} = \bar{e}_2 + a_3\bar{e}_3, \bar{b} = \bar{e}_4$  be the basis (5) for  $h$ . Evaluate the product  $[\bar{a}, \bar{b}]$ , and determine when it belongs to  $h$ . We have:

$$[\bar{a}, \bar{b}] = [\bar{e}_2 + a_3\bar{e}_3, \bar{e}_4] = \bar{e}_2 - a_3\bar{e}_3 = x\bar{a} + y\bar{b}.$$

The last vector equality is satisfied if and only if  $a_3 = 0$ . So, the following subalgebra is obtained  $h_7 = \text{Span}\{\bar{e}_2, \bar{e}_4\}$ .

Let  $\bar{a} = \bar{e}_3, \bar{b} = \bar{e}_4$  be the basis (6) for  $h$ . Evaluate the product  $[\bar{a}, \bar{b}]$ , and determine if it belongs to  $h$ . We have  $[\bar{a}, \bar{b}] = [\bar{e}_3, \bar{e}_4] = -\bar{e}_3 = -\bar{a} \in \text{Span}\{\bar{a}, \bar{b}\}$ .

So, the new 2-dimensional subalgebra  $h_8 = \text{Span}\{\bar{e}_3, \bar{e}_4\}$  of Lie algebra  $g$  is found.

The next theorem summarizes all results of Part II.

**Theorem 1:** Lie algebra  $g$  has two different 2-parameters sets of 2-dimensional subalgebras:

$$h_1 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4\};$$

$h_2 = \text{Span}\{\bar{e}_1 + a_3\bar{e}_3 + a_4\bar{e}_4, \bar{e}_2 - \frac{a_4a_3^2}{(1 - a_4)^2}\bar{e}_3 + \frac{1 + a_4}{1 - a_4}a_3\bar{e}_4\}, a_4 \neq 1$ ; three different 1-parameter sets of 2-dimensional subalgebras.

$h_3 = \text{Span}\{\bar{e}_1 + a_4\bar{e}_4, \bar{e}_3\}, a_4 \neq 1, h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\}, h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}$  and three special 2-dimensional subalgebras,

$$h_6 = \text{Span}\{\bar{e}_1, \bar{e}_4\}; \quad h_7 = \text{Span}\{\bar{e}_2, \bar{e}_4\}; \quad h_8 = \text{Span}\{\bar{e}_3, \bar{e}_4\}.$$

### Part III. Reductive pairs with 2-dimensional subalgebras of Lie algebra $g$

How many of 2-dimensional subalgebras  $h$  form reductive pairs  $\{h, m\}$  of Lie algebra  $g$ ? To answer this question, we will find all reductive complements  $m$  for each subalgebra  $h$  such that the conditions  $[h, m] \subset m$  and  $g = h \oplus m$  are satisfied. A complement  $m$  for any 2-dimensional subalgebra  $h$  should be a 2-dimensional subspace, and we can describe it as  $m = \text{Span}\{\bar{c}, \bar{d}\}$  where vectors  $\bar{c}$  and  $\bar{d}$  form some canonical basis for  $m$ . Remind all canonical bases for 2-dimensional subspaces:

- (1)  $\bar{c} = \bar{e}_1 + c_3\bar{e}_3 + c_4\bar{e}_4, \bar{d} = \bar{e}_2 + d_3\bar{e}_3 + d_4\bar{e}_4$ ;
- (2)  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_4\bar{e}_4, \bar{d} = \bar{e}_3 + d_4\bar{e}_4$ ;
- (3)  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_3\bar{e}_3, \bar{d} = \bar{e}_4$ ; (4)  $\bar{c} = \bar{e}_2 + c_4\bar{e}_4, \bar{d} = \bar{e}_3 + d_4\bar{e}_4$ ;
- (5)  $\bar{c} = \bar{e}_2 + c_3\bar{e}_3, \bar{d} = \bar{e}_4$ ; (6)  $\bar{c} = \bar{e}_3, \bar{d} = \bar{e}_4$ .

We start to utilize five sets of 2-dimensional subalgebras and three special 2-dimensional subalgebras listed in below Theorem 1.

**Theorem 1:** Subalgebra  $h_1 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_2 + b_3\bar{e}_3 + b_4\bar{e}_4\}$ .

Find all reductive complements for  $h_1$  if they exist.

Let  $\vec{c} = \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4$ ,  $\vec{d} = \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$ ,  $\vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$ , by  $\vec{c}$  and  $\vec{d}$ .

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4] = -c_3\vec{e}_3 + c_3\vec{e}_3 = \vec{0}; \text{ it is the identity.}$$

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4] = \vec{e}_2 - d_3\vec{e}_3 - \vec{e}_2 + d_3\vec{e}_3 = \vec{0}; \text{ it's}$$

the identity.

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4] = -\vec{e}_2 + c_3(\vec{e}_1 - \vec{e}_4) + c_4\vec{e}_2 + b_3\vec{e}_3 - b_3c_3\vec{e}_3 + b_4c_3\vec{e}_3 = x_3\vec{c} + y_3\vec{d}.$$

$$\text{So, } x_3 = c_3, y_3 = c_4 - 1, x_3c_3 + y_3d_3 = b_3 - b_3c_3 + b_4c_3, x_3c_4 + y_3d_4 = -c_3.$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4] = d_3(\vec{e}_1 - \vec{e}_4) + d_4\vec{e}_2 - b_3(\vec{e}_1 - \vec{e}_4) - b_3d_3\vec{e}_3 - b_4\vec{e}_2 + b_4d_3\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

$$\text{So, } x_4 = d_3 - b_3, y_4 = d_4 - b_4, x_4c_3 + y_4d_3 = -b_3d_4 + b_4d_3, x_4c_4 + y_4d_4 = b_3 - d_3.$$

The conditions above generate the following system of 4 nonlinear equations for components  $c_3, c_4, d_3, d_4$ :

$$c_3^2 + (c_4 - 1)d_3 = b_3 - b_3c_4 + b_4c_3, c_3c_4 + (c_4 - 1)d_4 = -c_3,$$

$$(d_3 - b_3)c_3 + (d_4 - b_4)d_3 = -b_3d_4 + b_4d_3, (d_3 - b_3)c_4 + (d_4 - b_4)d_4 = b_3 - d_3.$$

It makes sense to solve the system in two different cases:  $c_4=1$ , and  $c_4 \neq 1$ .

If  $c_4=1$ , then  $c_3=0$ ,  $(d_4 - b_4)d_3 = b_4d_3 - b_3d_4$ ,  $2(d_3 - b_3) = (b_4 - d_4)d_4$ . We see that  $\vec{c} = \vec{e}_1 + \vec{e}_4$ , and  $\vec{d}$  is some vector determined by the last two equations above. This pair  $\{h_1, m\}$  with  $m = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{d}\}$  is not a reductive pair because the intersection  $h_1 \cap m = \{\vec{e}_1 + \vec{e}_4\} \neq \vec{0}$  isn't zero vector for any  $\vec{d}$ .

$$\text{If } c_4 \neq 1, \text{ then } d_4 = \frac{1+c_4}{1-c_4}c_3, d_3 = \frac{c_3-b_4}{1-c_4}c_3 - b_3, d_3 = \frac{b_4-d_4}{c_4}d_4 + b_3$$

$$\text{and } (d_3 - b_3)c_3 + (d_4 - b_4)d_3 = b_4d_3 - b_3d_4.$$

Utilizing the 1<sup>st</sup> formula for  $d_4$ , we compare two different values for  $d_3$ . The corresponding procedure is long, we omit details. At the end of it, we obtain the following equation for  $c_3$ :

$$c_3^2 - b_4(1 - c_4)c_3 - b_3(1 - c_4)^2 = 0. (**)$$

Substitute now formulas for  $d_3$  and  $d_4$  obtained above into the 3<sup>rd</sup> equation that was not used before:  $(d_3 - b_3)c_3 + (d_4 - b_4)d_3 = b_4d_3 - b_3d_4$ . Simplifying step by step this equation and utilizing several times the equation (\*\*), we obtain the following result:  $c_3c_4=0$ . So,  $c_3=0$  or  $c_4=0$ .

If  $c_3=0$  then from the equation (\*\*) we obtain  $c_4=1$  or  $b_3=0$ . For  $c_4=1$  we have  $d_3=b_3$ ,  $d_4=0$ , and the following pair appears  $h_1 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4\}$ ,  $m = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3\vec{e}_3\}$ . This pair is not reductive because the intersection  $h_1 \cap m = \{\vec{e}_1 + \vec{e}_4\}$  is not zero vectors. For the case  $b_3=0$  we have  $d_3=0$ ,  $d_4=0$ . The corresponding pair for this case is  $h_1 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_4\vec{e}_4\}$ ,  $m_1 = \text{Span}\{\vec{e}_1 + c_4\vec{e}_4, \vec{e}_2\}$ . This pair is reductive if  $b_4 \neq 0$ ,  $c_4 \neq 1$ .

If  $c_4=0$  then  $d_4=c_3$ ,  $d_3=0$ , and  $c_3$  is the solution of the equation  $c_3^2 - b_4c_3 - b_3 = 0$ . We have the following reductive pair:

$$h_1 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4\}, m_2 = \text{Span}\{\vec{e}_1 + c_3\vec{e}_3, \vec{e}_2 + c_3\vec{e}_4\}, c_3 \neq b_4, \text{ and } c_3 \text{ is the solution of the equation } c_3^2 - b_4c_3 - b_3 = 0.$$

2. Let  $\vec{c} = \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4$ ,  $\vec{d} = \vec{e}_3 + d_4\vec{e}_4$ , be the basis (2). Multiply vectors  $\vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$\vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4 \text{ by } \vec{c} \text{ and } \vec{d}. \text{ We have:}$$

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4] = c_2\vec{e}_2 - c_2\vec{e}_2 = \vec{0}; \text{ it is the identity.}$$

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_3 + d_4\vec{e}_4] = -\vec{e}_3 + \vec{e}_3 = \vec{0}; \text{ it is the identity.}$$

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4] = -\vec{e}_2 + c_4\vec{e}_2 + b_3\vec{e}_3 - b_3c_2(\vec{e}_1 - \vec{e}_4) - b_3c_4\vec{e}_3 - b_4c_2\vec{e}_2 = x_3\vec{c} + y_3\vec{d}.$$

$$\text{So, } x_3 = -b_3c_2, y_3 = b_3 - b_3c_4, x_3c_2 = c_4 - 1 - b_4c_2, x_3c_4 + y_3d_4 = b_3c_2.$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_3 + d_4\vec{e}_4] = (\vec{e}_1 - \vec{e}_4) + d_4\vec{e}_2 - b_3d_4\vec{e}_3 + b_4\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

$$\text{So, } x_4 = 1, y_4 = b_4 - b_3d_4, x_4c_2 = d_4, x_4c_4 + y_4d_4 = -1.$$

We obtain the next system of 4 equations for the components  $c_2, c_4, d_4$ :

$$-b_3c_2^2 = c_4 - 1 - b_4c_2, -b_3c_2c_4 + b_3(1 - c_4)d_4 = b_3c_2, c_2 = b_3, c_4 + (b_4 - b_3d_4)d_4 = -1.$$

Analyzing this system of 4 equations, we obtain the following solution of it (details are omitted):  $d_4=c_2$ ,  $c_4=0$ , and  $c_2$  is the solution of the equation  $b_3c_2^2 - b_4c_2 - 1 = 0$ . The corresponding reductive pair is:

$$h_1 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4\}, m_3 = \text{Span}\{\vec{e}_1 + c_2\vec{e}_2, \vec{e}_3 + c_2\vec{e}_4\}, \text{ where } b_3c_2^2 - b_4c_2 - 1 = 0.$$

3. Let  $\vec{c} = \vec{e}_1 + c_2\vec{e}_2 + c_3\vec{e}_3$ ,  $\vec{d} = \vec{e}_4$ , be the basis (3). Multiply vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$ ,  $\vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$ , by  $\vec{c}$  and  $\vec{d}$ . We obtain the following products:

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_3\vec{e}_3] = c_2\vec{e}_2 - c_3\vec{e}_3 - c_2\vec{e}_2 + c_3\vec{e}_3 = \vec{0}; \text{ it is the identity.}$$

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_4] = \vec{0}; \text{ it is the identity.}$$

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_3\vec{e}_3] = -\vec{e}_2 + c_4\vec{e}_2 + b_3\vec{e}_3 - b_3c_2(\vec{e}_1 - \vec{e}_4) - b_3c_3\vec{e}_3 - b_4c_2\vec{e}_2 = x_3\vec{c} + y_3\vec{d}.$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_4] = \vec{e}_2 - b_3\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

The third and fourth equalities generate the following conditions:

$$x_3 = c_3 - b_3c_2, y_3 = b_3c_2 - c_3, x_3c_2 = -1 - b_4c_2, x_3c_3 = b_3 + b_4c_3, x_4 = 0, y_4 = 0, x_4c_2 = 1, x_4c_3 = -b_3.$$

These conditions produce the obvious contradiction:  $0=1$ . This means that no reductive pair for  $h_1$  exists when the complement has the basis (3).

Let  $\vec{c} = \vec{e}_2 + c_4\vec{e}_4$ ,  $\vec{d} = \vec{e}_3 + d_4\vec{e}_4$  be the basis (4). Multiply vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$ ,  $\vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$  by  $\vec{c}$  and  $\vec{d}$ . We have two identities  $[\vec{a}, \vec{c}] = \vec{0}$ ,  $[\vec{a}, \vec{d}] = \vec{0}$ , and

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_2 + c_4\vec{e}_4] = c_4\vec{e}_2 - b_3(\vec{e}_1 - \vec{e}_4) - b_3c_4\vec{e}_3 - b_4\vec{e}_2 = x_3\vec{c} + y_3\vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_3 + d_4\vec{e}_4] = (\vec{e}_1 - \vec{e}_4) + d_4\vec{e}_2 - b_3d_4\vec{e}_3 + b_4\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

The product  $\vec{c}, \vec{d}$  can't be generated by vectors  $\vec{c}, \vec{d}$ . So, no reductive complement with basis (4) exists for subalgebra  $h_1$ .

Let  $\vec{c} = \vec{e}_2 + c_3\vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (5). Multiply vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4, \vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$ , by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_2 + c_3\vec{e}_3] = \vec{e}_2 - c_3\vec{e}_3 - \vec{e}_2 + c_3\vec{e}_3 = \vec{0}; \text{ it is the identity.}$$

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_4] = \vec{0}; \text{ it is the identity.}$$

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_2 + c_3\vec{e}_3] = c_3(\vec{e}_1 - \vec{e}_4) - b_3(\vec{e}_1 - \vec{e}_4) + b_3c_3\vec{e}_3 - b_4\vec{e}_2 = x_3\vec{c} + y_3\vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_4] = \vec{e}_2 - b_3\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

The solution for those vector equations is  $c_3=0, b_3=0$ . As the result, the following pair  $h_1 = Span\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_4\vec{e}_4\}, m = Span\{\vec{e}_2, \vec{e}_4\}$  is obtained but it is not a reductive pair because  $\vec{c} = \vec{e}_3, \vec{d} = \vec{e}_4$

6. Let  $\vec{c} = \vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (6). Multiply vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4, \vec{b} = \vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4$  by  $\vec{c}$  and  $\vec{d}$ . We have two identities  $[\vec{a}, \vec{c}] = \vec{0}, [\vec{a}, \vec{d}] = \vec{0}$ , and

$$[\vec{b}, \vec{c}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_3] = (\vec{e}_1 - \vec{e}_4) + b_4\vec{e}_3 = x_3\vec{c} + y_3\vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 + b_3\vec{e}_3 + b_4\vec{e}_4, \vec{e}_4] = \vec{e}_2 - b_3\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

The products  $[\vec{b}, \vec{c}]$  and  $[\vec{b}, \vec{d}]$  are not generated by vectors  $\vec{c}, \vec{d}$ . This means that no reductive complement with the basis (6) exists for subalgebra  $h_1$ .

Sub algebra  $h_2 = Span\{\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4\}, a_4 \neq 1$ .

Find reductive complements for  $h_2$  if they exist.

Let  $\vec{c} = \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4, \vec{d} = \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a}, \vec{b}$  by  $\vec{c}, \vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4] = -c_3\vec{e}_3 + a_3\vec{e}_3 - a_3c_3\vec{e}_3 + a_4c_3\vec{e}_3 = x_3\vec{c} + y_3\vec{d}.$$

So,  $x_1=0, y_1=0, x_1c_3+y_1d_3=a_3-c_3-a_3c_3+a_4c_3, x_1c_4+y_1d_4=0$ .

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4] = \vec{e}_2 - d_3\vec{e}_3 - a_3(\vec{e}_1 - \vec{e}_4) - a_4d_3\vec{e}_3 - a_4\vec{e}_2 + a_4d_3\vec{e}_3 = x_2\vec{c} + y_2\vec{d}.$$

So,  $x_2=-a_3, y_2=a_4, x_2c_3+y_2d_3=-d_3-a_3d_3+a_4d_3, x_2c_4+y_2d_4=a_3$ .

$$[\vec{b}, \vec{c}] = [\vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4, \vec{e}_1 + c_3\vec{e}_3 + c_4\vec{e}_4] = -\vec{e}_2 + c_3(\vec{e}_1 - \vec{e}_4) + c_4\vec{e}_4 -$$

$$\frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{a_4a_3^2}{(1-a_4)^2}c_4\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3c_3\vec{e}_3 = x_3\vec{c} + y_3\vec{d}.$$

So,  $x_3=c_3, y_3=c_4-1, x_3c_3+y_3d_3 = \frac{a_4a_3^2}{(1-a_4)^2}(c_4-1) + \frac{1+a_4}{1-a_4}a_3c_3,$

$x_3c_4+y_3d_4=-c_3,$

$$[\vec{b}, \vec{d}] = [\vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4, \vec{e}_2 + d_3\vec{e}_3 + d_4\vec{e}_4] = d_3(\vec{e}_1 - \vec{e}_4) + d_4\vec{e}_2 +$$

$$\frac{a_4a_3^2}{(1-a_4)^2}d_4\vec{e}_3 + \frac{a_4a_3^2}{(1-a_4)^2}(\vec{e}_1 - \vec{e}_4) - \frac{1+a_4}{1-a_4}a_3\vec{e}_2 + \frac{1+a_4}{1-a_4}a_3d_3\vec{e}_3 = x_4\vec{c} + y_4\vec{d}.$$

$x_4 = d_3 + \frac{a_4a_3^2}{(1-a_4)^2}, y_4 = d_4 - \frac{1+a_4}{1-a_4}a_3, x_4c_3+y_4d_3 = \frac{a_4a_3^2d_4}{(1-a_4)^2} + \frac{1+a_4}{1-a_4}a_3d_3,$

So,  $x_4c_4+y_4d_4 = -d_3 - \frac{a_4a_3^2}{(1-a_4)^2}$

From the equalities above, we obtain the following system of 7 nonlinear equations for components  $c_3, c_4, d_3, d_4$ :

$$a_3-c_3-a_3c_4+a_4c_3=0, \quad -a_3c_3+(1-a_4)d_3=-d_3-a_3d_4+a_4d_3, \quad -a_3c_4+(1-a_4)d_4=a_3,$$

$$c_3c_4+(c_4-1)d_4=-c_3, \quad c_3c_3+(c_4-1)d_3 = \frac{a_4a_3^2}{(1-a_4)^2}(c_4-1) + \frac{1+a_4}{1-a_4}a_3c_3,$$

$$[d_3 + \frac{a_4a_3^2}{(1-a_4)^2}]c_3 + (d_4 - \frac{1+a_4}{1-a_4}a_3)d_3 = \frac{a_4a_3^2}{(1-a_4)^2}d_4 + \frac{1+a_4}{1-a_4}a_3d_3,$$

$$[d_3 + \frac{a_4a_3^2}{(1-a_4)^2}]c_4 + (d_4 - \frac{1+a_4}{1-a_4}a_3)d_4 = -d_3 - \frac{a_4a_3^2}{(1-a_4)^2}.$$

The 1<sup>st</sup> equation gives us  $c_3 = \frac{1-c_4}{1-a_4}a_3$ , the 2<sup>nd</sup> equation gives  $d_3 = \frac{a_3(c_3-d_4)}{2(1-a_4)}$ , the 3<sup>rd</sup> equation gives  $d_4 = \frac{1+c_4}{1-a_4}a_3$ , and the 4<sup>th</sup> equation gives  $d_4 = \frac{1+c_4}{1-c_4}c_3$ . Substitute the values found for  $c_3, d_3, d_4$  into

the 5<sup>th</sup> equation, and simplify the corresponding expression. We obtain the identity  $1-c_4=1-c_4$ . Substituting the values found for  $c_3, d_3, d_4$  into the 6<sup>th</sup> equation, we obtain the identity  $0=0$  as well. Substitute the same values for  $c_3, d_3, d_4$  into the 7<sup>th</sup> equation. Simplifying the corresponding expression, we obtain the equation  $2c_4^2+c_4(1-a_4)=0$ . This equation has two solutions for  $c_4$ :  $c_4=0$  or  $c_4 = \frac{a_4-1}{2}$ . So, we obtain two

pairs for subalgebra  $h_2$ . The first pair is  $\{h_2, m_1\}$  where:

$$h_2 = Span\{\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4\}, a_4 \neq 1,$$

$m_1 = Span\{\vec{e}_1 + \frac{a_3}{1-a_4}\vec{e}_3, \vec{e}_2 + \frac{a_3}{1-a_4}\vec{e}_4\}$ . This pair is reductive if  $a_3 \neq 0, a_4 \neq 0$ .

The second pair is  $\{h_2, m_2\}$  where:

$$h_2 = Span\{\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4\}, a_4 \neq 1,$$

$$m_2 = Span\{\vec{e}_1 + \frac{3-a_4}{2(1-a_4)}a_3\vec{e}_3 + \frac{a_4-1}{2}\vec{e}_4, \vec{e}_2 + \frac{a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{2(1-a_4)}a_3\vec{e}_4\}.$$

This pair is reductive if  $a_3 \neq 0, a_4 \neq 1$ .

2. Let  $\vec{c} = \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4, \vec{d} = \vec{e}_3 + d_4\vec{e}_4$  be the basis (2) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a}, \vec{b}$  from  $h_2$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4] = c_2\vec{e}_2 + a_3\vec{e}_3 - a_3c_2(\vec{e}_1 - \vec{e}_4) - a_3c_2\vec{e}_3 - a_4c_2\vec{e}_2 = x_1\vec{c} + y_1\vec{d}.$$

So,  $x_1=-a_3c_2, y_1=a_3-a_3c_4, x_1c_2=c_2-a_4c_2, x_1c_4+y_1d_4=a_3c_2,$

$$[\vec{a}, \vec{d}] = [\vec{e}_1 + a_3\vec{e}_3 + a_4\vec{e}_4, \vec{e}_3 + d_4\vec{e}_4] = -\vec{e}_3 - a_3d_4\vec{e}_3 + a_4\vec{e}_3 = x_2\vec{c} + y_2\vec{d}.$$

So,  $x_2=0, y_2=a_4-a_3d_4-1, x_2c_2=0, x_2c_4+y_2d_4=0.$

$$[\vec{b}, \vec{c}] = [\vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 + \frac{1+a_4}{1-a_4}a_3\vec{e}_4, \vec{e}_1 + c_2\vec{e}_2 + c_4\vec{e}_4] = -\vec{e}_2 + c_4\vec{e}_2 - \frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 +$$

$$c_2\frac{a_4a_3^2}{(1-a_4)^2}(\vec{e}_1 - \vec{e}_4) + c_4\frac{a_4a_3^2}{(1-a_4)^2}\vec{e}_3 - \frac{1-a_4}{1-a_4}a_3c_2\vec{e}_2 = x_3\vec{c} + y_3\vec{d}.$$

$$\text{So, } x_3 = \frac{a_4 a_3^2}{(1-a_4)^2} c_2, y_3 = \frac{a_4 a_3^2}{(1-a_4)^2} (c_4 - 1), x_3 c_2 = c_4 - 1 - \frac{1+a_4}{1-a_4} a_3 c_2,$$

$$x_3 c_4 + y_3 d_4 = -\frac{a_4 a_3^2}{(1-a_4)^2} c_2$$

$$[\bar{b}, \bar{d}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_3 + d_4 \bar{e}_4 \right] = (\bar{e}_1 - \bar{e}_4) + d_4 \bar{e}_2 + d_4 \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_3 = x_4 \bar{c} + y_4 \bar{d}.$$

$$\text{So, } x_4 = 1, y_4 = \frac{a_4 a_3^2}{(1-a_4)^2} d_4 + \frac{1+a_4}{1-a_4} a_3, x_4 c_2 = d_4, x_4 c_4 + y_4 d_4 = -1.$$

These equalities produce the system of 7 nonlinear equations for components  $c_2, c_4, d_4$ :

$$-a_3 c_2^2 = c_2(1-a_4), -a_3 c_2 c_4 + (a_3 - a_3 c_4) d_4 = a_3 c_2, (a_4 - a_3 d_4 - 1) d_4 = 0,$$

$$\frac{a_4 a_3^2}{(1-a_4)^2} c_2^2 = c_4 - 1 - \frac{1+a_4}{1-a_4} a_3 c_2,$$

$$\frac{a_4 a_3^2}{(1-a_4)^2} c_2 c_4 + (c_4 - 1) \frac{a_4 a_3^2}{(1-a_4)^2} d_4 = -\frac{a_4 a_3^2}{(1-a_4)^2} c_2,$$

$$d_4 = c_2, c_4 + \frac{a_4 a_3^2}{(1-a_4)^2} d_4^2 + \frac{1+a_4}{1-a_4} a_3 d_4 = -1.$$

Consider two cases for the system. If  $c_2=0$  then  $c_4=1, d_4=0$ , and from the last equation  $c_4=-1$ . We obtain a contradiction, so the system has no solution at this case.

Let  $c_2 \neq 0$ . Then from the 1<sup>st</sup> equation we have  $c_2 = \frac{a_4 - 1}{a_3}$ , and from the 2<sup>nd</sup> equation we obtain  $a_3 c_4 = 0$ . The last equation gives two possible results:  $a_3=0$  or  $c_4=0$ . If  $a_3=0$  then from the 4<sup>th</sup> and 7<sup>th</sup> equations we obtain the contradiction  $c_4=1, c_4=-1$  again. So,  $c_4=0$ . Substitute the values of  $c_2, d_4, c_4$  into the 4<sup>th</sup> equation, the 5<sup>th</sup> equation, and the 7<sup>th</sup> equation. We obtain the identities at all these cases. So, the following reductive pair appears:

$$h_2 = \text{Span}\{\bar{e}_1 + a_3 \bar{e}_3 + a_4 \bar{e}_4, \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4\}, a_4 \neq 1,$$

$$m_3 = \text{Span}\{\bar{e}_1 + \frac{a_4 - 1}{a_3} \bar{e}_2, \bar{e}_3 + \frac{a_4 - 1}{a_3} \bar{e}_4\}, a_3 \neq 0.$$

3. Let  $\bar{c} = \bar{e}_1 + c_2 \bar{e}_2 + c_3 \bar{e}_3, \bar{d} = \bar{e}_4$  be the basis (3). Multiply basic vectors  $\bar{a}, \bar{b}$  by  $\bar{c}$  and  $\bar{d}$ .

One product is very important for this case. Compute  $[\bar{b}, \bar{d}]$ :

$$[\bar{b}, \bar{d}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_4 \right] = \bar{e}_2 + \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 = x_4 \bar{c} + y_4 \bar{d}.$$

$$\text{So, } x_4 = 0, y_4 = 0, x_4 c_2 = 1, x_4 c_3 = \frac{a_4 a_3^2}{(1-a_4)^2}.$$

This product  $[\bar{b}, \bar{d}]$  produces the obvious contradiction  $x_4=0, x_4 c_2=1$ . So, subalgebra  $h_2$  has no reductive complement with basis (3).

4. Let  $\bar{c} = \bar{e}_2 + c_4 \bar{e}_4, \bar{d} = \bar{e}_3 + d_4 \bar{e}_4$  be the basis (4). Multiply vector  $\bar{b}$  from  $h_2$  and vector  $\bar{d}$ . We have:

$$[\bar{b}, \bar{d}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_3 + d_4 \bar{e}_4 \right] = (\bar{e}_1 - \bar{e}_4) + d_4 \bar{e}_2 + \frac{a_4 a_3^2}{(1-a_4)^2} d_4 \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_3 = x_4 \bar{c} + y_4 \bar{d}.$$

This product  $[\bar{b}, \bar{d}]$  contains vector  $\bar{e}_1$  which can't be generated by vectors  $\bar{c}$  and  $\bar{d}$ . This means that  $h_2$  has no reductive complement with basis (4).

5. Let  $\bar{c} = \bar{e}_2 + c_3 \bar{e}_3, \bar{d} = \bar{e}_4$  be the basis (5). Multiply basic vectors  $\bar{b}, \bar{a}$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_3 \bar{e}_3 + a_4 \bar{e}_4, \bar{e}_2 + c_3 \bar{e}_3] = \bar{e}_2 - c_3 \bar{e}_3 - a_3(\bar{e}_1 - \bar{e}_4) - a_4 \bar{e}_2 + a_3 c_3 \bar{e}_3 = x_1 \bar{c} + y_1 \bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_3 \bar{e}_3 + a_4 \bar{e}_4, \bar{e}_4] = -a_3 \bar{e}_3 = x_2 \bar{c} + y_2 \bar{d},$$

$$[\bar{b}, \bar{c}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_2 + c_3 \bar{e}_3 \right] = c_3(\bar{e}_1 - \bar{e}_4) + \frac{a_4 a_3^2}{(1-a_4)^2} (\bar{e}_1 - \bar{e}_4) - \frac{a_3(1+a_4)}{1-a_4} \bar{e}_2 + \frac{1+a_4}{1-a_4} a_3 c_3 \bar{e}_3.$$

$$[\bar{b}, \bar{d}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_4 \right] = \bar{e}_2 + \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 = x_4 \bar{c} + y_4 \bar{d}.$$

Vector equalities  $[\bar{a}, \bar{c}]$  and  $[\bar{b}, \bar{c}]$  give immediately  $a_3=0, c_3=0$ . So, we obtain the following pair  $h_2 = \text{Span}\{\bar{e}_1 + a_4 \bar{e}_4, \bar{e}_2\}, m = \text{Span}\{\bar{e}_2, \bar{e}_4\}$  which is not reductive because  $h_2 \cap m \neq \bar{0}$ .

6. Let  $\bar{c} = \bar{e}_3, \bar{d} = \bar{e}_4$  be the basis (6). Multiply vectors  $\bar{a}, \bar{b}$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_3 \bar{e}_3 + a_4 \bar{e}_4, \bar{e}_3] = -\bar{e}_3 + a_4 \bar{e}_3 = x_1 \bar{c} + y_1 \bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_3 \bar{e}_3 + a_4 \bar{e}_4, \bar{e}_4] = -a_3 \bar{e}_3 = x_2 \bar{c} + y_2 \bar{d},$$

$$[\bar{b}, \bar{c}] = \left[ \bar{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \bar{e}_3 + \frac{1+a_4}{1-a_4} a_3 \bar{e}_4, \bar{e}_3 \right] = (\bar{e}_1 - \bar{e}_4) + \frac{1+a_4}{1-a_4} a_3 \bar{e}_3 = x_3 \bar{c} + y_3 \bar{d}.$$

The product  $[\bar{b}, \bar{c}]$  contains vector  $\bar{e}_1$  which can't be generated by vectors  $\bar{c}$  and  $\bar{d}$ . This means that  $h_2$  has no reductive complement with basis (6).

Subalgebra  $h_3 = \text{Span}\{\bar{e}_1 + a_4 \bar{e}_4, \bar{e}_3\}$ . This subalgebra has no reductive complement at all. The corresponding evaluation procedure is long as that for subalgebra  $h_1$ , therefore it's omitted.

$$\text{Subalgebra } h_4 = \text{Span}\{\bar{e}_1 + a_2 \bar{e}_2, \bar{e}_3 + a_2 \bar{e}_4\}.$$

Find reductive complements for  $h_4$  if they exist.

1. Let  $\bar{c} = \bar{e}_1 + c_3 \bar{e}_3 + c_4 \bar{e}_4, \bar{d} = \bar{e}_2 + d_3 \bar{e}_3 + d_4 \bar{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2 \bar{e}_2, \bar{b} = \bar{e}_3 + a_2 \bar{e}_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2 \bar{e}_2, \bar{e}_1 + c_3 \bar{e}_3 + c_4 \bar{e}_4] = -c_3 \bar{e}_3 - a_2 \bar{e}_2 + a_2 c_3 (\bar{e}_1 - \bar{e}_4) + a_2 c_4 \bar{e}_2 = x_1 \bar{c} + y_1 \bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2 \bar{e}_2, \bar{e}_2 + d_3 \bar{e}_3 + d_4 \bar{e}_4] = \bar{e}_2 - d_3 \bar{e}_3 + a_2 d_3 (\bar{e}_1 - \bar{e}_4) + a_2 d_4 \bar{e}_2 = x_2 \bar{c} + y_2 \bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2 \bar{e}_4, \bar{e}_1 + c_3 \bar{e}_3 + c_4 \bar{e}_4] = \bar{e}_3 - c_4 \bar{e}_3 + a_2 c_3 \bar{e}_3 = x_3 \bar{c} + y_3 \bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2 \bar{e}_4, \bar{e}_2 + d_3 \bar{e}_3 + d_4 \bar{e}_4] = -(\bar{e}_1 - \bar{e}_4) - d_4 \bar{e}_3 - a_2 \bar{e}_2 + a_2 d_3 \bar{e}_3 = x_4 \bar{c} + y_4 \bar{d}.$$

These vector equalities produce the following system of equations:

$$a_2 c_3 c_3 + a_2 d_3 (c_4 - 1) = -c_3, a_2 c_3 c_4 + a_2 d_3 (c_4 - 1) = -a_2 c_3, a_2 c_3 d_3 + (1 + a_2 d_4) d_3 = -d_3,$$

$$a_2 d_3 c_4 + (1 + a_2 d_4) d_4 = -a_2 d_3, c_4 = 1 + a_2 c_3, -a_2 d_3 c_3 = a_2 d_3 - d_4, c_4 = -1 - a_2 d_4.$$

This system of equations has the solution:  $c_4=1+a_2c_3$ ,  $d_3=-\frac{a_2c_3+1}{a_2}$ ,  $d_4=-\frac{a_2c_3+2}{a_2}$ ,  $a_2 \neq 0$ . If  $a_2=0$  then the system of equations generates the contradiction  $c_4=1$  and  $c_4=-1$ . So, we obtain one new reductive pair:

$$h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\},$$

$$m_1 = \text{Span}\{\bar{e}_1 + c_3\bar{e}_3 + (1+a_2c_3)\bar{e}_4, \bar{e}_2 - \frac{a_2c_3+1}{a_2}\bar{e}_3 - \frac{a_2c_3+2}{a_2}\bar{e}_4\}.$$

2. Let  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_4\bar{e}_4$ ,  $\bar{d} = \bar{e}_3 + d_4\bar{e}_4$  be the basis (2) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2$ ,  $\bar{b} = \bar{e}_3 + a_2\bar{e}_4$  of  $h_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_1 + c_2\bar{e}_2 + c_4\bar{e}_4] = c_2\bar{e}_2 - a_2\bar{e}_2 + a_2c_4\bar{e}_4 = x_1\bar{c} + y_1\bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + d_4\bar{e}_4] = -\bar{e}_3 + a_2(\bar{e}_1 - \bar{e}_4) + a_2d_4\bar{e}_2 = x_2\bar{c} + y_2\bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_1 + c_2\bar{e}_2 + c_4\bar{e}_4] = \bar{e}_3 - c_2(\bar{e}_1 - \bar{e}_4) - c_4\bar{e}_3 - a_2c_2\bar{e}_3 = x_3\bar{c} + y_3\bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_3 + d_4\bar{e}_4] = -d_4\bar{e}_3 + a_2\bar{e}_3 = x_4\bar{c} + y_4\bar{d}.$$

The system of equations has two solutions:  $a_2=0$ ,  $c_2=0$ ,  $d_4=0$ , and  $a_2 \neq 0$ ,  $c_2=0$ ,  $d_4=0$ ,  $c_4=1$ . For the first case, the corresponding pair  $h_4 = \text{Span}\{\bar{e}_1, \bar{e}_3\}$ ,

$m = \text{Span}\{\bar{e}_1 + c_4\bar{e}_4, \bar{e}_3\}$  is not reductive because  $h_4 \cap m \neq \bar{0}$ . For the second case, the corresponding pair  $h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\}$ ,  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_3\bar{e}_3$  is reductive if  $a_2 \neq 0$ .

3. Let  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_3\bar{e}_3$ ,  $\bar{d} = \bar{e}_4$  be the basis (3) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2$ ,  $\bar{b} = \bar{e}_3 + a_2\bar{e}_4$  of  $h_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_1 + c_2\bar{e}_2 + c_3\bar{e}_3] = c_2\bar{e}_2 - c_3\bar{e}_3 - a_2\bar{e}_2 + a_2c_3(\bar{e}_1 - \bar{e}_4) = x_1\bar{c} + y_1\bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_4] = a_2\bar{e}_2 = x_2\bar{c} + y_2\bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_1 + c_2\bar{e}_2 + c_3\bar{e}_3] = \bar{e}_3 - c_2(\bar{e}_1 - \bar{e}_4) + a_2c_3\bar{e}_3 - a_2c_2\bar{e}_2 = x_3\bar{c} + y_3\bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_4] = -\bar{e}_3 = x_4\bar{c} + y_4\bar{d}, \text{ where } x_4=0, y_4=0.$$

The last vector equality generates a contradiction  $\bar{e}_3 = \bar{0}$ , so subalgebra  $h_4$  has no reductive complement with basis (3).

4. Let  $\bar{c} = \bar{e}_2 + c_4\bar{e}_4$ ,  $\bar{d} = \bar{e}_3 + d_4\bar{e}_4$  be the basis (4) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2$ ,  $\bar{b} = \bar{e}_3 + a_2\bar{e}_4$  of  $h_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_2 + c_4\bar{e}_4] = \bar{e}_2 + a_2c_4\bar{e}_2 = x_1\bar{c} + y_1\bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + d_4\bar{e}_4] = -\bar{e}_3 + a_2(\bar{e}_1 - \bar{e}_4) + a_2d_4\bar{e}_2 = x_2\bar{c} + y_2\bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_2 + c_4\bar{e}_4] = -(\bar{e}_1 - \bar{e}_4) - c_4\bar{e}_3 - a_2\bar{e}_2 = x_3\bar{c} + y_3\bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_3 + d_4\bar{e}_4] = -d_4\bar{e}_3 + a_2\bar{e}_4 = x_4\bar{c} + y_4\bar{d}.$$

The product  $[\bar{b}, \bar{c}]$  contains vector  $\bar{e}_1$  that can't be generated by  $\bar{c}$  and  $\bar{d}$ , so subalgebra  $h_4$  has no reductive complement with basis (4).

5. Let  $\bar{c} = \bar{e}_2 + c_3\bar{e}_3$ ,  $\bar{d} = \bar{e}_4$  be the basis (5) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2$ ,  $\bar{b} = \bar{e}_3 + a_2\bar{e}_4$  of  $h_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_2 + c_3\bar{e}_3] = \bar{e}_2 - c_3\bar{e}_3 + a_2c_3(\bar{e}_1 - \bar{e}_4) = x_1\bar{c} + y_1\bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_4] = a_2\bar{e}_2 = x_2\bar{c} + y_2\bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_2 + c_3\bar{e}_3] = -(\bar{e}_1 - \bar{e}_4) + a_2c_3\bar{e}_3 - a_2\bar{e}_2 = x_3\bar{c} + y_3\bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_4] = -\bar{e}_3 = x_4\bar{c} + y_4\bar{d}.$$

The product  $[\bar{b}, \bar{c}]$  contains vector  $\bar{e}_1$  that can't be generated by vectors  $\bar{c}, \bar{d}$ , so subalgebra  $h_4$  has no reductive complement with basis (5).

6. Let  $\bar{c} = \bar{e}_3$ ,  $\bar{d} = \bar{e}_4$  be the basis (6) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + a_2\bar{e}_2$ ,  $\bar{b} = \bar{e}_3 + a_2\bar{e}_4$  of  $h_4$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3] = -\bar{e}_3 + a_2(\bar{e}_1 - \bar{e}_4) = x_1\bar{c} + y_1\bar{d},$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + a_2\bar{e}_2, \bar{e}_4] = a_2\bar{e}_2 = x_2\bar{c} + y_2\bar{d},$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_3] = a_2\bar{e}_3 = x_3\bar{c} + y_3\bar{d},$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + a_2\bar{e}_4, \bar{e}_4] = -\bar{e}_3 = x_4\bar{c} + y_4\bar{d}.$$

The system of vector equations has the solution  $a_2=0$ . The corresponding pair  $h_4 = \text{Span}\{\bar{e}_1, \bar{e}_3\}$ ,  $m = \text{Span}\{\bar{e}_3, \bar{e}_4\}$  is not reductive because  $h_4 \cap m \neq \bar{0}$ .

Subalgebra  $h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}$ .

Find reductive complements for  $h_5$  if they exist.

Let  $\bar{c} = \bar{e}_1 + c_3\bar{e}_3 + c_4\bar{e}_4$ ,  $\bar{d} = \bar{e}_2 + d_3\bar{e}_3 + d_4\bar{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\bar{a} = \bar{e}_1 + \bar{e}_4$  and  $\bar{b} = \bar{e}_3 + b_4\bar{e}_4$  of  $h_5$  by  $\bar{c}$  and  $\bar{d}$ . We have:

$$[\bar{a}, \bar{c}] = [\bar{e}_1 + \bar{e}_4, \bar{e}_1 + c_3\bar{e}_3 + c_4\bar{e}_4] = -c_3\bar{e}_3 + c_3\bar{e}_3 = \bar{0}, \text{ it is the identity;}$$

$$[\bar{b}, \bar{c}] = [\bar{e}_3 + b_4\bar{e}_4, \bar{e}_1 + c_3\bar{e}_3 + c_4\bar{e}_4] = \bar{e}_3 - c_4\bar{e}_3 + b_4c_3\bar{e}_3 = x_3\bar{c} + y_3\bar{d}, \text{ it is the identity;}$$

$$[\bar{b}, \bar{d}] = [\bar{e}_3 + b_4\bar{e}_4, \bar{e}_2 + d_3\bar{e}_3 + d_4\bar{e}_4] = \bar{e}_3 - c_4\bar{e}_3 + b_4c_3\bar{e}_3 = x_3\bar{c} + y_3\bar{d},$$

$$x_3=0, y_3=0;$$

$$[\bar{a}, \bar{d}] = [\bar{e}_1 + \bar{e}_4, \bar{e}_2 + d_3\bar{e}_3 + d_4\bar{e}_4] = -(\bar{e}_1 - \bar{e}_4) - d_4\bar{e}_3 - b_4\bar{e}_2 + b_4d_3\bar{e}_3 = x_4\bar{c} + y_4\bar{d}.$$

The solution for the system of vector equalities is  $c_4=1+b_4c_3$ ,  $d_3=-\frac{1+b_4c_3}{b_4}$ ,  $d_4=-\frac{2+b_4c_3}{b_4}$ ,  $b_4 \neq 0$ . This solution produces the pair  $h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}$ , but it is not reductive because  $h_5 \cap m = \{\bar{e}_1 + c_3\bar{e}_3 + (1+b_4c_3)\bar{e}_4\} \neq \bar{0}$ . If  $b_4=0$  then the system of equalities has no solution because the contradiction  $c_4=1$ ,  $c_4=-1$  appears.

Let  $\bar{c} = \bar{e}_1 + c_2\bar{e}_2 + c_4\bar{e}_4$ ,  $\bar{a} = \bar{e}_1 + \bar{e}_4$  be the basis (2) for a possible

reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$  and  $\vec{b} = \vec{e}_3 + b_4 \vec{e}_4$  of  $h_5$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$\begin{aligned} [\vec{a}, \vec{C}] &= [\vec{e}_1 + \vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = c_2 \vec{e}_2 - c_2 \vec{e}_2 = \vec{0} \text{ - it is the identity;} \\ [\vec{a}, \vec{D}] &= [\vec{e}_1 + \vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_3 + \vec{e}_3 = \vec{0} \text{ - it is the identity;} \\ [\vec{b}, \vec{C}] &= [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = \vec{e}_3 - c_2(\vec{e}_1 - \vec{e}_4) - c_4 \vec{e}_3 - b_4 c_2 \vec{e}_2 = x_3 \vec{c} + y_3 \vec{d}; \\ [\vec{b}, \vec{D}] &= [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = -d_4 \vec{e}_3 + b_4 \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}. \end{aligned}$$

These vector equalities generate the following system of equations:

$$c_2(b_4 - c_2) = 0, (1 - c_4)d_4 = (1 + c_4)c_2, d_4(b_4 - d_4) = 0.$$

The system has four different solutions:  $c_2=0, d_4=0; c_2=0, c_4=1, d_4=b_4;$

$$c_2=b_4, c_4=1, d_4=0; c_2=b_4, c_4=0, d_4=b_4.$$

These solutions produce the following pairs:  $h_5 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\}$ ,  $m_1 = \text{Span}\{\vec{e}_1 + c_4 \vec{e}_4, \vec{e}_3\}; h_5 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\}$ ,

$$m = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\}; h_5 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\},$$

$$m_2 = \text{Span}\{\vec{e}_1 + b_4 \vec{e}_2 - \vec{e}_4, \vec{e}_3\};$$

$$h_5 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\}, m = \text{Span}\{\vec{e}_1 + b_4 \vec{e}_2, \vec{e}_3 + b_4 \vec{e}_4\}.$$

Two pairs  $\{h_5, m_1\}$  and  $\{h_5, m_2\}$  are reductive if  $b_4 \neq 0$ , other pairs are not reductive.

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (3) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$  and  $\vec{b} = \vec{e}_3 + b_4 \vec{e}_4$  of  $h_5$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$[\vec{a}, \vec{C}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3] = c_2 \vec{e}_2 - c_2 \vec{e}_2 - c_3 \vec{e}_3 + c_3 \vec{e}_3 = \vec{0} \text{ - it is the identity;}$$

$$[\vec{a}, \vec{D}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_4] = \vec{0} \text{ - it is the identity;}$$

$$[\vec{b}, \vec{C}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3] = \vec{e}_3 - c_2(\vec{e}_1 - \vec{e}_4) - b_4 c_2 \vec{e}_2 + b_4 c_3 \vec{e}_3 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{D}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_4] = -\vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The product  $[\vec{b}, \vec{D}]$  generates a contradiction  $\vec{e}_3 = \vec{0}$ . So, subalgebra  $h_5$  has no reductive complement with basis (3).

Let  $\vec{c} = \vec{e}_2 + c_4 \vec{e}_4, \vec{d} = \vec{e}_3 + d_4 \vec{e}_4$  be the basis (4) for a possible reductive complement  $m$ .

Multiply basic vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$  and  $\vec{b} = \vec{e}_3 + b_4 \vec{e}_4$  of  $h_5$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$[\vec{a}, \vec{C}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_2 + c_4 \vec{e}_4] = \vec{e}_2 - \vec{e}_2 = \vec{0} \text{ - it is the identity;}$$

$$[\vec{a}, \vec{D}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_3 + \vec{e}_3 = \vec{0} \text{ - it is the identity;}$$

$$[\vec{b}, \vec{C}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_2 + c_4 \vec{e}_4] = (\vec{e}_1 - \vec{e}_4) - c_4 \vec{e}_3 - b_4 \vec{e}_2 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{D}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = -d_4 \vec{e}_3 + b_4 \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The product  $[\vec{b}, \vec{C}]$  contains vector  $\vec{e}_1$  that can't be generated by vectors  $\vec{c}, \vec{d}$ . This means that subalgebra  $h_5$  has no reductive complement with basis (4).

5. Let  $\vec{c} = \vec{e}_2 + c_3 \vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (5) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1 + \vec{e}_4$  and  $\vec{b} = \vec{e}_3 + b_4 \vec{e}_4$  of  $h_5$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$[\vec{a}, \vec{C}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_2 + c_3 \vec{e}_3] = \vec{e}_2 - c_3 \vec{e}_3 - \vec{e}_2 + c_3 \vec{e}_3 = \vec{0} \text{ - it is the identity;}$$

$$[\vec{a}, \vec{D}] = [\vec{e}_1 + \vec{e}_4, \vec{e}_4] = \vec{0} \text{ - it is the identity;}$$

$$[\vec{b}, \vec{C}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_2 + c_3 \vec{e}_3] = -(\vec{e}_1 - \vec{e}_4) - b_4 \vec{e}_2 + b_4 c_3 \vec{e}_3 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{D}] = [\vec{e}_3 + b_4 \vec{e}_4, \vec{e}_4] = -\vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The product  $[\vec{b}, \vec{C}]$  contains vector  $\vec{e}_1$  that can't be generated by vectors  $\vec{c}, \vec{d}$ . This means that subalgebra  $h_5$  has no reductive complement with basis (5).

6. Let  $\vec{c} = \vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (6) for a possible reductive complement  $m$ . Compare  $m = \text{Span}\{\vec{e}_3, \vec{e}_4\}$  and  $h_5 = \text{Span}\{\vec{e}_1 + \vec{e}_4, \vec{e}_3 + b_4 \vec{e}_4\}$ . It's obvious that  $m \cap h_5 = \{\vec{e}_3 + b_4 \vec{e}_4\}$ . So, this pair is not reductive.

$$\text{Subalgebra } h_6 = \text{Span}\{\vec{e}_1, \vec{e}_4\}$$

Let  $\vec{c} = \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4, \vec{d} = \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{b} = \vec{e}_4$  and  $\vec{b} = \vec{e}_4$  of  $h_6$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$[\vec{a}, \vec{C}] = [\vec{e}_1, \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4] = -c_3 \vec{e}_3 = x_1 \vec{c} + y_1 \vec{d},$$

$$[\vec{a}, \vec{D}] = [\vec{e}_1, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = \vec{e}_2 - d_3 \vec{e}_3 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{C}] = [\vec{e}_4, \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4] = c_3 \vec{e}_3 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{D}] = [\vec{e}_4, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_2 + d_3 \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The system of the vector equations has the solution  $c_3=0, d_3=0, d_4=0$ .

The corresponding subspace  $m = \text{Span}\{\vec{e}_1 + c_4 \vec{e}_4, \vec{e}_2\}$  is not a reductive complement for  $h_6$  because  $h_6 \cap m = \{\vec{e}_1 + c_4 \vec{e}_4\} \neq \vec{0}$ .

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4, \vec{d} = \vec{e}_3 + d_4 \vec{e}_4$  be the basis (2) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1$  and  $\vec{b} = \vec{e}_4$  of  $h_6$  by  $\vec{C}$  and  $\vec{D}$ . We have:

$$[\vec{a}, \vec{C}] = [\vec{e}_1, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = c_2 \vec{e}_2 = x_1 \vec{c} + y_1 \vec{d},$$

$$[\vec{a}, \vec{D}] = [\vec{e}_1, \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_3 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{C}] = [\vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = -c_2 \vec{e}_2 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{D}] = [\vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The system of the vector equalities has the solution  $c_2=0, d_4=0$ . The corresponding subspace  $m = \text{Span}\{\vec{e}_1 + c_4 \vec{e}_4, \vec{e}_3\}$  is not a reductive complement for  $h_6$  because  $h_6 \cap m = \{\vec{e}_1 + c_4 \vec{e}_4\} \neq \vec{0}$ .

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3, \vec{d} = \vec{e}_4$  be the basis (3) for a possible reductive complement  $m$ . It is obvious that  $h_6 \cap m = \{\vec{e}_4\} \neq \vec{0}$ , so subalgebra  $h_6$  has no reductive complement with basis (3).

Let  $\vec{c} = \vec{e}_2 + c_4 \vec{e}_4$ ,  $\vec{d} = \vec{e}_3 + d_4 \vec{e}_4$  be the basis (4) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_1$  and  $\vec{b} = \vec{e}_4$  of  $h_6$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_1, \vec{e}_2 + c_4 \vec{e}_4] = \vec{e}_2 = x_1 \vec{c} + y_1 \vec{d},$$

$$[\vec{a}, \vec{d}] = [\vec{e}_1, \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_3 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{c}] = [\vec{e}_4, \vec{e}_2 + c_4 \vec{e}_4] = \vec{e}_2 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The system of the vector equalities has the solution  $c_4=0, d_4=0$ . We obtain the new reductive pair  $h_6 = Span\{\vec{e}_1, \vec{e}_4\}$ ,  $m_1 = Span\{\vec{e}_2, \vec{e}_3\}$ .

Let  $\vec{c} = \vec{e}_2 + c_3 \vec{e}_3$ ,  $\vec{d} = \vec{e}_4$  be the basis (5) for a possible reductive complement  $m$ . It is obvious that  $h_6 \cap m = \{\vec{e}_4\} \neq \vec{0}$  in this case. So, subalgebra  $h_6$  has no reductive complement with basis (5).

Let  $\vec{c} = \vec{e}_3$ ,  $\vec{d} = \vec{e}_4$  be the basis (6) for a possible reductive complement  $m$ . Consider this pair  $m = Span\{\vec{e}_3, \vec{e}_4\}$ ,  $m = Span\{\vec{e}_3, \vec{e}_4\}$ . It's obvious that  $h_6 \cap m \neq \vec{0}$ . So, subalgebra  $h_6$  has no reductive complement with basis (6).

Subalgebra  $h_7 = Span\{\vec{e}_2, \vec{e}_4\}$

Let  $\vec{c} = \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4$ ,  $\vec{d} = \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_2$  and  $\vec{b} = \vec{e}_4$  of  $h_7$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{d}] = [\vec{e}_2, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = d_3(\vec{e}_1 - \vec{e}_4) + d_4 \vec{e}_2 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{a}, \vec{c}] = [\vec{e}_2, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = d_3(\vec{e}_1 - \vec{e}_4) + d_4 \vec{e}_2 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{c}] = [\vec{e}_4, \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4] = c_3 \vec{e}_3 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_4, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_2 + d_3 \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The solution of the system of vector equations is  $c_3=0, d_3=0, d_4=0$ . This solution produces the following pair  $h_7 = Span\{\vec{e}_2, \vec{e}_4\}$ ,  $m = Span\{\vec{e}_1 + c_4 \vec{e}_4, \vec{e}_2\}$  that is not reductive because  $h_7 \cap m = \{\vec{e}_2\} \neq \vec{0}$ .

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4$ ,  $\vec{d} = \vec{e}_3 + d_4 \vec{e}_4$  be the basis (2) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_2$  and  $\vec{b} = \vec{e}_4$  of  $h_7$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_2, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = -\vec{e}_2 + c_4 \vec{e}_2 = x_1 \vec{c} + y_1 \vec{d},$$

$$[\vec{a}, \vec{d}] = [\vec{e}_2, \vec{e}_3 + d_4 \vec{e}_4] = (\vec{e}_1 - \vec{e}_4) + d_4 \vec{e}_2 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{c}] = [\vec{e}_4, \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4] = -c_2 \vec{e}_2 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4] = \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

This system of vector equations has no solution because of the contradiction  $c_4=1$  from the product  $[\vec{a}, \vec{c}]$  and  $c_4=-1$  (from the product  $[\vec{b}, \vec{c}]$ ). This means that subalgebra  $h_7$  has no reductive complement with basis (2).

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3$ ,  $\vec{d} = \vec{e}_4$  be the basis (3) for a possible reductive complement  $m$ . Subalgebra  $h_7$  has the basis  $\vec{a} = \vec{e}_2$ ,  $\vec{b} = \vec{e}_4$ . It's obvious that  $h_7 \cap m = \{\vec{e}_4\} \neq \vec{0}$ , so subalgebra  $h_7$  has no reductive complement with basis (3).

For the bases (4), (5), and (6) we have the similar cases as for the basis (3). Subalgebra  $h_7$  has no reductive complements at all.

Subalgebra  $h_8 = Span\{\vec{e}_3, \vec{e}_4\}$

Let  $\vec{c} = \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4$ ,  $\vec{d} = \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4$  be the basis (1) for a possible reductive complement  $m$ . Multiply basic vectors  $\vec{a} = \vec{e}_3$  and  $\vec{b} = \vec{e}_4$  of  $h_8$  by  $\vec{c}$  and  $\vec{d}$ . We have:

$$[\vec{a}, \vec{c}] = [\vec{e}_3, \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4] = \vec{e}_3 - c_4 \vec{e}_3 = x_1 \vec{c} + y_1 \vec{d},$$

$$[\vec{a}, \vec{d}] = [\vec{e}_3, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = -(\vec{e}_1 - \vec{e}_4) - d_4 \vec{e}_3 = x_2 \vec{c} + y_2 \vec{d},$$

$$[\vec{b}, \vec{c}] = [\vec{e}_4, \vec{e}_1 + c_3 \vec{e}_3 + c_4 \vec{e}_4] = c_3 \vec{e}_3 = x_3 \vec{c} + y_3 \vec{d},$$

$$[\vec{b}, \vec{d}] = [\vec{e}_4, \vec{e}_2 + d_3 \vec{e}_3 + d_4 \vec{e}_4] = -\vec{e}_2 + d_3 \vec{e}_3 = x_4 \vec{c} + y_4 \vec{d}.$$

The system of vector equations has no solution because the contradiction  $c_4=1, c_4=-1$  appears. This means that subalgebra  $h_8$  has no reductive complement with basis (1).

Let  $\vec{c} = \vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4$ ,  $\vec{d} = \vec{e}_3 + d_4 \vec{e}_4$  be the basis (2) for a possible reductive complement  $m$ . It is obvious that the corresponding pair  $h_8 = Span\{\vec{e}_3, \vec{e}_4\}$ ,  $m = Span\{\vec{e}_1 + c_2 \vec{e}_2 + c_4 \vec{e}_4, \vec{e}_3 + d_4 \vec{e}_4\}$  is not reductive because  $h_8 \cap m = \{\vec{e}_3 + d_4 \vec{e}_4\} \neq \vec{0}$ .

For the bases (3), (4), (5), and (6) we have the similar situations like for the bases (1) and (2). The subalgebra  $h_8$  has no reductive complements with all these bases.

The next theorem describes all different reductive pairs that were found.

**Theorem 2:** Each reductive pair  $\{h, m\}$  with 2-dimensional subalgebra  $h$  and 2-dimensional complement  $m$  of Lie algebra  $g$  is equal to one and only one pair from the next list of them:

- $h_1 = Span\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_4 \vec{e}_4\}$ ,  $m_1 = Span\{\vec{e}_1 + c_4 \vec{e}_4, \vec{e}_2\}$ ,  $b_4 \neq 0, c_4 \neq 1$ ;
- $h_2 = Span\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3 \vec{e}_3 + b_4 \vec{e}_4\}$ ,  $m_2 = Span\{\vec{e}_1 + c_3 \vec{e}_3, \vec{e}_2 + c_4 \vec{e}_4\}$ , where  $c_3 \neq b_4$ , and  $c_3$  is any real solution of the equation  $c_3^2 - b_4 c_3 - b_3 = 0$ ;
- $h_3 = Span\{\vec{e}_1 + \vec{e}_4, \vec{e}_2 + b_3 \vec{e}_3 + b_4 \vec{e}_4\}$ ,  $m_3 = Span\{\vec{e}_1 + c_2 \vec{e}_2, \vec{e}_3 + c_2 \vec{e}_4\}$ , where  $c_2$  is any real solution of the equation  $b_3 c_2^2 - b_4 c_2 - 1 = 0$ ;
- $h_4 = Span\{\vec{e}_1 + a_3 \vec{e}_3 + a_4 \vec{e}_4, \vec{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \vec{e}_3 + \frac{1+a_4}{1-a_4} a_3 \vec{e}_4\}$ ,  $a_4 \neq 1$ ,  
 $m_1 = Span\{\vec{e}_1 + \frac{a_3}{1-a_4} \vec{e}_3, \vec{e}_2 + \frac{a_3}{1-a_4} \vec{e}_4\}$  where  $a_3 \neq 0, a_4 \neq 0$ .
- $h_5 = Span\{\vec{e}_1 + a_3 \vec{e}_3 + a_4 \vec{e}_4, \vec{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \vec{e}_3 + \frac{1+a_4}{1-a_4} a_3 \vec{e}_4\}$ ,  $a_4 \neq 1, a_3 \neq 0, a_4 \neq -1$ ,  
 $m_2 = Span\{\vec{e}_1 + \frac{3-a_4}{2(1-a_4)} a_3 \vec{e}_3 + \frac{a_4-1}{2} \vec{e}_4, \vec{e}_2 + \frac{a_3^2}{(1-a_4)^2} \vec{e}_3 + \frac{1+a_4}{2(1-a_4)} a_3 \vec{e}_4\}$ .
- $h_6 = Span\{\vec{e}_1 + a_3 \vec{e}_3 + a_4 \vec{e}_4, \vec{e}_2 - \frac{a_4 a_3^2}{(1-a_4)^2} \vec{e}_3 + \frac{1+a_4}{1-a_4} a_3 \vec{e}_4\}$ ,  $a_4 \neq 1$ ,



$$m_3 = \text{Span}\{\bar{e}_1 + \frac{a_4 - 1}{a_3}\bar{e}_2, \bar{e}_3 + \frac{a_4 - 1}{a_3}\bar{e}_4\}, a_3 \neq 0.$$

$$7. h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\},$$

$$m_1 = \text{Span}\{\bar{e}_1 + c_3\bar{e}_3 + (1 + a_2c_3)\bar{e}_4, \bar{e}_2 - \frac{a_2c_3 + 1}{a_2}\bar{e}_3 - \frac{a_2c_3 + 2}{a_2}\bar{e}_4\}, a_2 \neq 0.$$

$$8. h_4 = \text{Span}\{\bar{e}_1 + a_2\bar{e}_2, \bar{e}_3 + a_2\bar{e}_4\}, m_2 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3\}, a_2 \neq 0.$$

$$9. h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}, m_1 = \text{Span}\{\bar{e}_1 + c_4\bar{e}_4, \bar{e}_3\}, b_4 \neq 0.$$

$$10. h_5 = \text{Span}\{\bar{e}_1 + \bar{e}_4, \bar{e}_3 + b_4\bar{e}_4\}, m_2 = \text{Span}\{\bar{e}_1 + b_4\bar{e}_2 - \bar{e}_4, \bar{e}_3\}, b_4 \neq 0.$$

$$11. h_6 = \text{Span}\{\bar{e}_1, \bar{e}_4\}, m_1 = \text{Span}\{\bar{e}_2, \bar{e}_3\}.$$

**Remark:** It is unknown yet which reductive pairs from Theorem

2 are equivalent with respect to the inner automorphisms of the given Lie algebra.

#### References

1. Shtukar U (2016) Classification of Canonical Bases for  $(n-2)$ -dimensional Subspaces of  $n$ -dimensional Vector Space. Journal of Generalized Lie Theory and Applications 10(1).
2. Shtukar U (2017) Classification of Maximal Subalgebras and Corresponding Reductive Pairs of Lie Algebra of All  $2 \times 2$  Real Matrices. Journal of Generalized Lie Theory and Applications (Submitted).
3. Nomizu K (1954) Invariant Affine Connections on Homogeneous Spaces. American Journal of Mathematics 76: 33-65.
4. Patera J, Winternitz P (1977) Subalgebras of real three- and four-dimensional Lie algebras. Journal of Mathematical Physics 18: 1449-1455.
5. Sagle A, Winter DJ (1967) On Homogeneous Spaces and Reductive Subalgebras of Simple Lie Algebras. Transactions of the American Mathematics Society 128: 142-147.