

Cognition and Intervention in Traumatic Brain Injury

Trevor Archer*

Department of Psychology, University of Gothenburg, Gothenburg, Sweden

Introduction

The destructive ravages of externally- or internally-induced damage to the brain and CNS trauma, whether this be of mild, moderate or severe extent, will invariably be the consequence of proximal-distal causation, proliferation and severity [1-3], such as cerebral ischemia which afflicts millions of individuals worldwide with eventual survivors suffering from long-term functional and cognitive deficits. Cerebral damage in traumatic brain injury (TBI) may be accompanied by headache, of varying intensities and localizations, dizziness and ill-feeling, loss of consciousness, blurring of vision, hearing impediments, confusion states, loss of memory and cognitive capacity, seizure activity, paralysis and coma as the primary-registered symptoms and expressions, with the expectation of accompanying damage to blood-brain-barrier integrity, accelerated apoptosis and excitotoxicity [4-6]. Further, there is a profusion of balance and attentive disorders in TBI independent of several other injury parameters. Neurobehavioral problems, mood and cognition disorder, especially with regard to memory performance, attention, planning and executive functioning are invariably impaired in individuals afflicted by TBI, a major and leading cause of chronic disability over the globe. All of these signs and symptoms in various wound-combinations, intensities and severities, locations and stages will offer the determination of the dimensions of short-term and long-term disruption of physical, sensory-motor, cognitive-emotional, behavioral and emotional domains [7-9]. Efforts to establish diagnostic and symptomatic instruments for establishing deficits in cognitive domains of TBI, not least regarding self-awareness [10,11] and mindfulness [12-14], have produced several essential methodological ingredients.

Rehabilitative progress has been obtained despite much destruction of the higher-order cognitive systems [15]. One obstacle barring interventional progress is related to insufficiency of development and the norm-achieving basis of instruments/methods applicable for detection of impairments to social cognition [16]. Examinations of the sub-acute effects of TBI upon gray matter volumetric analyses in children presenting severe TBI have shown the existence of marked reductions in the Cerebro-Cerebellar Mentalizing Network, the Salience Network, the Mirror Neuron/Empathy Network, the Central Executive Network and the Default Mode Network, as well as reduced gray-matter volumes of several central-hub regions of these essential neural networks [17]. Similar volumetric deterioration in the Cerebro-Cerebellar Mentalizing Network and several of its hub regions, including the cerebellum, predicted poorer cognitive ToM. Contrastingly, the disrupted cognitive-affective and conative aspects of Theory of Mind were predicted by volumetric reductions in the Salience Network and Mirror Neuron/Empathy Network, respectively. Overall, these findings imply that cognitive, affective and conative ToM may be predicted imminently through the individual differences in structures of different neural systems-the Cerebro-Cerebellar Mentalizing Network, Salience Network, and Mirror Neuron/Empathy Network, in each turn. TBI patients were observed to perform markedly worse than the healthy-volunteer control group with regard to all of the tests of emotion recognition, intention-comprehension, and on one particular task of response selection that together describe social learning [18].

The assessed ratings of current behavioral activity implied important alterations among the TBI patients' behavioral expressions relative to the situation prior to injury through a notable deterioration of community integration and interpersonal behavior than the healthy control group. Of the above three functions that were assessed, emotion recognition was associated with both deleterious post-injury behavior and community integration with the evidence that the association was not to be completely explained by injury severity, time since injury, or education.

Nevertheless, several useful interventions have been found to be effective [19]. For example, in an experimental laboratory setting it was observed that 6-hours of environmental enrichment provision on a daily basis was effective in conferring neurobehavioral and histological amelioration following TBI with extended improvements through the realization that the acquired benefits were comparable despite the manner in which the 6-hours of enriched environment were accrued [20]. The development of coping strategies for thriving and disorder adaptation, as well as the psychological adaptation to stressors and serious life events, have been discovered to exert a marked influence on the progressive improvement and against the persistence of posttraumatic injury complaints. Ongoing and prospective interventional therapies are aimed at the restoration brain plasticity in that they are designed to induce the recovery of cognitive functioning, the achievement which may implement the alleviation of other more-or-less related disorders which commonly are defined by the disruptions of diverse cognitive processing [21]. Wong et al. [22] have argued that psychosocial and physical environments appear to mediate the influence of systems, services, and policies on participation after acquired neurological disorders [22]. Furthermore, randomized clinical trials upon pediatric patients presenting moderate-to-severe TBI and involving more than 600 participants from eight clinical centers have proven relatively efficacious [23]. The active, as opposed to passive, style of coping strategy has been found to be significant for patient well-being and self-esteem progressive since the former are much more effective [24].

In afflicted individuals, the loss of structural connectivity following mild-to-moderate TBI may be applied as a biomarker estimation both of damage extent and of any putative improvements following intervention. In adolescents presenting TBI, global efficiency level, mean local efficiency scores, modularity, normalized clustering coefficient expression, normalized characteristic path length, a measure of structural connectivity abnormality, and small-worldness assessments, as well as Post-Concussion Symptom Inventory scoring,

*Corresponding author: Archer T, Department of Psychology, University of Gothenburg, 40530 Gothenburg, Sweden, Tel: +46 31 7864694; E-mail: trevor.archer@psy.gu.se

Received: May 29, 2017; Accepted: May 30, 2017; Published: June 07, 2017

Citation: Archer T (2017) Cognition and Intervention in Traumatic Brain Injury. Clin Exp Psychol 3: e108. doi: [10.4172/2471-2701.1000e108](https://doi.org/10.4172/2471-2701.1000e108)

Copyright: © 2017 Archer T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

were all inferior to healthy control adolescents, but following aerobic training, global efficiency level was increased and normalized characteristic path length was decreased [25]. Also, the enhanced Post-Concussion Symptom Inventory scores were correlated significantly with the global efficiency level improvements and normalized characteristic path length decrease. A novel interventional approach designed to improve both mobility and attention deficits through application a reward-learning procedure is offered through video-gaming therapy [26,27]. In a study of TBI patients, it was observed that the patients receiving video-gaming therapy performed better than those receiving 'balance-platform' therapy on the Community Balance & Mobility Scale, the Unified Balance Scale, the Timed Up and Go test (TUG), the static balance test and the selective visual attention evaluation test thereby implying positive assets with the technique [28]. In a recent study performed on army (US) veterans suffering from mild TBI, it was observed that Compensatory Cognitive Training presented higher levels of improvement in cognitive difficulties and application of cognitive strategies as well as on neuropsychological measures of attention, learning, and executive functioning than the comparison group that received only 'usual-care' treatment [29]. Interestingly, it has been reported also that pharmaco-therapeutic interventions using a compound co-administration regimen of the antioxidants, 2,4-disulfonyl α -phenyl tertiary butyl nitron (HPN-07), a spin-trapagent, and N-acetylcysteine (NAC) decreased markedly both the pathologic Tau accumulation and signs of ongoing neurodegeneration in the cochlea and the auditory cortex of laboratory rats [30]. Physical, psychological and mental fatigue distress 70-90% of TBI with serious aggravation of other symptoms and biomarkers. In a study of adults with a history of TBI that were presented cognitive behavior therapy or 'treatment-as-usual', the former recipients reported better sleep quality together a significant decrease in daily fatigue levels compared with the latter [31].

Taken together, the evidence implies that the cognitive-emotion condition of TBI patient is hazardous from the point of view of health and well-being, particularly with the mild, moderate and severe levels of infirmity [32]. Despite the cognitive impairments arising from limited-to-substantial connectivity with white and gray matter that result in widespread disharmony among essential network systems, the potential for promising interventional therapies seems both efficacious and technologically innovative, such as at synaptic sites of neural plasticity [33]. It is becoming increasingly apparent that the social determinants of TBI, e.g. physical environment, gender, interpersonal violence-propensity, personal health practices and coping skills and availability of rehabilitation services, require greater attention, particularly among minority ethnic groups [34].

References

- Hinson HE, Schreiber MA, Laurie AL, Baguley IJ, Bourdette D, et al. (2017) Early fever as a predictor of paroxysmal sympathetic hyperactivity in traumatic brain injury. *J Head Trauma Rehabil*.
- Omar M, Moore L, Lauzier F, Tardif PA, Dufresne P, et al. (2017) Complications following hospital admission for traumatic brain injury: A multicenter cohort study. *J Crit Care* 41: 1-8.
- Spurlock MS, Ahmed AI, Rivera KN, Yokobori S, Lee SW, et al. (2017) Amelioration of penetrating ballistic-like brain injury induced cognitive deficits after neuronal differentiation of transplanted human neural stem cells. *J Neurotrauma* 34: 1981-1995.
- Braun M, Vaibhav K, Saad NM, Fatima S, Vender JR, et al. (2017) White matter damage after traumatic brain injury: A role for damage associated molecular patterns. *Biochim Biophys Acta*.
- Paci M, Infante-Rivard C, Marcoux J (2017) Traumatic brain injury in the workplace. *Can J Neurol Sci*, pp: 1-7.
- Sutton JA, Clauss RP (2017) A review of the evidence of zolpidem efficacy in neurological disability after brain damage due to stroke, trauma and hypoxia: A justification of further clinical trials. *Brain Inj*, pp: 1-9.
- Belchev Z, Levy N, Berman I, Levinzon H, Hoofien D, et al. (2017) Psychological traits predict impaired awareness of deficits independently of neuropsychological factors in chronic traumatic brain injury. *Br J Clin Psychol*.
- Maneewong J, Maneeton B, Maneeton N, Vaniyapong T, Traisathit P, et al. (2017) Delirium after a traumatic brain injury: Predictors and symptom patterns. *Neuropsychiatr Dis Treat* 13: 459-465.
- Winter L, Moriarty HJ, Short TH (2017) Self-reported driving difficulty in veterans with traumatic brain injury: Its central role in psychological well-being.
- Lloyd O, Ownsworth T, Fleming J, Zimmer-Gembeck MJ (2017) Development and preliminary validation of the Paediatric Awareness Questionnaire for children and adolescents with traumatic brain injury. *Child Neuropsychol* 23: 1-21.
- Sherer M, Poritz JMP, Tulsy D, Kisala P, Leon-Novelo L, et al. (2017) Conceptual structure of health related quality of life for persons with traumatic brain injury: Confirmatory factor analysis of the TBI-QOL. *Arch Phys Med Rehabil*.
- Haller CS, Bosma CM, Kapur K, Zafonte R, Langer EJ (2017) Mindful creativity matters: trajectories of reported functioning after severe traumatic brain injury as a function of mindful creativity in patients' relatives: A multilevel analysis. *Qual Life Res* 26: 893-902.
- Kristofersson GK, Beckers T, Krueger R (2017) Perceptions of an adapted mindfulness program for persons experiencing substance use disorders and traumatic brain injury. *J Addict Nurs* 27: 247-253.
- Xu GZ, Li YF, Wang MD, Cao DY (2017) Complementary and alternative interventions for fatigue management after traumatic brain injury: A systematic review. *Ther Adv Neurol Disord* 10: 229-239.
- Dang B, Chen W, He W, Chen G (2017) Rehabilitation treatment and progress of traumatic brain injury dysfunction. *Neural Plast*.
- Kelly M, McDonald S, Frith MHJ (2016) A survey of clinicians working in brain injury rehabilitation: Are social cognition impairments on the radar? *J Head Trauma Rehabil*.
- Ryan NP, Catroppa C, Beare R, Silk TJ, Hearps SJ, et al. (2017) Uncovering the neuroanatomical correlates of cognitive, affective, and conative theory of mind in pediatric traumatic brain injury: A neural systems perspective. *Soc Cogn Affect Neurosci*.
- May M, Milders M, Downey B, Whyte M, Higgins V, et al. (2017) Social behavior and impairments in social cognition following traumatic brain injury. *J Int Neuropsychol Soc* 23: 400-411.
- Archer T (2012) Influence of physical exercise on traumatic brain injury deficits: Scaffolding effect. *Neurotox Res* 21: 418-34.
- Radabaugh HL, LaPorte MJ, Greene AM, Bondi CO, Lajud N, et al. (2017) Refining environmental enrichment to advance rehabilitation based research after experimental traumatic brain injury. *Exp Neurol* 294: 12-18.
- Stradecki-Cohan HM, Cohan CH, Raval AP, Dave KR, Reginensi D, et al. (2017) Cognitive deficits after cerebral ischemia and underlying dysfunctional plasticity: potential targets for recovery of cognition. *J Alzheimers Dis*.
- Wong AWK, Ng S, Dashner J, Baum MC, Hammel J, et al. (2017) Relationships between environmental factors and participation in adults with traumatic brain injury, stroke, and spinal cord injury: a cross-sectional multi-center study. *Qual Life Res*.
- Wade SL, Kurowski BG (2017) Behavioral clinical trials in moderate to severe pediatric traumatic brain injury: Challenges, potential solutions, and lessons learned. *J Head Trauma Rehabil*.
- Scheenen ME, van der Horn HJ, de Koning ME, van der Naalt J, Spikman JM (2017) Stability of coping and the role of self-efficacy in the first year following mild traumatic brain injury. *Soc Sci Med*. 181:184-190.
- Yuan W, Wade SL, Quatman-Yates C, Hugentobler JA, Gubanich PJ, et al. (2017) Structural Connectivity Related to Persistent Symptoms After Mild TBI in Adolescents and Response to Aerobic Training: Preliminary Investigation. *J Head Trauma Rehabil*.
- Archer T (2017) Educational-exercise stimulation by video-gaming. *BAOJ Psychology* 2: 020.

27. Archer T, Garcia D, Moradi S (2016) The enigmatic influence of video-internet gaming: liabilities and assets over the lifespan. *Sports Medicine and Rehabilitation Journal* 1: 1008-1017.
28. Straudi S, Severini G, Sabbagh CA, Pavarelli C, Gamberini G, et al. (2017) The effects of video game therapy on balance and attention in chronic ambulatory traumatic brain injury: An exploratory study. *BMC Neurol* 17: 86.
29. Pagulayan KF, O'Neil M, Williams RM, Turner AP, Golshan S, et al. (2017) Mental health does not moderate compensatory cognitive training efficacy for veterans with a history of mild traumatic brain injury. *Arch Phys Med Rehabil*.
30. Du X, West MB, Cai Q, Cheng W, Ewert DL, et al. (2017) Antioxidants reduce neurodegeneration and accumulation of pathologic Tau proteins in the auditory system after blast exposure. *Free Radic Biol Med* 108: 627-643.
31. Nguyen S, McKay A, Wong D, Rajaratnam SM, Spitz G, et al. (2017) Cognitive behavior therapy to treat sleep disturbance and fatigue following traumatic brain injury: A pilot randomized controlled trial. *Arch Phys Med Rehabil*.
32. von Steinbüchel N, Real RGL, Sasse N, Wilson L, Otto C, et al. (2017) German validation of Quality of Life after Brain Injury (QOLIBRI) assessment and associated factors. *PLoS One* 12: e0176668.
33. Wen Z, Li D, Shen M, Chen G (2017) Therapeutic potentials of synapses after traumatic brain injury: A comprehensive review. *Neural Plast* 2017: 4296075.
34. Zeiler KJ, Zeiler FA (2017) Social Determinants of traumatic brain injury in the north american indigenous population: A review. *Can J Neurol Sci*, pp: 1-7.

Citation: Archer T (2017) Cognition and Intervention in Traumatic Brain Injury. Clin Exp Psychol 3: e108. doi: [10.4172/2471-2701.1000e108](https://doi.org/10.4172/2471-2701.1000e108)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>