

Commentary on *Psilocybe cyanescens*

Alexander Giessler*

University of Göttingen, Germany

Psilocybe cyanescens Wakef. is a potently psychoactive synanthrope increasingly found in parks and flowerbeds in North American and European cities [1-3]. It is spreading by mulch usage, garden waste and outdoor cultivations of private consumers. Dennis and Wakefield [4] described the holotype from a botanical garden in London. After decades of globalization, the question is standing to reason, where this saprobic fungal alien initially came from.

Although a species delimitation to the European *P. serbica* [5] meanwhile is accepted among mycologists, *P. cyanescens*' relation to the North American *P. azurescens* [6] and *P. allenii* [7], as well as the Australian *P. subaeruginosa* [8], is still unclear. They more or less distinguish in morphology and distribution, but show a very close molecular phylogeny. Furthermore, *P. cyanescens*, *P. azurescens* and *P. allenii* are known to mate by plasmogamy. The taxonomical discussion requires more international studies with comprehensive strain collections.

P. cyanescens from England, Germany and the Pacific Northwest of the United States are characterized by the same, merely four mating types in a tetrapolar incompatibility system (Giessler, unpublished). Basidiomycetes usually have more mating types, to promote outbreeding, especially between farer related populations. The lack of mating types may be used as an argument for a recent, anthropogenously promoted spreading, because it indicates a small gene pool due to a missing allopatric differentiation. *P. cyanescens* obviously represents a genetic bottleneck, but where is the bottle's belly?

Maybe *P. subaeruginosa* colonies were the origin of *P. cyanescens*, *P. azurescens* and *P. allenii*. This species varies between the characteristics of the other three. It often resembles

P. cyanescens, but has a higher number of mating types [9,10]. Simple mating experiments of *P. cyanescens*, *P. azurescens* and *P. allenii* with the Australian relative, accompanied by fructification experiments considering the ability to produce fertile offspring, would answer a lot of questions. An export of *P. subaeruginosa* to America and Europe, for example, by the shipping of wood or exotic plants, in the early 20th century is well imaginable.

References

1. Gartz J (1996) Observations on the *Psilocybe cyanescens* complex of Europe and North America. *Annali dei Musei Civici di Rovereto* 12: 209-218.
2. Stamets P (2000) Growing gourmet and medicinal mushrooms. Ten Speed Press, Berkeley.
3. Shaw PJA, Kibby G (2001) Aliens in the flowerbeds. The fungal biodiversity of ornamental woodchips. *Field Mycology* 2: 6-11.
4. Dennis RWG, Wakefield EM (1946) New or interesting British fungi. *Transactions of the British Mycological Society* 24: 141-166.
5. Borovička J, Noordeloos ME, Gryndler M, Obornik M (2011) Molecular phylogeny of *Psilocybe cyanescens* complex in Europe, with reference to the position of the secotioid *Weraroa novae-zelandiae*. *Mycol Prog* 10: 149-155.
6. Stamets P, Gartz J (1995) A new caerulescent *Psilocybe* from the pacific coast of northwestern America. *Integration* 6: 21-27.
7. Borovička J, Rockefeller A, Werner PG (2012) *Psilocybe allenii*-A new bluing species from the Pacific Coast, USA. *Czech Mycology* 64: 181-195.
8. Cleland J, Burton J (1927) Australian fungi: notes and descriptions-No. 6. *Transactions and Proceedings of the Royal Society of South Australia* 51: 298-306.
9. Chang YS, Mills AK (1992) Re-examination of *Psilocybe subaeruginosa* and related species with comparative morphology, isozymes and mating compatibility studies. *Mycol Res* 96: 429-441.
10. Johnston P, Buchanan PK (1995) The genus *Psilocybe* (Agaricales) in New Zealand. *N Z J Botany* 33: 379-388.

*Corresponding author: Alexander Giessler, University of Göttingen, Germany, E-mail: oeke.junior@gmx.de

Received February 13, 2017; Accepted February 16, 2017; Published February 23, 2017

Citation: Giessler A (2017) Commentary on *Psilocybe cyanescens*. *Adv Tech Biol Med* 5: 205. doi: [10.4172/2379-1764.1000205](https://doi.org/10.4172/2379-1764.1000205)

Copyright: © 2017 Giessler A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.