

Former Male Elite Athletes Consume More Alcohol than Controls, but There is No Difference in Alcohol-Related Morbidity or Mortality

Kontro TK*

Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

Introduction

There is a widespread scientific consensus that heavy use of alcohol, smoking, and lack of physical activity contribute to morbidity and mortality [1-3]. The increased mortality related to heavy alcohol consumption is associated with multiple somatic diseases (such as cirrhosis, pancreatitis, certain cancers, stroke, accidents and other external causes of deaths) [4,5], but also with several mental health outcomes (such as depression and suicide). Studies have found that people who drink are much more likely to smoke [6,7], and dependence on alcohol and tobacco is also correlated [8,9]. Many studies have indicated that athletic participation is associated with more excessive use of alcohol, but most studies are focused on drinking habits among college athletes [10,11].

According to previous studies former athletes have lower morbidity and mortality than general population [12-15], but there is lack of evidence on how continuation of physical activity, alcohol consumption and smoking habits contributes to former athletes' health in later years. So, a recent study includes novel data and findings analysed previously neither from the Finnish former athlete cohort nor from any other comparable cohort worldwide [16].

A recent study [16] had three aims, the first of which was to investigate how former competitive sports career was associated with alcohol-related morbidity and mortality and with use of alcohol among former athletes. Secondly, the risk of alcohol-related diseases or deaths and alcohol consumption were compared between different sports groups. Thirdly, it was examined how different factors, such as smoking status and physical activity, were associated with use of alcohol among former athletes after their active athletic career.

Methods

In a recent study [16] occurrence of alcohol-related diseases and deaths were followed using national hospital discharge register from 1970 to 2008 among Finnish male former elite athletes (n=2202) and matched controls (n=1403) alive in 1970. Age-adjusted hazard ratios (HRs) for alcohol-related diseases and deaths were calculated by Cox proportional hazards model. In 1985 cohort members answered a questionnaire including items on use of alcohol, smoking habits and physical activity.

Principal Findings and Conclusions

It was found in a recent study [16] that there was no significant difference in alcohol-related morbidity or mortality between former athletes and controls. However, morbidity and mortality was higher among both weightlifters and combat sports athletes compared to endurance sports athletes, jumpers & hurdlers or shooters. Former athletes reported higher alcohol consumption than controls, especially team sports athletes used more alcohol and had heavier drinking occasions than other sports athletes. Use of alcohol after active sports career was greater if participation in physical activities or leisure-time sports was discontinued. Correspondingly, current and ex-smokers used more alcohol than non-smokers [16].

In conclusion, it seems that engagement to competitive sports does not become replaced with alcohol dependence among former athletes after elite level sports career termination. Former athletes' overall better health habits (such as less smoking), and the directly or indirectly subsequent biological factors are likely to explain observation that there was no difference in the risk of alcohol-related morbidity or mortality among former elite athletes compared to controls. This was found despite that former elite athletes consumed more alcohol than controls [16].

The novel findings from a recent long-term follow-up study [16] give new evidence for the benefits of continuing physical activity at an older age. It is important to notice that the risk of excessive alcohol consumption may increase among individuals, who discontinue leisure-time sports after their active sports career. The results may motivate coaches, sports clubs and public health officials to develop models to prevent such as excessive alcohol consumption among risk sports during career termination. Mostly due to the apparent association with a healthier lifestyle, especially physical activity engagement, and consequently, a lower prevalence of risk factors associated with major chronic diseases. Recent study findings will be considered as a stimulus for engaging in sports at an older age among athletes and non-athletes [16].

Future Directions

It is not known exactly how well the results can be generalized to today's athletes, non-athletes or women. So, it is important to investigate alcohol consumption, alcohol-related morbidity and mortality among those subgroups in the future. Furthermore, a deeper understanding of the relationships between use of alcohol and the risk of alcohol-related diseases and sports is necessary to determine the true consequences of alcohol on health and well-being in athletes. Models should be developed how to avoid an excessive use of alcohol among risk sports after an active athletic career.

References

1. Batista CHX, Soares JM (2013) Are former elite athletes more protected against metabolic syndrome? *J Cardiol* 61: 440-445.
2. Batista CHX, Soares JM (2014) Is athletic background associated with a future lower prevalence of risk factors for chronic disease? *J Exercise Sci Fitness* 12: 47-54.
3. Bobo JK, Husten C (2000) Sociocultural influences on smoking and drinking. *Alcohol Res Health* 24: 225-232.

*Corresponding author: Kontro TK, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland, E-mail: titta.k.kontro@student.jyu.fi

Received May 17, 2017; Accepted June 05, 2017; Published June 12, 2017

Citation: Kontro TK (2017) Former Male Elite Athletes Consume More Alcohol than Controls, but There is No Difference in Alcohol-Related Morbidity or Mortality. *J Yoga Phys Ther* 7: 265. doi: [10.4172/2157-7595.1000265](https://doi.org/10.4172/2157-7595.1000265)

Copyright: © 2017 Kontro TK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

4. Carmody TP, Brischetto CS, Matarazzo JD (1985) Co-occurrent use of cigarettes, alcohol, and coffee in healthy, community-living men and women. *Health Psychol* 4: 323-335.
5. Di Castelnuovo A, Costanzo S, Bagnardi V (2006) Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Arch Intern Med* 166: 2437-2445.
6. Fine LJ, Philogene GS, Gramling R (2004) Prevalence of multiple chronic disease risk factors. 2001 National Health Interview Survey. *Am J Prev Med* 27: 18-24.
7. Gajewski AK, Poznanska A (2008) Mortality of top athletes, actors and clergy in Poland: 1924-2000 follow-up study of the long term effect of physical activity. *Eur J Epidemiol* 23: 335-340.
8. Grant BF, Hasin DS, Chou SP (2004) Nicotine dependence and psychiatric disorders in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry* 61: 1107-1115.
9. Kontro TK, Sarna S, Kaprio J, Kujala UM (2017) Use of alcohol and alcohol-related morbidity in Finnish former elite athletes. *Med Sci Sports Exerc* 49: 492-499.
10. Kujala UM, Marti P, Kaprio J (2003) Occurrence of chronic disease in former top-level athletes. Predominance of benefits, risks or selection effects? *Sports Med* 33: 553-561.
11. Li TK, Volkow ND, Baler RD (2007) The biological bases of nicotine and alcohol co-addiction. *Biol Psychiatry* 61: 1-3.
12. Lim SS, Vos T, Flaxman AD (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2224-2260.
13. Lisha NE, Sussman S (2010) Relationship of high school and college sports participation with alcohol, tobacco, and illicit drug use: a review. *Addict Behav* 35: 399-407.
14. Martens MP, Dams-O'Connor K, Beck NC (2006) A systematic review of college student-athlete drinking: prevalence rates, sport-related factors, and interventions. *J Subst Abuse Treat* 31: 305-316.
15. Nelson TF, Wechsler H (2001) Alcohol and college athletes. *Med Sci Sports Exerc* 33: 43-47.
16. World Health Organization (WHO) (2014) Global Status Report on Alcohol and Health. Geneva.

Citation: Kontro TK (2017) Former Male Elite Athletes Consume More Alcohol than Controls, but There is No Difference in Alcohol-Related Morbidity or Mortality. *J Yoga Phys Ther* 7: 265. doi: [10.4172/2157-7595.1000265](https://doi.org/10.4172/2157-7595.1000265)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>