

Common Fixed Point Theorem in T_0 Quasi Metric Space

Balaji R Wadkar¹, Ramakant Bhardwaj², Lakshmi Narayan Mishra^{3*} and Basant Singh¹

¹Department of Mathematics, AISECT University, Bhopal-Chiklod Road, Bhopal, Madhya Pradesh, India

²Department of Mathematics, TIT Group of Institutes, Anand Nagar, Bhopal, Madhya Pradesh, India

³Department of Mathematics, Mody University of Science and Technology, Lakshmangarh, Sikar Road, Sikar, Rajasthan, India

Abstract

In this paper, we prove fixed point theorems for generalized C-contractive and generalized S-contractive mappings in a bi-complete di-metric space. The relationship between q- spherically complete T_0 Ultra-quasi-metric space and bi-complete diametric space is pointed out in proposition 3.2. This work is motivated by Petals and Fvidalis in a T_0 -ultra-quasi-metric space

2000 AMS Subject Classification: 47H17, 74H05, 47H09.

Keywords: Fixed Point; Generalized C-Contraction; Generalized S-Contraction; Spherically Complete; Bi-Complete Di-metric

Introduction

In Agyingi [1] proved that every generalized contractive mapping defined in a q- spherically complete T_0 -ultra-quasi metric space has a unique fixed point. In Petals and Fvidalis [2] proved that every contractive mapping on a spherically complete non Archimedean normed space has a unique fixed point. Agyingi and Gega proved fixed point theorems in a T_0 -ultra-quasi-metric space [3-5]. Later many authors published number of papers in this space [6-10].

In this paper we shall prove a fixed point theorem for generalized c- contractive and generalized s-contractive mappings in a bi-complete di-metric space.

If we delete, in the used definition of the pseudo metric d on the set X. the symmetry condition, $d(x, y)=d(y, x)$, whenever $x, y \in X$ we are led to the concept of quasi-pseudo metric.

Definition 1.1: Let (X, m) be a metric space. Let $T: X \rightarrow X$ map is called a C-contraction if there exist, $0 \leq k < \frac{1}{2}$ such that for all $x, y \in X$ the following inequality holds [10],

$$m(Tx, Ty) \leq k [m(x, Tx) + m(y, Ty)]$$

Definition 1.2: Let (X, m) be a metric space. A map $T: X \rightarrow X$ is called a S-contraction if there exist $0 \leq k < \frac{1}{3}$ such that for all $x, y \in X$ the following inequality holds [10]

$$m(Tx, Ty) \leq k [m(x, Tx) + m(y, Ty) + m(x, y)]$$

Preliminaries

Now we recall some elementary definitions and terminology from the asymmetric topology which are necessary for a good understanding of the work below.

Definition 2.1: Let X be a non empty set. A function $d: X \times X \rightarrow [0, \infty)$ is called quasi pseudo metric on X if

$$d(x, x) = 0, \forall x \in X$$

$$d(x, z) \leq d(x, y) + d(y, z), \forall x, y, z \in X$$

Moreover if $d(x, y) = 0 = d(y, x) \Rightarrow x = y$ then d is said to be a T_0 quasi metric or di-metric. The latter condition is referred as the T_0 condition.

Example 2.1: On $R \times R$, we define the real valued map d given by $d(x, y) = |x - y| = \max\{|x - y|, 0\}$ then (R, d) is a di metric space.

Remark 2.1

Let d be quasi-pseudo metric on X, then the map d^{-1} defined by $d^{-1}(x, y) = d(y, x)$ whenever $x, y \in X$ is also a quasi-pseudo metric on X, called the conjugate of d.

It is also denoted by d^t or d^c . It is easy to verify that the function d^s defined by $d^s = d \vee d^{-1}$

$$\text{i.e. } d^s(x, y) = \max\{d(x, y), d(y, x)\}$$

defines a metric on X whenever d is a T_0 quasi pseudo metric.

In some cases, we need to replace $[0, \infty)$ by $[0, \infty]$ (where for a d attaining the value ∞ , the triangle inequality is interpreted in the obvious way). In such case we speak of extended quasi- pseudo metric.

Definition 2.2: The di metric space (X, d) is said to be bi complete if the metric space (R, d^s) is complete.

Example 2.2: Let $X = [0, \infty)$ define for each $x, y \in X$, $n(x, y) = x$ if $x > y$ and $n(x, y) = 0$ if $x < y$. It is not difficult to check that (X, n) is a T_0 quasi pseudo metric space. Notice that, for $x, y \in [0, \infty)$, we have $n^s(x, y) = \max\{x, y\}$ if $x \neq y$ and $n^s(x, y) = 0$ if $x = y$, the matrix n^s is complete on (X, d) .

Definition 2.3: Let (X, d) be quasi pseudo metric space, for $x, \epsilon \in X$ & $\epsilon > 0$

$$B_d(x, \epsilon) = \{y \in X : d(x, y) < \epsilon\}$$

denotes the open $\epsilon -$ ball at x. The collection of such balls is a base for a topology $\tau(d)$ induced by d on X. Similarly for $x, \epsilon \in X$ & $\epsilon \geq 0$

***Corresponding author:** Mishra LN, Department of Mathematics, Mody University of Science and Technology, Lakshmangarh, Sikar Road, Sikar, Rajasthan 332 311, India. Tel: +919838375431; E-mail: lakshminarayanmishra04@gmail.com

Received January 02, 2017; **Accepted** January 30, 2017; **Published** February 08, 2017

Citation: Wadkar BR, Bhardwaj R, Mishra LN, Singh B (2017) Common Fixed Point Theorem in T_0 Quasi Metric Space. Fluid Mech Open Acc 4: 143. doi: 10.4172/2476-2296.1000143

Copyright: © 2017 Wadkar BR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$$C_d(x, \epsilon) = \{y \in X : d(x, y) \leq \epsilon\}$$

denotes the closed ϵ -ball at x .

Definition 2.4: Let (X, d) be quasi pseudo metric space Let $(x_i), i \in I$ be a family of points in X and let $(r_i), i \in I$ & $(s_i), i \in I$ be a family of non negative real numbers.

We say that $(C_d(x_i, r_i), C_{d^{-1}}(x_i, s_i)), i \in I$ has the mixed binary intersection property provided that

$$(C_d(x_i, r_i), C_{d^{-1}}(x_i, s_i)) \neq \emptyset \text{ for all } i, j \in I$$

Definition 2.5: Let (X, d) be quasi pseudo metric space we say that (X, d) is Isbell complete provided that each family $(C_d(x_i, r_i), C_{d^{-1}}(x_i, s_i)), i \in I$ that has the mixed binary intersection property is such that

$$\bigcap_{i \in I} (C_d(x_i, r_i)) \cap (C_{d^{-1}}(x_i, s_i)) \neq \emptyset$$

Proposition 2.1: If (X, d) is an extended Isbell-complete quasi-pseudo metric space then (X, d^s) is hyper complete. An interesting class of quasi pseudo metric space, for which, we investing a type of completeness are the ultra quasi pseudo metric.

Definition 2.6: Let X be a set & $d : X \times X \rightarrow [0, \infty)$ be function a function mapping into the set $[0, \infty)$ of non negative real's then d is ultra quasi pseudo metric on X if

$$d(x, x) = 0 \text{ for all } x \text{ in } X \text{ \& } d(x, y) = d(y, x)$$

$$d(x, z) \leq \max\{d(x, y), d(y, z)\} \text{ whenever } x, y, z \in X$$

The conjugate d^{-1} of d where $d^{-1}(x, y) = d(y, x)$ whenever $x, y \in X$ is also an ultra quasi pseudo metric on X .

If d also satisfies the T_0 -condition, then d is called a T_0 -ultra quasi metric on X . Notice that $d^s = \sup\{d, \bar{d}\} = d \vee d^{-1}$ is an ultra-metric on X whenever d is a T_0 -ultra quasi metric.

In a literature, T_0 -ultra quasi metric spaces are also known as non Archimedean T_0 -quasi metric.

q-spherically Completeness

In this section we shall recall some results about q-spherical completeness belonging mainly to [8].

Definition 3.1: Let (X, d) be an ultra -quasi pseudo metric space Let $(x_i), i \in I$ be a family of points in X and let $(r_i), i \in I$ & $(s_i), i \in I$ be a family of non negative real numbers we say that (X, d) is q-spherical complete provided that each family [2]

$$(C_d(x_i, r_i), C_{d^{-1}}(x_i, s_i)), i \in I$$

Satisfying $d(x_i, s_i) \leq \max\{r_i, s_j\}$, whenever $i, j \in I$ is such that

$$\bigcap_{i \in I} (C_d(x_i, r_i)) \cap (C_{d^{-1}}(x_i, s_i)) \neq \emptyset$$

Proposition 3.2: Each q-spherically complete T_0 ultra quasi metric space (X, d) is bi-complete[8].

Main Results

We recall the following interesting results respectively due to Chatterji [10] and to Shukla [11]

Theorem 4.1a A C-contraction on a complete metric space has a unique fixed point.

Theorem 4.1b A S-contraction on a complete metric space has a unique fixed point.

Following results generalizes the above theorem to setting of a bi-complete di-metric space.

Definition 4.1: Let (X, d) be a quasi pseudo metric space. A map $T : X \rightarrow X$ is called a c-pseudo contraction if there exist $k, 0 \leq k < \frac{1}{2}$ such that for all $x, y \in X$ the following inequality holds.

$$d(Tx, Ty) \leq k[d(Tx, x) + d(y, Ty)]$$

Definition 4.2: Let (X, d) be a quasi-pseudo metric space. A map $T : X \rightarrow X$ is called a S-pseudo contraction if there exist $k, 0 \leq k < \frac{1}{3}$ such that for all $x, y \in X$ the following inequality holds.

$$d(Tx, Ty) \leq k[d(Tx, x) + d(y, Ty) + d(x, y)]$$

Now we define following definitions

Definition 4.3: Let (X, d) be a quasi-pseudo metric space. A map $T : X \rightarrow X$ is called a generalized c-pseudo contraction if there exist $k, 0 \leq k < \frac{1}{4}$ such that for all $x, y \in X$ the following inequality holds.

$$d(Tx, Ty) \leq k\{d(Tx, x) + d(y, Ty) + d(Tx, y) + d(x, Ty)\}$$

Definition 4.4: Let (X, d) be a quasi-pseudo metric space. A map $T : X \rightarrow X$ is called a generalized S-pseudo contraction if there exist $k, 0 \leq k < \frac{1}{8}$ such that for all $x, y \in X$ the following inequality holds.

$$d(Tx, Ty) \leq k\{d(x, Tx) + d(x, Ty) + d(y, Tx) + d(y, Ty) + d(Tx, x) + d(Ty, x) + d(Tx, y) + d(Ty, y)\}$$

Theorem 4.1:

Let (X, d) be a bi complete di metric space and let $T : X \rightarrow X$ be a generalized c-pseudo contraction then T has a unique fixed point.

Proof: Since $T : X \rightarrow X$ is a generalized c-pseudo contraction then there exist $k, 0 \leq k < \frac{1}{4}$ such that for all $x, y \in X$ the following inequality holds:

$$d(Tx, Ty) \leq k\{d(Tx, x) + d(y, Ty) + d(Tx, y) + d(x, Ty)\}$$

We shall first show that $T : (X, d^s) \rightarrow (X, d^s)$ is a generalized c-contraction.

Since for any $x, y \in X$ we have

$$d^{-1}(Tx, Ty) = d(Ty, Tx)$$

$$\leq k\{d(Ty, y) + d(x, Tx) + d(Ty, x) + d(y, Tx)\}$$

$$\leq k\{d^{-1}(y, Ty) + d^{-1}(Tx, x) + d^{-1}d(x, Ty) + d^{-1}(Tx, y)\}$$

$$d^{-1}(Tx, Ty) \leq k\{d^{-1}(y, Ty) + d^{-1}(Tx, x) + d^{-1}d(x, Ty) + d^{-1}(Tx, y)\}$$

We see that $T : (X, d^{-1}) \rightarrow (X, d^{-1})$ is a generalized C-pseudo contraction therefore

$$d(Tx, Ty) \leq k\{d(Tx, x) + d(y, Ty) + d(Tx, y) + d(x, Ty)\}$$

$$\leq k\{d^s(x, Tx) + d^s(y, Ty) + d^s(Tx, y) + d^s(x, Ty)\}$$

$$\text{and } d^{-1}(Tx, Ty) \leq k\{d^{-1}(y, Ty) + d^{-1}(Tx, x) + d^{-1}(x, Ty) + d^{-1}(Tx, y)\}$$

$$\leq k \left\{ d^s(y, Ty) + d^s(x, Tx) + d^s(x, Ty) + d^s(Tx, y) \right\} \text{ for all } x, y \in X$$

Hence $d^s(Tx, Ty) \leq k \left\{ d^s(x, Tx) + d^s(y, Ty) + d^s(Tx, y) + d^s(x, Ty) \right\}$, for all $x, y \in X$

and so $T : (X, d^s) \rightarrow (X, d^s)$ is a generalized C- contraction.

By assumption (X, d) is a bi complete. Hence (X, d^s) is complete. There fore by theorem (4a) T has a unique fixed point. This completes the proof.

Corollary 4.1: Let (X, d) be a T_0 -Isbell-Complete quasi pseudo metric spaces and $T : X \rightarrow X$ be a generalized c - pseudo contraction then T has a unique fixed point.

The proof follows from the proposition 2.1

Corollary 4.2: Any generalized c- pseudo contraction on a q-spherically complete T_0 ultra quasi metric space has a unique fixed point.

The proof follows from the proposition 3.1

Theorem 4.2:

Let (X, d) be a bi complete di metric space and let $T : X \rightarrow X$ be an generalized S pseudo contraction then T has a unique fixed point

Proof: As in the previous proof it is enough to prove that $T : (X, d^s) \rightarrow (X, d^s)$ is an generalized S -contraction.

Since $T : X \rightarrow X$ be a S -pseudo contraction then there exist $k, 0 < k < \frac{1}{8}$ such that for all $x, y \in X$ the following inequality holds:

$$d(Tx, Ty) \leq k \{ d(x, Tx) + d(x, Ty) + d(y, Tx) + d(y, Ty) + d(Tx, x) + d(Ty, x) + d(Tx, y) + d(Ty, y) \}$$

We shall first show that $T : (X, d^s) \rightarrow (X, d^s)$ is a generalized C- contraction.

Since for any $x, y \in X$ we have

$$d^{-1}(Tx, Ty) = d(Ty, Tx)$$

$$d(Ty, Tx) \leq k \{ d(y, Ty) + d(y, Tx) + d(x, Ty) + d(x, Tx) + d(Ty, y) + d(Tx, y) + d(Ty, x) + d(Tx, x) \}$$

$$\leq k \left\{ \begin{aligned} & d^{-1}(Ty, y) + d^{-1}(Tx, y) + d^{-1}(Ty, x) + d^{-1}(Tx, x) \\ & + d^{-1}(y, Ty) + d^{-1}(y, Tx) + d^{-1}(x, Ty) + d^{-1}(x, Tx) \end{aligned} \right\}$$

$$d^{-1}(Tx, Ty) \leq k \left\{ \begin{aligned} & d^{-1}(x, Tx) + d^{-1}(x, Ty) + d^{-1}(y, Tx) + d^{-1}(y, Ty) \\ & + d^{-1}(Tx, x) + d^{-1}(Ty, x) + d^{-1}(Tx, y) + d^{-1}(Ty, y) \end{aligned} \right\}$$

We see that $T : (X, d^{-1}) \rightarrow (X, d^{-1})$ is a pseudo contraction.

Therefore

$$d(Tx, Ty) \leq k \{ d(x, Tx) + d(x, Ty) + d(y, Tx) + d(y, Ty) + d(Tx, x) + d(Ty, x) + d(Tx, y) + d(Ty, y) \}$$

$$d(Tx, Ty) \leq k \{ d^s(Tx, x) + d^s(x, Ty) + d^s(y, Tx) + d^s(y, Ty) + d^s(Tx, x) + d^s(Ty, x) + d^s(Tx, y) + d^s(Ty, y) \}$$

and

$$d^{-1}(Tx, Ty) \leq k \left\{ \begin{aligned} & d^{-1}(x, Tx) + d^{-1}(x, Ty) + d^{-1}(y, Tx) + d^{-1}(y, Ty) \\ & + d^{-1}(Tx, x) + d^{-1}(Ty, x) + d^{-1}(Tx, y) + d^{-1}(Ty, y) \end{aligned} \right\}$$

$$\leq k \left\{ \begin{aligned} & d^n(Tx, x) + d^n(Ty, x) + d^n(Tx, y) + d^n(Ty, y) \\ & + d^n(x, Tx) + d^n(x, Ty) + d^n(y, Tx) + d^n(y, Ty) \end{aligned} \right\} \text{ For all } x, y \in X$$

Hence

$$d^n(Tx, Tx) \leq k \left\{ \begin{aligned} & d^n(Tx, x) + d^n(Ty, x) + d^n(Tx, y) + d^n(Ty, y) \\ & + d^n(x, Tx) + d^n(x, Ty) + d^n(y, Tx) + d^n(y, Ty) \end{aligned} \right\}, \text{ for all } x, y \in X$$

and so $T : (X, d^s) \rightarrow (X, d^s)$ is a generalized s-contraction.

By assumption (X, d) is a bi complete. Hence (X, d^s) is complete. There fore by theorem (4a) T has a unique fixed point. This completes the proof.

Corollary 4.3: Let (X, d) be a T_0 -Isbell-Complete quasi pseudo metric spaces and $T : X \rightarrow X$ be a pseudo contraction then T has a unique fixed point.

The proof follows from the proposition 2.1

Corollary 4.4: Any s-pseudo contraction on a q-spherically complete T_0 ultra quasi metric space has a unique fixed point.

References

1. Agyingi CA (2013) A fixed point theorem in non-Archimedean T_0 -quasi-metric spaces. Adv Fixed Point theory 3: 667-674.
2. Petals C, Fvidalis T (1993) A fixed point theorem in non-Archimedean vector spaces. Pro Amer Math Soc118: 819-821.
3. Agyingi CA, Gaba YU (2014) A fixed point like theorem in a T_0 -ultra-quasi-metric space. Advances in Inequalities and Applications Article-ID.
4. Agyingi CA, Gaba YU (2014) Common fixed point theorem for maps in a T_0 -ultra-quasi-metric space. Sci J Math Resin press 4: 117-124.
5. Gaba YE (2014) Unique fixed point theorems for contractive maps type in T_0 quasi metric spaces. AdvFixed point theory 4: 117-125.
6. Kemajou E, PAK'unzi H, Otafudu OO (2012) The Isbell-hull of a di-space. Topology Appl 159: 2463-2475.
7. Kemajou E, PAK'unzi H, Otafudu OO (2012) The Isbell-hull of a di-space. Topology Appl 159: 2463-2475.
8. Deza MM, Deza E (2009) Encyclopedia of Distances, Springer, Berlin.
9. Bonsangue MM, Breugel FV, Rutten JMM (1995) Generalized Ultrametric Spaces: completion, topology, and power domains via the Yoneda embedding.
10. Chatterji KS, Acad CR (1972) Fixed point theorems, Bulgare Sci 25: 727-730.
11. Shukla DP, Tiwari SK (23012) Unique fixed point for s-weak contractive mappings, Gen Math 4: 28-34.