

## Computational Approach to Identify the Major Histocompatibility Complex Binding Antigenic Peptides from ‘Ascariasis’

Mishra S\* and Gomase VS\*

Department of Biotechnology, Mewar University, Chittorgarh, India

### Abstract

The *Ascaris lumbricoides* cause ‘ascariasis’ in human and its elimination is the major public concern. In this current investigation, we predicted the MHC (Major Histocompatibility Complex) class I & II binding peptides with computational approach like PSSM (Position Specific Scoring Matrices) and SVM (Support Vector Machine) algorithms. We predict the peptide binders of Cytochrome b (mitochondrion) protein from *Ascaris lumbricoides* sequence to MHC-I molecules are as 11mer\_H2\_Db, 10mer\_H2\_Db, 9mer\_H2\_Db, 8mer\_H2\_Db. Also study integrates prediction of peptide MHC class I binding; proteasomal C terminal cleavage and TAP transport efficiency by using sequence and properties of the amino acids. We also found the binding of peptides to different alleles by using Position Specific Scoring Matrix. Cytochrome B from *Ascaris lumbricoides* (365 residues long) with 357 nonamers having antigenic MHC binding peptides. PSSM based server will predict the peptide binders from sequence to MHCII molecules are as I\_Ab.p, I\_Ad.p, I\_Ag7, which are found antigenic epitopes region in Cytochrome B from *Ascaris lumbricoides*. This investigation can be useful in rational vaccine design and simultaneously increase the understanding the role of the immune system against antigenic.

**Keywords:** Ascariasis; *Ascaris lumbricoides*; Epitopes; Antigenic peptides; MHC-Binders; TapPred; PSSM; SVM; Nonamers; Cytochrome b (mitochondrion)

**Abbreviations:** STH: Soil Transmitted Helminthes; SAC: School Age Children; MDA: Mass Drug Administration; MHC I: Major Histocompatibility Complex Class I; MHC II: Major Histocompatibility Complex Class II; PSSM: Position Specific Scoring Matrices; SVM: Support Vector Machine; UniProt: The Universal Protein Resource; NCBI: National Center for Biotechnology Information; TAP: Transporter Associated with Antigen Processing

### Introduction

“Ascariasis” infection caused in humans and other mammals via intestinal round worms (*A. lumbricoides*) of genus ‘Ascaris’ with incubation period of 10-24 months in the jejunum and middle ileum of the human intestine. The adult worm are in size varies up to 15-35 cm long and appeared in color white or yellow. Female worm produces 240,000 eggs per day. The fertilized eggs turn infectious within 5-10 days when it expelled into favorable soil [1] and stay viable for up to 17 months. The route of the disease propagation is majorly from the worm contaminated food hand and soil, through ingestion of contaminated food and the opportunistic parasite get success in hatching of eggs in the small intestine. This worm infectious occurrence are also observed as a zoonotic infection in pigs and use of hog manure [2] but, in endemic areas, the transmission of this infection is basically from person to person [3].

The diseases symptoms experienced by the individuals are pneumonitis include wheezing, dyspnea, nonproductive cough, hemoptysis, and fever and eosinophilia. The larvae mature in jejunum in approximately 65 days which ultimately feeds on host (human) digestion (Figure 1). The infected children’s with nutritional conditional like marginal diet may found to be susceptible to protein, caloric, or suffer from Vitamin A deficiency, which ultimately interferes in their normal growth and causes growth [4] and the relative immunodeficiency also been observed in overexposed situation [5]. In the minimal load this worm expression remain asymptomatic whereas in the overload situation it creates severe complications. The predominant occurrence of this disease found in the warm and moist climatic geographical regions and also in those tropical and subtropical

regions where the poor sanitation and hygiene are practice. In the worldwide the estimated preponderance of this disease outbreak is recorded is 25% (0.8-1.22 billion people) [6]. Children’s are supposed to be the majorly prone to this disease of tropical and arising countries [7-8]. The differential diagnoses includes are acute pancreatitis, biliary colic and community-acquired pneumonia. The recent advanced strategy for STH infection control is preventive chemotherapy with combinational therapy of albendazole or mebendazole [9]. In another study, conducted by researcher in Chench district, where the mass drug administration (MDA) therapy is implied to most risky population and SAC as the central strategy to control soil-transmitted helminthes (STH) infection [10]. However, Identification of antigenic peptide that binds to MHC [Major Histocompatibility Complex] molecule improves the understanding of specificity of immune responses and is important for discovery of vaccines. MHC molecules are cell surface proteins that found in all vertebrate animals and are play an important role in the immune system.

### MHC class I antigen

In the process of the antigen-presentation in the proteasome, antigenic protein is cleaved into oligopeptides [11-13]. The cleaved antigenic peptide fragment were brought through TAP protein into the endoplasmic reticulum (ER) and then these peptides combines with MHC class I molecule and forms a complex [14-18]. The resultant complexes were translocated towards the surface of the antigen presenting cell, which are thereafter recognized and identified by T-cell

\*Corresponding authors: Sonu Mishra, Department of Biotechnology, Mewar University, Chittorgarh, India, Tel: +91-9560808369; E-mail: [sonumishra1014@gmail.com](mailto:sonumishra1014@gmail.com)

Virendra S Gomase, Department of Biotechnology, Mewar University, Chittorgarh, India, Tel: +91 27743967; E-mail: [gomase.viren@gmail.com](mailto:gomase.viren@gmail.com)

Received May 06, 2016; Accepted May 17, 2016; Published May 26, 2016

**Citation:** Mishra S, Gomase VS (2016) Computational Approach to Identify the Major Histocompatibility Complex Binding Antigenic Peptides from ‘Ascariasis’. J Data Mining Genomics Proteomics 7: 198. doi:10.4172/2153-0602.1000198

**Copyright:** © 2016 Mishra S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Lifecycle of the *Ascaris lumbricoides*.

and enable immune system to elicit an immune response [19-23]. Henceforth, TAP binding peptides prediction is very important for identification of the MHC class-I restricted T cell epitopes.

### Proteasomal degradation

The process of antigen presentation the proteasomal degradation is important step to regulate and maintain the equilibrium among intracellular proteins [24]. Through the action of the proteinase inside the antigenic protein are cleaved into oligopeptides [25] and then after these oligopeptides binds to the TAP, which later on transports peptides into the ER.

### Transport of peptide by TAP

TAP is heterodimeric transmembrane protein, and belongs from ABC transporter family. This transports antigenic peptide to ER [26]. Study suggest that most of the MHC binding peptides are unable to diffuse across membrane, but TAP able to carry them inside the ER where it binds to MHC class I molecules. These MHC-peptide complexes will be translocate on the surface of antigen presenting cells [21] and are recognized by T-cell receptors to elicit an immune response.

### MHC class II antigen

The prediction of peptides binding to a MHC class II molecule is difficult due to different side chains and longer length in the extracellular antigen presentation [27-29]. In the MHC class II antigen presentation process, antigenic protein is ingested by antigen-presenting cells via endocytosis or phagocytosis process, and the after cleaved by cathepsins (a class of protease) into oligopeptides in the endosomes, than are fuse with lysosomes containing MHC class II molecules [30] and present them at the cell surface for recognition by T cells [31-39]. Where T helper cells trigger an immune response by inflammation and swelling due to phagocytes or may lead to an antibody-mediated immune response via B-cell activation. Since MHCs have a key role in immune system by stimulating cellular and humoral immunity against antigenic peptide are used for controlling specific immunological processes by creating peptides to bind to specific MHC alleles and this binding affinity to specific peptides are used for designing synthetic peptide vaccines [40-43].

### Methodology

#### Retrieval of specific data from database

The protein data of cytochrome b (mitochondrion) from *Ascaris lumbricoides* was retrieved from www.ncbi.nlm.nih.gov, UniProt databases for the further analysis [44-47].

### Antigenicity identification of protein

The identification of the antigenicity of the Cytochrome b (mitochondrion) protein are predicted through utilizing the different types of the computational tools, methods and relevant scales like Hopp and Woods, Welling, Parker, Bepipred, Kolaskar and Tongaonkar antigenicity methods [48-59] and identified the antigenic peptide which are capable to eliciting an antibody response.

### MHC binding peptide prediction

The earlier sequence patterns method for prediction of MHC molecules have proven to be too elementary and also the complexity of the binding motif cannot be precisely interpreted by the few residues present in the pattern [60]. MHC binding peptide is predicted using neural networks trained on C terminals of known epitopes. By using RANKPEP, we predict peptide binders to MHCI molecules from protein sequences or sequence alignments using Position Specific Scoring Matrices (PSSMs) whose C terminal end is likely to be the result of proteosomal cleavage [61-64].

### Antigenic peptides prediction by cascade SVM based TAP pred method

By using TAPPred we predict TAP binders on the basis of sequence and the properties of amino acids [65]. We found the MHCI binding regions, the binding affinity of protein from *Ascaris lumbricoides* having 365 amino acids long residues with 357 nonamers.

### Results

#### Retrieval of the protein sequence from database

The Cytochrome b protein from *Ascaris lumbricoides* [gi|319656158|gb|ADV58572.1|], contain a long residue of 365 amino acids with 357 nonamers.

```
>gi|319656158|gb|ADV58572.1|cytochrome b (mitochondrion) [Ascaris lumbricoides]
MKLDFVNSMVSLSKSKVLTYGWVNFSGSMLGMVLGFGQILGTGFLAFYYNSNDGALAFLSVQYIMY
EVNFGWIFRVSHFNGASMFIFLYLHLFKGMFFMSYRLKKAWSGVILLVMMEAFMGYVLLV
WAQMSFWASVVTLSLSPVWVGFVAIVTCIWSGFTVSSATLKKFFVHLVLPWGLLFLVLLHLVF
LHETRSTSKSYCHGHYDKVCFSPYVWKDFLNVVWVVFIFLGFPLLDGPEMFIESDPMMSF
VHIVPEWYFLFAYAILRAIPNKVLGVVSLFASILVLVVFLVNNYVSVMSKLNKFLVVFVIFVLV
VLSWLGQCLVEDPFVLSMVFSLYFFVIFLLFLVYFVGRVFM
```

### Prediction of antigenic peptides

In antigenic prediction, we focused on the area of greatest local Hydrophilicity through antigenic determinants. In Hopp-Woods scale

Hydrophilicity Prediction method, result data found high in Position: 197 with Score: 0.956 (max) 194-TRSTSK-200 in a protein, assuming the probability of antigenic determinants would be exposed on the surface of the protein and hence, there might be chances to be located in hydrophilic regions (Figure 2). Welling et al. antigenicity plot provides value as the log of the quotient between percentage in average proteins and percentage in a sample of known antigenic regions. The prediction result found highest in Position: 189 Score: 0.841 (max) 186-LLHLV FL-192 (Figure 3). We also study Hydrophobicity plot of HPLC/Parker Hydrophilicity Prediction Result Data found in Position: 197 (Residue: S) i.e., 194-ETRSTSK-200 with highest score: 5.871 (Figure 4), BepiPred predicts the location of linear B-cell epitopes Result found in Position: 197 (Residue: S) with highest score: 1.023 (193-LVFLHETR-200) (Figure 5), Kolaskar and Tongaonkar antigenicity methods (Figures 6 and 7), Table 1). Predicted peptides result found 4-DFVNSM VVSLPSSKVL T-20,28-MLGMVLGFQILTGTFLAFY-46,52-ALAFLSVQYIMY-63,69-WIFRVSH-75,80-SMFFIFLYLHLFK-92,100-

| n  | Strat | Sequence  | End Position |
|----|-------|---|--------------|
| 1  | 4     | DFVNSM VVSLPSSKVL T   | 20           |
| 2  | 28    | MLGMVLGFQILTGTFLAFY   | 46           |
| 3  | 52    | ALAFLSVQYIMY  | 63           |
| 4  | 69    | WIFRVSH   | 75           |
| 5  | 80    | SMFFIFLYLHLFK   | 92           |
| 6  | 100   | RLKKAWVSGVILLVM   | 116          |
| 7  | 120   | FMGYVLVWA   | 128          |
| 8  | 132   | FWASVVITSLLSVIPVWGFAIVTCIWSGFTVSSATLKFFVLHF<br>LVPWGLLFLVLLHLVFLH | 193          |
| 9  | 199   | SKSYCHGHYDKVCFSPYVWKDFLNVVVWFVFIFFSLGFPFL                         | 241          |
| 10 | 253   | NNSPVHIVPEWYFLFAYAILRAIPNKVLGVVSLFASILVLVF<br>VLVNNYVMS           | 306          |
| 11 | 308   | LMKFLVFVIFVLVLSWLQCLVEDPFVFLSMVFSFLYFFVI<br>FLLFLVYFVG            | 361          |

There are 11 antigenic determinants in your sequences.

Table 1: The 11 antigenic determinants of cytochrome b (mitochondrion) protein.




Figure 2: Hydrophobicity plot (result data found high in position: 197 with score: 0.956 (max) 194-TRSTSK-200) [49].




Figure 3: Hydrophobicity plot (result found highest in position: 189 Score: 0.841 (max) 186-LLHLV FL-192) [50].




Figure 4: Hydrophobicity plot of HPLC (result data found in position: 197 (Residue: S) i.e., 194-ETRSTSK-200 with highest score: 5.871) [51].




Figure 5: Bepipred linear epitope prediction plot (result found in position: 197 (Residue: S) with highest score: 1.023 and sequence is 193-LVFLHETR-200).




Figure 6: Antigenicity plot [53].

RLKKAWSGIVILLVM-116,120-FMGYVLVWA-128,132-FWASV  
VITSLLSVIPVWGFAIVTCIWSGFTVSSATLKFFVLHFLVPWGLL  
FLVLLHLVFLH-193,199-SKSYCHGHYDKVCFSPYVWKDFLNV  
VVWFVIFFLSGFPFL-241,253-MMSPVHIVPEWYFLFAYAILRA-  
IPNKVLGVVSLFASILVLVVFLVNNYVSMS-306,308-LNKFL  
VFVFIFVLVLSWLGQCLVEDPFVFLSMVFSFLYFFVIFLLFLVY  
YFVG-361 and there might be chances that the predicted antigenic

fragments can bind to MHC molecule and it can be the first chokepoints in synthetic vaccine design.

### MHC binding peptide prediction

We found the binding of peptides to a number of different alleles using Position Specific Scoring Matrix. MHC molecules are cell surface proteins, which actively participate in host immune responses to almost


all antigens. We have been able to predict MHC-I peptide binders of cytochrome b (mitochondrion) from *Ascaris lumbricoides*. We found predicted MHC-I peptide binders of protein for Matrix: 8mer\_H2\_Db.p.mtx, Consensus: QNWNCCCTI, Optimal Score: 52.494, Binding Threshold: 33.04; Matrix: 9mer\_H2\_Db.p.mtx, Consensus: FCIHNCDYIM, Optimal Score: 50.365, Binding Threshold: 17.96; Matrix: 10mer\_H2\_Db.p.mtx, Consensus: SGYYNFFWCL, Optimal Score: 58.858, Binding Threshold: 41.32; Matrix: 11mer\_H2\_Db.p.mtx, Consensus: CGVYNFYCCY, Optimal Score: 79.495, Binding Threshold: 56.96 (Table 2) and MHC-II peptide binders for Matrix: I\_Ab.p.mtx, Consensus: YYAPWCNNA, Optimal Score: 35.632, Binding Threshold: 9.52; Matrix: I\_Ad.p.mtx, Consensus: QMVHAAHAE, Optimal Score: 53.145, Binding Threshold: 7.10; Matrix: I\_Ag7.p.mtx, Consensus: WYAHAFKYV, Optimal Score: 40.873, Binding Threshold: 7.54 for MHC II allele (Table 3) was tested and opted the result.

### Antigenic peptides prediction by cascade SVM based TAP pred method

We also use a cascade SVM based TAPPred method which found 105 High affinity TAP Transporter (Transporter peptide regions) (Tables 3 and 4) peptide regions which represents predicted TAP binders residues which occur at N and C termini from *Ascaris lumbricoides* antigen cytochrome c oxidase subunit I (mitochondrion). The important transporter that transports antigenic peptides from cytosol to ER is TAP. It binds and translocate the selective antigenic peptides from cytosol to ER for MHC molecules specific binding. By using the approach of jackknife validation test the correlation coefficient of 0.88 was achieved.

### Conclusion

The antigenic proteins cytochrome b from *Ascaris lumbricoides* involved in different multiple antigenic components to direct and



| MHC-I Allele | Rank | POS. | N   | Sequence    | C   | MW (Da) | SCORE  | % OPT. |
|--------------|------|------|-----|-------------|-----|---------|--------|--------|
| 8mer_H2_Db   | 1    | 155  | IVT | CIWSGFTV    | SSA | 871.05  | 22.483 | 42.83% |
| 8mer_H2_Db   | 2    | 146  | SVI | PVWGFAIV    | TCI | 847.06  | 14.929 | 28.44% |
| 8mer_H2_Db   | 3    | 177  | HFL | VPWGLLFL    | VLL | 903.17  | 11.733 | 22.35% |
| 8mer_H2_Db   | 4    | 165  | VSS | ATLKFFV     | LHF | 954.18  | 11.706 | 22.30% |
| 8mer_H2_Db   | 5    | 224  | FLN | VVWFVFI     | FFS | 967.25  | 11.667 | 22.23% |
| 9mer_H2_Db   | 1    | 73   | IFR | VSHFNGASM   | FFI | 931.03  | 21.024 | 41.74% |
| 9mer_H2_Db   | 2    | 20   | KVL | TYGWNFGSM   | LGM | 1021.14 | 15.731 | 31.23% |
| 9mer_H2_Db   | 3    | 296  | VVF | VLVNNYVSV   | MSK | 988.14  | 15.156 | 30.09% |
| 9mer_H2_Db   | 4    | 233  | FIF | FSLGFPFL    | GDP | 1022.27 | 13.192 | 26.19% |
| 9mer_H2_Db   | 5    | 202  | SKS | YCHGHYDKV   | CFS | 1103.22 | 12.345 | 24.51% |
| 10mer_H2_Db  | 1    | 77   | SHF | NGASMFIFL   | YLH | 1128.36 | 16.914 | 28.74% |
| 10mer_H2_Db  | 2    | 299  | VLV | NNYVSVMSKL  | NKF | 1136.32 | 10.744 | 18.25% |
| 10mer_H2_Db  | 3    | 296  | VVF | VLVNNYVSV   | SKL | 1119.33 | 10.308 | 17.51% |
| 10mer_H2_Db  | 4    | 20   | KVL | TYGWNFGSM   | GMV | 1134.3  | 8.093  | 13.75% |
| 10mer_H2_Db  | 5    | 73   | IFR | VSHFNGASMF  | FIF | 1078.21 | 8.077  | 13.72% |
| 11mer_H2_Db  | 1    | 320  | IFV | LVLSWLQCL   | VED | 1189.51 | 13.592 | 17.10% |
| 11mer_H2_Db  | 2    | 115  | LLL | VMMEAFMGYVL | VWA | 1272.6  | 13.422 | 16.88% |
| 11mer_H2_Db  | 3    | 295  | LVV | FVLVNNYVSV  | SKL | 1266.51 | 12.705 | 15.98% |
| 11mer_H2_Db  | 4    | 20   | KVL | TYGWNFGSMLG | MVL | 1191.35 | 12.091 | 15.21% |
| 11mer_H2_Db  | 5    | 339  | VFL | SMVFSFLYFFV | IFL | 1368.67 | 11.309 | 14.23% |

\*The RANKPEP consists of a list of selected peptides binding potential (score) to the MHC molecule from the query given at a selected threshold. Peptides shown here contain a C-terminal residue that is predicted to be the result of proteasomal cleavage and also focus on the prediction of conserved epitopes that help to avoid immune evasion resulting from mutation. Proteasomal cleavage options are only applied to the prediction of MHC-I-restricted peptides.

Table 2: Promiscuous MHC ligands, having C-terminal ends are proteasomal cleavage sites of *Ascaris lumbricoides*.

| MHC-II Allele | Rank | POS. | N   | SEQUENCE  | C   | MW (Da) | SCORE  | % OPT. |
|---------------|------|------|-----|-----------|-----|---------|--------|--------|
| MHC-II I_Ab   | 1    | 150  | VWG | FAIVTCIWS | GFT | 998.24  | 18.998 | 53.32% |
| MHC-II I_Ab   | 2    | 123  | FMG | YVLVWAQMS | FWA | 1055.29 | 14.317 | 40.18% |
| MHC-II I_Ab   | 3    | 46   | LAF | YYSNDGALA | FLS | 955     | 13.721 | 38.51% |
| MHC-II I_Ab   | 4    | 132  | QMS | FWASVVITS | LLS | 968.15  | 12.184 | 34.19% |
| MHC-II I_Ab   | 5    | 175  | VLH | FLVPWGLLF | LVL | 1050.35 | 11.623 | 32.62% |
| MHC-II I_Ad   | 1    | 269  | LFA | YAILRAIPN | KVL | 1012.23 | 17.555 | 33.03% |
| MHC-II I_Ad   | 2    | 158  | CIW | SGFTVSSAT | LKF | 837.88  | 15.031 | 28.28% |
| MHC-II I_Ad   | 3    | 69   | NFG | WIFRVSHFN | GAS | 1164.37 | 14.098 | 26.53% |
| MHC-II I_Ad   | 4    | 267  | YFL | FAYAILRAI | PNK | 1019.27 | 11.974 | 22.53% |
| MHC-II I_Ad   | 5    | 280  | NKV | LGVVSLFAS | ILV | 874.05  | 10.238 | 19.26% |
| MHC-II_Ag7    | 1    | 86   | FIF | LYLHLFKGM | FFM | 1103.39 | 18.964 | 46.40% |
| MHC-II_Ag7    | 2    | 115  | LLL | VMMEAFMGY | VLV | 1060.31 | 10.534 | 25.77% |
| MHC-II_Ag7    | 3    | 270  | FAY | AILRAIPNK | VLG | 977.22  | 10.318 | 25.24% |
| MHC-II_Ag7    | 4    | 30   | SML | GMVLGFQIL | TGT | 959.21  | 10.11  | 24.74% |
| MHC-II_Ag7    | 5    | 216  | SPE | YVWKDFLNV | VVW | 1142.35 | 9.304  | 22.76% |

Table 3: Prediction of MHCII ligands all rows predicted binders to the MHC-II Allele i.e., MHC-II I\_Ab, MHC-II I\_Ad, MHC-II I\_Ag7.

| Peptide Rank | Start Position | Sequence  | Score | Predicted Affinity |
|--------------|----------------|-----------|-------|--------------------|
| 1            | 201            | SYCHGHYDK | 8.647 | High               |
| 2            | 144            | VIPVVGFAL | 8.64  | High               |
| 3            | 319            | VLVLSVVLG | 8.64  | High               |
| 4            | 301            | YVSVMSKLN | 8.624 | High               |
| 5            | 160            | FTVSSATLK | 8.628 | High               |
| 6            | 56             | LSVQIMYE  | 8.627 | High               |
| 7            | 127            | WAQMSFWAS | 8.625 | High               |
| 8            | 41             | TFLAFYYSN | 8.634 | High               |
| 9            | 213            | PSEYVWKDF | 8.624 | High               |
| 10           | 217            | VVWKDFLNV | 8.624 | High               |

\*TAPPred showing Cascade SVM based High affinity TAP Binders sites, their sequence, rank, position and scores are displayed in the tabular output are to be found 15 High affinity TAP Transporter peptide regions which represents predicted TAP binders residues which occur at N and C termini Cytochrome B from *Ascaris lumbricoides*.

Table 4: Cascade SVM based high affinity TAP binders.

empower the immune system to protect the host from the antigenic peptide. Major histocompatibility complexes (MHC-I and MHC-II) display specificity to bind with their respective epitopes. MHC class molecules are cell surface proteins that take active part in host immune reactions to response almost all antigens. The predicted result for the MHCII-I-Ab peptide regions are, 150-FAIVTCIWS, 123-YVLVWAQMS, 46-YYSNDGALA (optimal score is 35.632); MHCII-I\_Ad peptide regions, 158-SGFTVSSAT, 69-WIFRVSHFN, 267-FAYAILRAI (optimal score is 53.145); MHC-II I\_Ag7 peptide regions 86-LYLHLFKGM, 115-VMMEAFMGY, 270-AILRAIPNK (optimal score is 40.873) which represented the predicted binders from Cytochrome b (mitochondrion) protein. The complete investigational outcomes is an encouraging breakthrough with very high accuracies. The knowledge of the immune responses to an antigen protein (cytochrome b) clear that the whole protein is not necessary for raising the immune response, but a small fragment of antigen can induce immune response against whole antigen. This means the increase in affinity of MHC binding peptides may result in enhancement of immunogenicity of cytochrome b from *Ascaris lumbricoides*, hence are helpful *in silico* to design and develop highly predictive computational tools for the identification of T-cell epitopes. In future this methodology will be useful in cellular immunology, vaccine design, immunodiagnostics, immunotherapeutic and molecular understanding of susceptibility of autoimmune. Finally, accurate prediction remains vital for the future to design synthetic peptide vaccine.

## References

- Peng W, Zhou X, Gasser RB (2003) *Ascaris* egg profiles in human faeces: biological and epidemiological implications. *Parasitology* 127: 283-290.
- Hoenigl M, Seeber K, Valentin T, Zollner-Schwetz I, Krause R (2012) Pulmonary ascariasis in patients from wealthy countries: shift in epidemiology? *Int J Infect Dis* 16: e888.
- Centers for Disease Control and Prevention (CDC) (2013) Notes from the field: ascariasis associated with pig farming - Maine, 2010-2013. *MMWR Morb Mortal Wkly Rep* 62: 413.
- Peng W, Yuan K, Hu M, Gasser RB (2007) Recent insights into the epidemiology and genetics of *Ascaris* in China using molecular tools. *Parasitology* 134: 325-330.
- Le Hesran JY, Akiana J, Ndiaye HM, Dia M, Senghor P, et al. (2004) Severe malaria attack is associated with high prevalence of *Ascaris lumbricoides* infection among children in rural Senegal. *Trans R Soc Trop Med Hyg* 98: 397-399.
- Dold C1, Holland CV (2011) *Ascaris* and ascariasis. *Microbes Infect* 13: 632-637.
- Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, et al. (2006) Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* 367: 1521-1532.
- Walker M, Hall A, Basáñez MG (2011) Individual predisposition, household clustering and risk factors for human infection with *Ascaris lumbricoides*: new epidemiological insights. *PLoS Negl Trop Dis* 5: e1047.
- Zhou Q, Liu CF, Zhang LX, Zhou H, Chen YD (2015) [Research progress in soil-transmitted helminth infection control among children at home and abroad]. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* 27: 431-435.
- Speich B, Moser W, Ali SM (2016) Efficacy and reinfection with soil-transmitted helminths 18-weeks post-treatment with albendazole-ivermectin, albendazole-mebendazole, albendazole-oxantel pamoate and mebendazole. *Parasit Vectors* 9: 123.
- Zerdo Z, Yohanes T, Tariku B (2016) Soil-Transmitted Helminth Reinfection and Associated Risk Factors among School-Age Children in Chench District, Southern Ethiopia: A Cross-Sectional Study. *J Parasitol Res* 2016.
- Quintero-Hernández V, Jiménez-Vargas JM, Gurrola GB, Valdivia HH, Possani LD (2013) Scorpion venom components that affect ion-channels function. *Toxicon* 76: 328-342.
- Holzthütter HG, Frömmel C, Kloetzel PM (1999) A theoretical approach towards the identification of cleavage-determining amino acid motifs of the 20 S proteasome. *J Mol Biol* 286: 1251-1265.
- Rock KL, Gramm C, Rothstein L, Clark K, Stein R, et al. (1994) Inhibitors of the proteasome block the degradation of most cell proteins and the generation of peptides presented on MHC class I molecules. *Cell* 78: 761-771.
- Bhasin M, Raghava GP (2005) Pcleavage: an SVM based method for prediction of constitutive proteasome and immunoproteasome cleavage sites in antigenic sequences. *Nucleic Acids Res* 33: W202-W207.
- Vyas JM, Van der Veen AG, Ploegh HL (2008) The known unknowns of antigen processing and presentation. *Nat Rev Immunol* 8: 607-618.

17. Kelly A, Powis SH, Kerr LA, Mockridge I, Elliott T, et al. (2014) Assembly and function of the two ABC transporter proteins encoded in the human major histocompatibility complex. *Nature* 355: 641-644.
18. Lautscham G, Rickinson A, Blake N (2003) TAP-independent antigen presentation on MHC class I molecules: lessons from Epstein-Barr virus. *Microbes Infect* 5: 291-299.
19. Nussbaum AK, Kuttler C, Tenzer S, Schild H (2003) Using the World Wide Web for predicting CTL epitopes. *Curr Opin Immunol* 15: 69-74.
20. Lankat-Buttgereit B, Tampe R (2002) The transporter associated with antigen processing: Function and implications in human diseases. *Physiol Rev* 82: 187-204.
21. Rock KL, York A, Goldberg AL (2004) Post-proteasomal antigen processing for major histocompatibility complex class I presentation. *Nat Immunol* 5: 670-677.
22. Hämmerling GJ, Vogt AB, Kropshofer H (1999) Antigen processing and presentation--towards the millennium. *Immunol Rev* 172: 5-9.
23. Bhasin M1, Raghava GP (2007) A hybrid approach for predicting promiscuous MHC class I restricted T cell epitopes. *J Biosci* 32: 31-42.
24. Brusic V1, Bajic VB, Petrovsky N (2004) Computational methods for prediction of T-cell epitopes--a framework for modelling, testing, and applications. *Methods* 34: 436-443.
25. Reidesel H, Kolbeck B, Schmetzer O, Knapp EW (2004) Peptide binding at class I major histocompatibility complex scored with linear functions and support vector machines. *Genome Informatics* 15: 198-212.
26. Princiotta MF, Finzi D, Qian SB, Gibbs J, Schuchmann S, et al. (2003) Quantitating protein synthesis, degradation, and endogenous antigen processing. *Immunity* 18: 343-354.
27. Procko E, Gaudet R (2009) Antigen processing and presentation: TAPping into ABC transporters. *Curr Opin Immunol* 21: 84-91.
28. Yewdell JW, Reits E, Neefjes J (2003) Making sense of mass destruction: quantitating MHC class I antigen presentation. *Nat Rev Immunol* 3: 952-961.
29. Stern LJ, Wiley DC (1994) Antigenic peptide binding by class I and class II histocompatibility proteins. *Behring Inst Mitt* 94: 1-10.
30. Hammer J, Bono E, Gallazzi F, Belunis C, Nagy Z, et al. (1994) Precise prediction of major histocompatibility complex class II-peptide interaction based on peptide side chain scanning. *J Exp Med* 180: 2353-2358.
31. Jardetzky TS, Brown JH, Gorga JC, Stern LJ, Urban RG, et al. (1996) Crystallographic analysis of endogenous peptides associated with HLA-DR1 suggests a common, polyproline II-like conformation for bound peptides. *Proc Natl Acad Sci USA* 93: 734-738.
32. Cresswell P (1994) Assembly, transport, and function of MHC class II molecules. *Annu Rev Immunol* 12: 259-293.
33. Rudolph MG, Stanfield RL, Wilson IA (2006) How TCRs bind MHCs, peptides, and coreceptors. *Annu Rev Immunol* 24: 419-466.
34. Nielsen M, Lund O, Buus S, Lundegaard C (2010) MHC class II epitope predictive algorithms. *Immunology* 130: 319-328.
35. Nielsen M, Lundegaard C, Wornig P, Hvid CS, Lambeth K, et al. (2004) Improved prediction of MHC class I and class II epitopes using a novel Gibbs sampling approach. *Bioinformatics* 20: 1388-1397.
36. Murugan N, Dai Y (2005) Prediction of MHC class II binding peptides based on an iterative learning model. *Immunome Res* 1: 6.
37. Salomon J, Flower DR (2006) Predicting Class II MHC-Peptide binding: a kernel based approach using similarity scores. *BMC Bioinformatics* 7: 501.
38. Bordner AJ, Mittelmann HD (2010) Prediction of the binding affinities of peptides to class II MHC using a regularized thermodynamic model. *BMC Bioinformatics* 11: 41.
39. Nielsen M, Lundegaard C, Lund O (2007) Prediction of MHC class II binding affinity using SMM-align, a novel stabilization matrix alignment method. *BMC Bioinformatics* 8: 238.
40. Gomase VS, Chitlange NR (2002) Sensitive Quantitative Predictions of MHC Binding Peptides and Fragment Based Peptide Vaccines from *Taeniocrassiceps*. *J Vaccines Vaccin* 3: 131.
41. Changbhale SS, Chitlange NR, Gomase VS, Kale KV (2012) An Immunoinformatics Approach to Design Synthetic Peptide Vaccine from *Dendroaspis polylepis* Dendrotoxin-K (DTX-K). *Journal of Environmental & Analytical Toxicology* 2: 157.
42. Reche PA, Glutting JP, Zhang H, Reinherz EL (2004) Enhancement to the RANKPEP resource for the prediction of peptide binding to MHC molecules using profiles. *Immunogenetics* 56: 405-419.
43. Gomase VS, Chitlange NR, Changbhale SS, Kale KV (2013) Prediction of *Brugia malayi* antigenic peptides: candidates for synthetic vaccine design against lymphatic filariasis. *Protein Pept Lett* 20: 864-887.
44. Larsen JE, Lund O, Nielsen M (2006) Improved method for predicting linear B-cell epitopes. *Immunome Res* 2: 2.
45. <http://www.ncbi.nlm.nih.gov>
46. Sayers EW (2012) Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 40: D13-D25.
47. Bairoch A, Apweiler R, Wu CH, Barker WC, Boeckmann B, et al. (2005) The Universal Protein Resource (UniProt). *Nucleic Acids Res* 33: D154-D159.
48. Hopp TP, Woods KR (1981) Prediction of protein antigenic determinants from amino acid sequences. *Proc Natl Acad Sci USA* 78: 3824-3828.
49. Welling GW, Weijer WJ, van der Zee R, Welling-Wester S (1985) Prediction of sequential antigenic regions in proteins. *FEBS Lett* 188: 215-218.
50. Parker KC, Bednarek MA, Coligan JE (1994) Scheme for ranking potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J Immunol* 152: 163-175.
51. Kolaskar AS, Tongaonkar PC (1990) A semi-empirical method for prediction of antigenic determinants on protein antigens. *FEBS Lett* 276: 172-174.
52. Gomase VS, Kale KV, Chikhale NJ, Changbhale SS (2007) Prediction of MHC binding peptides and epitopes from alfalfa mosaic virus. *Curr Drug Discov Technol* 4: 117-215.
53. Gomase VS, Kale KV (2008) "In silico prediction of epitopes: a new approach for fragment based viral peptide vaccines". *Int J Applied Computing* 1: 39-46.
54. Gomase VS, Kale KV (2008) "Approach of proteomics system architecture in plant virus's database". *Int. J Applied Computing* 1: 33-38.
55. Gomase VS, Chitlange NR (2012) Sensitive Quantitative Predictions of MHC Binding Peptides and Fragment Based Peptide Vaccines from *Taeniocrassiceps*. *Journal of Vaccines and Vaccination* 3: 131.
56. Gomase VS, Chitlange NR (2012) Microbial Proteomics Approach for Sensitive Quantitative Predictions of MHC Binding Peptide from *Taenia* ovis. *J Data Mining Genomics Proteomics* 3: 121.
57. Mishra Sonu, Virendra S Gomase (2015) Prediction of antigenic epitope from *D. medinensis*: new paradigm of synthetic vaccine development. International Conference on "Recent Research Development in Environment, Social Sciences and Humanities" (ICRRDESH-15) pp: 103-107.
58. Mishra Sonu, Gomase VS (2015) Analysis of hydrophobicity and antigenic epitope prediction from *D. Medinensis*. International Conference on "Technologies for Sustainability-Engineering, Information Technology, Management and the Environment" (SUSTECH-15) pp: 44-51.
59. Mishra Sonu and, Gomase VS (2015) Analysis of Hydrophobicity and Antigenicity of Heat Shock Protein 70 from GWD. 2<sup>nd</sup> International Conference on "Recent Innovations in science, Engineering and Management" (ICRISEM-15) pp: 25-33.
60. Ruppert J, Sidney J, Celis E, Kubo RT, Grey HM, Sette A (1993) Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules. *Cell* 74: 929-937.
61. Reche PA, Glutting JP, Reinherz EL (2002) Prediction of MHC class I binding peptides using profile motifs. *Hum Immunol* 63: 701-709.
62. Reche PA, Reinherz EL (2003) Sequence variability analysis of human class I and class II MHC molecules: functional and structural correlates of amino acid polymorphisms. *J Mol Biol* 331: 623-641.
63. Craiu A, Akopian T, Goldberg A, Rock KL (1997) Two distinct proteolytic processes in the generation of a major histocompatibility complex class I-presented peptide. *Proc Nat Acad Sci U S A* 94: 10850-10855.
64. Pieters J (2000) MHC class II-restricted antigen processing and presentation. *Adv Immunol* 75: 159-208.
65. Bhasin M, Raghava GP (2004) Analysis and prediction of affinity of TAP binding peptides using cascade SVM. *Protein Sci* 13: 596-607.