

## Cord Blood with Low Cell Count: Re-Use, Rather than Discard

Jie Tan J\*

Advanced Medical and Dental Institute, Universiti Sains Malaysia, Penang, Malaysia

\*Corresponding author: Jie Tan J, Advanced Medical and Dental Institute, Universiti Sains Malaysia, Bertam 13200 Kepala Batas, Penang, Malaysia, Tel: +60 4-6533888; E-mail: jjtan@usm.my

Rec date: August 26, 2017; Acc date: August 28, 2017; Pub date: September 2, 2017

Copyright: © 2017, Tan JJ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Tan JJ (2017) Cord Blood with Low Cell Count: Re-Use, Rather than Discard. Single Cell Biol 6: 167.

### Short Communication

Single cell technology has been widely used in developmental and stem cell biology. Umbilical Cord Blood (UCB) has been accepted as a rich source of stem cells for treating various hematological diseases. Its unique therapeutic potential has sparked enormous scientific interest and business opportunity to establish cord blood banks to cryopreserve UCB cells for future use.

Umbilical Cord Blood (UCB) has been accepted as a rich source of stem cells for treating various hematological diseases. Its unique therapeutic potential has sparked enormous scientific interest and business opportunity to establish cord blood banks to cryopreserve UCB cells for future use. However, the clinical efficacy with regards to improved engraftment and reduced graft failure is reliant upon cell dosage, measured by the total nucleated cell count, the proportion of CD34+ cells, the colony-forming units, and the total volume of the each transplanted UCB unit [1,2]. This finding has also become the basis of cryopreservation criteria in most public cord blood banks after UCB collection from donors, as an established checkpoint prior to further processing for cryopreservation [3]. UCB units that fail to meet such criteria are discarded. Although the checkpoint criteria mitigate risk and further costs, unfortunately about 36% to 39% of the total collected UCB is discarded, which incurs substantial economics and depletion of resources [3-5].

It is likely that unwanted UCB units can be re-directed to laboratories for further processing and manipulations to improve their quality, making them favorable for clinical use instead of disposal. Studies have shown that mesenchymal stem cells (MSCs), a subset of plastic-adherent, multipotent regenerative cells which are capable of differentiation into osteo-, chondro- and adipo- lineages, can also be isolated from UCB in addition to CD34+ hematopoietic stem cells (HSCs) [6]. MSCs are not only a proven valuable therapeutic biologic source to regenerate tissues, but they also hold the key to address the cell dose in each UCB unit. Evidence has shown that the total nucleated cells in UCB can be markedly increased by co-culturing with MSCs [7, 8]. Such manipulation does not seem to affect the function of CD34+ HSCs, including its red blood cell differentiation capability [7]. Moreover, co-transplantation of UCB and expanded UCB with MSCs in co-culture in patients with hematological cancers has revealed a shorter neutrophil and platelet recovery time, as compared to patients who received two unmanipulated UCB units [9]. These results suggest an alternate transplantation strategy in lieu of UCB with low cell dose.

With current advances in stem cell technologies, the use of UCB is no longer limited to treating blood disorders. Rather, UCB can now

reprogrammed to generate induced pluripotent stem (IPS) cells [10], which serve as precursors for differentiating multiple cell types necessary for most tissue engineering and regenerative medicine efforts. Hence, collaborations between UCB banks and research institutions should be fostered to exploit UCB, and it should be a better option than discarding potentially life-saving, regenerative cells to the trash.

### References

- Rodrigues CA, Sanz G, Brunstein CG, Sanz J, Wagner JE, et al (2009) Analysis of risk factors for outcomes after unrelated cord blood transplantation in adults with lymphoid malignancies: A study by the Eurocord-Netcord and lymphoma working party of the European group for blood and marrow transplantation. *J Clin Oncol* 27: 256-263.
- Page KM, Zhang L, Mendizabal A, Wease S, Carter S, et al (2011) Total colony-forming units are a strong, independent predictor of neutrophil and platelet engraftment after unrelated umbilical cord blood transplantation: a single-center analysis of 435 cord blood transplants. *Biol Blood Marrow Transplant* 17: 1362-1374.
- Solves P, Perales A, Mirabet V, Blasco I, Blanquer A, et al (2004) Carbonell-Uberos, M. Angeles Soler, Optimizing donor selection in a cord blood bank. *Eur J Haematol* 72: 107-112.
- Kurtzberg J, Cairo MS, Fraser JK, Baxter-Lowe L, Cohen G, et al (2005) Results of the cord blood transplantation (COBLT) study unrelated donor banking program. *Transfusion* 45: 842-855.
- Bart T, Boo M, Balabanova S, Fischer Y, Nicoloso G, et al (2013) Impact of selection of cord blood units from the United States and swiss registries on the cost of banking operations. *Transfus Med Hemother* 40: 14-20.
- Yoshioka S, Miura Y, Iwasa M, Fujishiro A, Yao H, et al (2015) Isolation of mesenchymal stromal/stem cells from small-volume umbilical cord blood units that do not qualify for the banking system. *Int J Hematol* 102: 218-229.
- Lau SX, Leong YY, Ng WH, Ng AWP, Ismail IS, et al (2017) Human mesenchymal stem cells promote CD34+ hematopoietic stem cell proliferation with preserved red blood cell differentiation capacity. *Cell Biol Int* 41: 697-704.
- Robinson SN, Niu JT, Yang H, McMannis JD, Karandish S, et al (2006) Superior *ex vivo* cord blood expansion following co-culture with bone marrow-derived mesenchymal stem cells. *Bone Marrow Transplant* 37: 359-366.
- Lima MD, McNiece I, Robinson SN, Munsell M, Eapen M, et al (2012) Cord-blood engraftment with *ex vivo* mesenchymal-cell coculture. *N Engl J Med* 367: 2305-2315.
- Hu K, Yu J, Suknuntha K, Tian S, Montgomery K, et al (2011) Efficient generation of transgene-free induced pluripotent stem cells from normal and neoplastic bone marrow and cord blood mononuclear cells. *Blood* 117: 109-119.