

Current Perspective on Hydrogen Sulfide: Stinky Outside Precious Inside

Sangita Singh*

DuPont, DuPont Knowledge Center, Hyderabad, India

Abstract

Hydrogen sulfide is the newest member of gasotransmitter family. Endogenous production of hydrogen sulfide in brain was first reported in 1996. Since then a great deal of interest has been paid to understand various physiological roles of this molecule, still there are a lot of gaps that need to be addressed. This article provides author's perspective on research accomplishments in this area briefly, gaps and future opportunities, which may transform the area of medicine.

Keywords: Hydrogen sulfide; Cardiovascular; Cystathionine beta-synthase; Cystathionine-gamma-lyase; Gasotransmitter

Many of us experienced foul rotten egg like smell of hydrogen sulfide (H_2S) in chemistry laboratories during school days. Did you know back then that it is actually produced in our body? H_2S is the third and newest member of gasotransmitter family, along with nitric oxide (NO) and carbon monoxide (CO). Since the first report on physiological role of endogenously produced H_2S in mammals [1], there has been an explosion of research articles on its varied biological roles. Availability of grants from American Health Association (AHA), United States Centers for Disease Control and Prevention (CDC), National Institutes of Health (NIH) and other government agencies over two decades enabled deep dive research on the role of H_2S in cardiovascular health and other diseases. For example, H_2S induced vasorelaxation in cardiovascular system via its interaction with K_{ATP} channels [2,3], role of H_2S in protein S-sulfhydration [4,5], redox balance [6,7], and regulation of levels of second messenger like free calcium, cGMP, cAMP [8-10], were discovered. Abnormal endogenous metabolism of H_2S was found to have direct relationship with disorders in cardiovascular, gastrointestinal, nervous system and other diseases [Figure 1; details are reviewed in [11,12]. Even human genetic diseases were linked to H_2S -generating enzymes.

H_2S is a weak acid with pKa1 and pKa2 values 6.9 and >12, respectively. At physiological pH (~7.4), the ratio of HS^- to H_2S is around 3:1. It is unclear which one among hydrosulfide ion (HS^-) or H_2S is the dominant mediator of various physiological effects. Above 20 μM , H_2S inhibits cytochrome c oxidase in intact cells and is known to mess up electron transfer chain. Three major enzymes have been




Figure 2: H_2S -generating reactions catalyzed by CBS, CSE and CAT/MST. Arrows in black are the oldest known reactions in the trans-sulfuration pathway. Arrows in blue represents recently discovered alternative trans-sulfuration reactions catalyzed by CBS and CSE that produce H_2S , whereas arrows in green are reactions catalyzed by CAT followed by MST to produce H_2S .

identified in mammalian cells that catalyze endogenous production of H_2S (Figure 2). These are cystathionine γ -lyase (CSE), cystathionine β -synthase (CBS) and mercaptopyruvate sulfurtransferase (MST). CSE and CBS are PLP dependent and utilize various substrates including cysteine and homocysteine. MST does it from 3-mercaptopyruvate [13-15]. While CBS and CSE are located in cytosol, MST is partly localized in cytosol and partly in mitochondria [16,17]. CSE produces H_2S in peripheral tissues like cardiovascular and liver. CSE knockout mice were shown to have decreased level of H_2S level in serum [18,19]. CBS is mostly responsible for H_2S production in brain and kidney [20]. MST is believed to be responsible for H_2S production in brain and in vascular endothelial tissues [21,22]. Mutations in human CSE gene could cause hereditary cystathioninuria and hypercystathioninaemia [23,24]. Similarly, inborn errors in CBS gene were shown to be associated with human hereditary diseases like hyperhomocysteinaemia and




Figure 1: An overview of physiological effects of H_2S .

*Corresponding author: Sangita Singh, DuPont, DuPont Knowledge Center, Hyderabad, India, E-mail: Sangita.Singh@dupont.com; singh1.sangita@gmail.com

Received July 25, 2015; Accepted August 11, 2015; Published August 18, 2015

Citation: Singh S (2015) Current Perspective on Hydrogen Sulfide: Stinky Outside Precious Inside. Biochem Physiol 4: 175. doi: [10.4172/2168-9652.1000175](https://doi.org/10.4172/2168-9652.1000175)

Copyright: © 2015 Singh S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

homocystinuria. These were directly related to long-term problems like systemic inflammation, cardiovascular complications and damage of organs [25-27]. Based on *in vitro* data, kinetic simulations predicted that under normal condition and assuming equal concentrations of CBS and CSE, CBS is capable of contributing from 25 to 70% of total H₂S production depending on its extent of allosteric activation by S-adenosylmethionine. However, these simulations predicted that total H₂S formation and relative contribution of CBS in H₂S genesis could decrease under hyperhomocysteinemic conditions where CSE produced 74%, while CBS contributed only 26% of total H₂S in cell.

Extensive investigation on H₂S brought to light that lanthionine and homolanthionine were formed as by products during H₂S biogenesis [13-14]. Presence of lanthionine in mammalian brain is well known for long time, but recent studies have identified neuroprotective and anti-inflammatory properties of a related metabolite, called lanthionine ketamine [28,29]. Researchers are interested to explore mammalian LanC1-like proteins, which are homologous to bacterial enzymes responsible for prokaryotic lantibiotic synthesis [30,31]. Although none of these LanC-like proteins (LanCL) have been shown to catalyze lanthionine formation, there is evidence that LanCL1 directly binds and inhibits CBS [32]. Lanthionine and lanthionine ketamine molecules could be biomarkers for pathologies associated with altered metabolism of H₂S in cell [33,34]. The discovery that CBS is the only source of enzymatic production of lanthionine in cells till date could be useful for treatment of metabolic dysfunctions and neurodegenerative diseases by controlling its metabolism.

So much is done, still there is plenty of blue sky in the horizon. Better understanding of physiological role of H₂S and mechanism of metabolism over last decade led to the discovery of new drug candidates / inhibitors that either aimed to improve or to suppress its endogenous production. These molecules have different mode of actions and they are at different stages in their development pipelines or clinical trials [35,36]. So H₂S based therapeutics is yet several years away from reality or commercialization. Progress in H₂S field has been also hampered by the lack of availability of selective inhibitors of various enzymes that contribute to biosynthesis of this neurotransmitter [37]. Compounds like aminooxyacetic acid (AOAA), trifluoroalanine and hydroxylamine (HA) inhibit both CBS and CSE with different IC₅₀ values. There are claims that propargylglycine (PAG), β-cyano-L-alanine (BCA), L-aminoethoxyvinylglycine (AVG) could be selective inhibitors of CSE but the hunt for specific inhibitors of CBS still continues. Companies need to invest more on R&D and resources to discover more and drive some of those key candidates forward. Clear understanding of signaling mechanism will bring new targets and help to develop drug candidates with new mode of actions. Health data compiled from more than 190 countries shows that heart disease remains the number one global cause of death with 17.3 million deaths each year. These numbers are high, so H₂S based therapeutics can be an enormous opportunity for business, if any of the trial candidates become a break through success.

This field also needs attention from fundamental science and technology professionals from diverse background for novel insights. For example nanotechnology or nanomedicine can contribute to the development of organelle specific targeted delivery devices with controlled released properties. It is equally important to be able to accurately measure levels of H₂S in tissues. This area will require synergy between chemistry and biology for development of new sensors, which will glow up in presence of trace amount of H₂S. Several reaction-based fluorescent probes were already developed which offered advantages like high sensitivity, while maintaining selectivity for H₂S over other RSONs and free thiols [38-41]. Only a few were used successfully in live-

animal imaging and more such probes need to be developed [42,43]. In addition to that development of high end imaging techniques, better contrast to clearly understand H₂S biogenesis in various animal models will enable monitoring the effect of drugs on enhancement or inhibition of H₂S synthesis in live cells. The exact ratio of ionic forms of H₂S inside cell is unknown, but based on its two pKa values; it is expected to be a mixture of HS⁻ and H₂S under physiological pH, although they are in equilibrium. In such cases, knowledge about the reaction mechanism of a dye and reasons for its selectivity towards H₂S over other reactants inside cell need to be clearly understood as it will help researchers to come up with even better solutions. Nonetheless, studies by Qian et al. [42] and Matthew et al. [43] are quite promising for researchers as well as clinical studies. Not only H₂S, sensitive tools are required to detect and estimate quantities of other known metabolites like lanthionine and lanthionine ketamine in live-cells. Among other areas omics will play crucial role in future. For example, targeted or untargeted proteomics and metabolomics using advanced mass spectrometry based detection technologies will help to profile proteins or metabolites related to H₂S metabolism that are either up or down regulated under pathological conditions. These advancements will have power to solve the toughest problems in H₂S research and, in turn, transform cardiovascular and other fields of medicine. Solutions to these are at the interfaces of technologies, so collaboration will continue to remain as critical element for successful, high-impact research.

As we all know, taking pills is a reactive action; an alternate way to enjoy good health is through proper diet planning. Methionine cannot be synthesized by our body and must be supplied through diet. Cysteine, which is synthesized by our body, requires a steady supply of dietary sulfur in order to do so [44]. These amino acids are directly or indirectly utilized in catalytic production of H₂S in cell. Recently polysulfides and similar classes of compounds were identified as H₂S donors. Garlic, onion, Chinese chive, shallot and many other vegetables are rich sources of organosulfides like acyclic disulfide, trisulfide, tetrasulfide and cyclic polysulfide [45,46]. Regular dietary supplementations of polysulfides or organosulfides were found to be beneficial for prevention of cardiovascular diseases [47,48]. Along with that improved H₂S signaling helped sustain reduced blood pressure, promote growth of new arterioles, reduce bad cholesterol and triglyceride levels. However it should be kept in mind that excess of H₂S is toxic. So a controlled consumption of rich natural sources of sulfides through good food practices could be an inexpensive and sustainable way to healthy heart and life.

Footnotes

Sangita Singh worked on H₂S biogenesis during her Ph.D. at the University of Nebraska, Lincoln and University of Michigan, Ann Arbor under the tutelage of Professor Ruma Banerjee

References

1. Abe K, Kimura H (1996) The possible role of hydrogen sulfide as an endogenous neuromodulator. *J Neurosci* 16: 1066-1071
2. Zhao W, Zhang J, Lu Y, Wang R (2001) The vasorelaxant effect of H₂S as a novel endogenous gaseous K(ATP) channel opener. *EMBO J* 20: 6008-6016.
3. Yang W, Yang G, Jia X, Wu L, Wang R (2005) Activation of KATP channels by H₂S in rat insulin-secreting cells and the underlying mechanisms. *J Physiol* 569: 519-531.
4. Kimura H (2014) The physiological role of hydrogen sulfide and beyond. *Nitric Oxide* 41: 4-10.
5. Yang G, Zhao K, Ju Y, Mani S, Cao Q, et al. (2013) Hydrogen sulfide protects against cellular senescence via S-sulphydration of Keap1 and activation of Nrf2. *Antioxid Redox Signal* 18: 1906-1919.

6. Yan SK, Chang T, Wang H, Wu L, Wang R, et al. (2006) Effects of hydrogen sulfide on homocysteine-induced oxidative stress in vascular smooth muscle cells. *Biochem Biophys Res Commun* 351: 485-491.
7. Módis K, Asimakopoulou A, Coletta C, Papapetropoulos A, Szabo C (2013) Oxidative stress suppresses the cellular bioenergetic effect of the 3-mercaptopyruvate sulfur transferase / hydrogen sulfide pathway. *Biochem Biophys Res Commun* 433: 401-407.
8. Bucci M, Papapetropoulos A, Vellecco V, Zhou Z, Pyriochou A, et al. (2010) Hydrogen sulfide is an endogenous inhibitor of phosphodiesterase activity. *Arterioscler Thromb Vasc Biol* 30:1998-2004.
9. Coletta C, Papapetropoulos A, Erdelyi K, Olah G, Módis K, et al. (2012) Hydrogen sulfide and nitric oxide are mutually dependent in the regulation of angiogenesis and endothelium-dependent vasorelaxation. *Proc Natl Acad Sci USA* 109: 9161-9166.
10. Wang R (2012) Shared signaling pathways among gasotransmitters. *Proc Natl Acad Sci USA* 109: 8801-8802.
11. Chen Y, Wang R (2012) The message in the air: hydrogen sulfide metabolism in chronic respiratory diseases. *Respir Physiol Neurobiol* 184: 130-138.
12. Wallace JL, Blackler RW, Chan MV, Da Silva GJ, Elsheikh W, et al. (2015) Anti-inflammatory and cytoprotective actions of hydrogen sulfide: translation to therapeutics. *Antioxid Redox Signal* 22: 398-410.
13. Singh S, Padovani D, Leslie RA, Chiku T, Banerjee R (2009) Relative contributions of cystathionine beta-synthase and gamma-cystathionase to H₂S biogenesis via alternative trans-sulfuration reactions. *J Biol Chem* 284: 22457-22466.
14. Chiku T, Padovani D, Zhu W, Singh S, Vitvitsky V, et al. (2009) H₂S biogenesis by human cystathionine gamma-lyase leads to the novel sulfur metabolites lanthionine and homolanthionine and is responsive to the grade of hyperhomocysteinemia. *J Biol Chem* 284: 11601-11612.
15. Yadav PK, Yamada K, Chiku T, Koutmos M, Banerjee R (2013) Structure and kinetic analysis of H₂S production by human mercaptopyruvate sulfurtransferase. *J Biol Chem* 288: 20002-20013.
16. Nagahara N, Ito T, Kitamura H, Nishino T (1998) Tissue and subcellular distribution of mercaptopyruvate sulfurtransferase in the rat: confocal laser fluorescence and immunoelectron microscopic studies combined with biochemical analysis. *Histochem Cell Biol* 110: 243-250.
17. Taniguchi T, Kimura T (1974) Role of 3-mercaptopyruvate sulfur transferase in the formation of the iron-sulfur chromophore of adrenal ferredoxin. *Biochim Biophys Acta* 364: 284-295.
18. Hosoki R, Matsuki N, Kimura H (1997) The possible role of hydrogen sulfide as an endogenous smooth muscle relaxant in synergy with nitric oxide. *Biochem Biophys Res Commun* 237: 527-531.
19. Yang G, Wu L, Jiang B, Yang W, Qi J, et al. (2008) H₂S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 322: 587-590.
20. Kabil O, Vitvitsky V, Xie P, Banerjee R (2011) The quantitative significance of the trans-sulfuration enzymes for H₂S production in murine tissues. *Antioxid Redox Signal* 15: 363-372.
21. Shibuya N, Mikami Y, Kimura Y, Nagahara N, Kimura H (2009) Vascular endothelium expresses 3-mercaptopyruvate sulfur transferase and produces hydrogen sulfide. *J Biochem* 146: 623-626.
22. Shibuya N, Tanaka M, Yoshida M, Ogasawara Y, Togawa T, et al. (2009) 3-Mercaptopyruvate sulfurtransferase produces hydrogen sulfide and bound sulfane sulfur in the brain. *Antioxid Redox Signal* 11: 703-714.
23. Wang J, Hegele RA (2003) Genomic basis of cystathioninuria (MIM 219500) revealed by multiple mutations in cystathionine gamma-lyase (CTH). *Hum Genet* 112: 404-408.
24. Zhu W, Lin A, Banerjee R (2008) Kinetic properties of polymorphic variants and pathogenic mutants in human cystathionine gamma-lyase. *Biochemistry* 47: 6226-6232.
25. Jhee KH, Kruger WD (2005) The role of cystathionine beta-synthase in homocysteine metabolism. *Antioxid Redox Signal* 7: 813-822.
26. Banerjee R, Zou CG (2005) Redox regulation and reaction mechanism of human cystathionine-beta-synthase: a PLP-dependent hemesensor protein. *Arch Biochem Biophys* 433: 144-156.
27. Yamanishi M, Kabil O, Sen S, Banerjee R (2006) Structural insights into pathogenic mutations in heme-dependent cystathionine-beta-synthase. *J Inorg Biochem* 100: 1988-1995.
28. Ricci G, Vecsi L, Nardini M, Arduini A, Storto S, et al. (1989) Detection of 2H-,4-thiazine-,6-dihydro-,5-dicarboxylic acid (lanthionine ketimine) in the bovine brain by a fluorometric assay. *Biochim Biophys Acta* 990: 211-215.
29. Hensley K, Venkova K, Christov A (2010) Emerging biological importance of central nervous system lanthionines. *Biosynthesis and mode of action of antibiotics. Molecules* 15: 5581-5594.
30. Chatterjee C, Paul M, Xie L, van der Donk WA (2005) Biosynthesis and mode of action of antibiotics. *Chem Rev* 105: 633-684.
31. Chung CH, Kurien BT, Mehta P, Mhatre M, Mou S, et al. (2007) Identification of lanthionine synthase C-like protein-1 as a prominent glutathione binding protein expressed in the mammalian central nervous system. *Biochemistry* 46: 3262-3269.
32. Zhong WX, Wang YB, Peng L, Ge XZ, Zhang J, et al. (2012) Lanthionine synthetase C-like protein 1 interacts with and inhibits cystathionine β²-synthase: a target for neuronal antioxidant defense. *J Biol Chem* 287: 34189-34201.
33. Whiteman M, Le Trionnaire S, Chopra M, Fox B, Whatmore J (2011) Emerging role of hydrogen sulfide in health and disease: critical appraisal of biomarkers and pharmacological tools. *Clin Sci (Lond)* 121: 459-488.
34. DeRatt BN, Ralat MA, Kabil O, Chi YY, Banerjee R, et al. (2014) Vitamin B-6 restriction reduces the production of hydrogen sulfide and its biomarkers by the transsulfuration pathway in cultured human hepatoma cells. *J Nutr* 144: 1501-1508.
35. Wallace JL, Wang R (2015) Hydrogen sulfide-based therapeutics: exploiting a unique but ubiquitous gasotransmitter. *Nat Rev Drug Discov* 14: 329-345.
36. Meng G, Wang J, Xiao Y, Bai W, Xie L, et al. (2015) GYY4137 protects against myocardial ischemia and reperfusion injury by attenuating oxidative stress and apoptosis in rats. *J Biomed Res* 29: 203-213.
37. Asimakopoulou A, Panopoulos P, Chasapis CT, Coletta C, Zhou Z, et al. (2013) Selectivity of commonly used pharmacological inhibitors for cystathionine β² synthase (CBS) and cystathionine β² lyase (CSE). *Br J Pharmacol* 169: 922-932.
38. Miller EW, Chang CJ (2007) Fluorescent probes for nitric oxide and hydrogen peroxide in cell signaling. *Curr Opin Chem Biol* 11: 620-625.
39. Lippert AR, New EJ, Chang CJ (2011) Reaction-based fluorescent probes for selective imaging of hydrogen sulfide in living cells. *J Am Chem Soc* 133: 10078-10080.
40. Yu F, Hanab X, Chen L (2014) Fluorescent probes for hydrogen sulfide detection and bioimaging. *Chem Commun* 50: 12234-12249.
41. Shimamoto K, Hanaoka K (2015) Fluorescent probes for hydrogen sulfide (H₂S) and sulfane sulfur and their applications to biological studies. *Nitric Oxide* 46: 72-79.
42. Qian Y, Karpus J, Kabil O, Zhang SY, Zhu HL, et al. (2011) Selective fluorescent probes for live-cell monitoring of sulphide. *Nat Commun* 2: 495.
43. Hammers MD, Taormina MJ, Cerda MM, Montoya LA, Seidenkranz DT, et al. (2015) A Bright Fluorescent Probe for H₂S Enables Analyte-Responsive, 3D Imaging in Live Zebrafish Using Light Sheet Fluorescence Microscopy. *J Am Chem Soc* .
44. Stipanuk MH (1986) Metabolism of sulfur-containing amino acids. *Annual Review of Nutrition* 6: 179-209.
45. Corzo-Martínez M, Corzo N, Villamiel M (2007) Biological properties of onions and garlic. *Trends Food Sci Tech* 18: 609-625.
46. Tocmo R, Liang D, Lin Y, Huang D (2015) Chemical and biochemical mechanisms underlying the cardioprotective roles of dietary organopolysulfides. *Front Nutr* 2: 1.
47. Benavides GA, Squadrito GL, Mills RW, Patel HD, Isbell TS, et al. (2007) Hydrogen sulfide mediates the vasoactivity of garlic. *Proc Natl Acad Sci U S A* 104: 17977-17982.
48. Tsai CW, Chen HW, Sheen LY, Lii CK (2012) Garlic: health benefits and actions. *Biomedicine* 2: 17-29.