

Cuttings as an Alternative for Propagation of *Tillandsia recurvata*: An Epiphytic Species of Ethnobotanic Value

Díaz SV^{1*}, España PC¹, Arango ICP¹, Zapata EV², Aparicio AR² and Palacios AF³

¹Biotechnology Research Center (CEIB), Autonomous University of the State of Morelos, Mexico

²Biotic Products Development Center, National Polytechnic Institute, Mexico

³Research Center on Biodiversity and Conservation, Autonomous University of the State of Morelos, Mexico

Abstract

Tillandsia recurvata has anticancer properties as well as horticultural value. Production through seed germination is non-viable since the species produces low numbers of seeds with low viability, longevity and germination rates. We explored propagation using cuttings and tested different inert substrates and plant growth regulators. In 2008, cuttings were taken from *T. recurvata* plants and sown in/on three different substrates (red lava rock, gravel and electricity cable) to which growth regulating cytokinins [6-Benzilaminopurine (BAP) and Kinetin (KIN)] were added. The effects of substrate and cytokinin addition were determined in terms of cutting survival and four different relative growth indices (length, weight, number of live leaves and number of new leaves). Cutting weight, number of new leaves and survival did not differ among the different substrate types and plant growth regulators. The cuttings sown on cable presented a higher quantity of live leaves. Cutting length was greater in the control or with addition of KIN; length was also greater when the cuttings were sown in red lava rock. The red lava rock and cable are both substrates with potential for propagation of *T. recurvata*; however, given the lack of differences found between the control and KIN treatments, addition of these plant growth regulators appears to be unnecessary. We conclude that sowing cuttings in/on inert substrates constitutes an inexpensive and easily implemented technique for propagation of this species.

Keywords: Epiphytes; *Tillandsia recurvata*; Propagation; Plant growth regulators

Introduction

Many wild plant species are extracted from their environment because they are of ornamental value [1] and/or represent a source of chemical compounds (i.e., natural products with therapeutic properties) [2], which give them a measure of ethnobotanical value [3]. However, chronic extraction of plants can negatively impact populations and have repercussions at both community and ecosystem levels [4]. As a consequence, efforts have been made towards the generation of biotechnological [5] and/or agricultural (e.g. domestication [6]) protocols that permit the sustainable management of populations under extraction. The search for adequate substrates for plant propagation is also important [7], as is examination of the effects of plant growth regulators [8]. These factors can act to optimize the survival and growth of artificially propagated plants.

Members of the family Bromeliaceae have different ethnobotanical uses [9]; species of the genus *Tillandsia* are often used in religious festivals [1,10,11] and are also useful from a horticultural perspective [12], while others have therapeutic properties [9,13,14]. The epiphyte *Tillandsia recurvata* (L.) has multiple uses, since extracts of this species are cytotoxic to some human carcinogenic cellular lines [15], in addition to the allelopathic [16] and horticultural [12] potential of some individuals.

T. recurvata is an important species, since it reincorporates nutrients to the ecosystem [17], interacts with its phorophytes [18] and other epiphytes [16] and participates in multi specific interactions (xylophage-host-epiphyte [19,20]). It is an atmospheric species (i.e., the roots are for anchorage only and nutrients are acquired directly from the environment) [21] of wide distribution and is found from the southern United States to Argentina [22]. It can complete its life cycle on inert substrates such as electricity cables [23] and is highly abundant in some forests [24,25]. However, its fruits contain few seeds (average 50 seeds) [12,26,27] of approximately 2.5 mm in

length with a white plumose appendage [22], which present lower longevity and germination than other sympatric *Tillandsia* species (e.g. *Tillandsia caput-medusae*) [12]. One alternative to propagation by seed germination is the development of propagation technologies that generate information about the factors that act to promote the growth of *Tillandsia recurvata* and allow the generation of individuals with similar characteristics suitable for use as ornamental plants and/or sources of natural products. Apart from one study [28] conducted in wild conditions, there is a lack of information about the growth of atmospheric *Tillandsia* species. In this study, we suggest the use of cuttings taken from *T. recurvata* individuals as a direct and rapid method of propagation. The objective of this study was to evaluate the growth and survival of *T. recurvata* cuttings in different inert substrates and assess the effect of adding different plant growth regulators. We hypothesized that, as an atmospheric species, cuttings of *T. recurvata* would present greater survival and growth in substrates where they are more exposed and that the addition of plant growth regulators would act to increase their survival and growth.

Material and Methods

Collection site and study species

In January 2008, 35 *Tillandsia recurvata* individuals were collected in San Andrés de la Cal, Tepoztlán, Morelos, Mexico (8°57'22.2" W;

*Corresponding author: Díaz SV, Biotechnology Research Center (CEIB), Autonomous University of the State of Morelos, Mexico, Tel: 52-777-329-70-57; E-mail: susana.valencia@uaem.mx

Received June 23, 2017; Accepted July 05, 2017; Published July 15, 2017

Citation: Díaz SV, España PC, Arango ICP, Zapata EV, Aparicio AR, et al. (2017) Cuttings as an Alternative for Propagation of *Tillandsia recurvata*: An Epiphytic Species of Ethnobotanic Value. J Hortic 4: 202. doi: [10.4172/2376-0354.1000202](https://doi.org/10.4172/2376-0354.1000202)

Copyright: © 2017 Díaz SV, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

99°06'50.2" N, 1495 m a.s.l.). The dominant vegetation type at the collection site is tropical deciduous forest (For more information about the collection) [12,20,24].

The epiphytic bromeliads found at the study site include *Tillandsia achyrostachys* E. Morren ex Baker, *Tillandsia caput-medusae* E. Morren, *Tillandsia hubertiana* Matuda, *Tillandsia schiedeana* Steud., *Tillandsia ionantha* Planch., *Tillandsia makoyana* Baker, *Tillandsia recurvata* and *Viridantha atroviridipetala* (Matuda) Espejo. *T. recurvata* accounts for 72% of the epiphytes present [24]. Plants of this species are herbaceous epiphytes, rupicolous or growing on suspended cables, acaulescent, the leaves have a dense covering of trichomes, which gives them a white-grey color, and they are verticillate, forming a tussock of leaves. A tussock is considered to be a group of *T. recurvata* plants that form a spherical (5-10 cm diameter) and discrete cluster, where all of the leaves come from the center of the sphere and delimit the tussock with their tips [29]. In each tussock, there are different branches with a few leaves distributed in a distichous form.

Experimental design

The collected plant material was transported to the Biotechnology Laboratory of the Centro de Desarrollo de Productos Bióticos-Instituto Politécnico Nacional (CeProBi-IPN) and 7-9 cuttings of similar size and shape were obtained from each plant. A cutting is a module of the plant (e.g. branches) separated from its progenitor that can give rise to an independent individual (clone) with genetic material identical to that of the plant from which it was originally obtained [30]. In this study, a *T. recurvata* branch was considered a cutting. Prior to sowing, the identity of each cutting was recorded by labeling them with progressive numbers (1 to 270) and the length (mm) and weight (g) of each was recorded, along with the number of leaves.

The cuttings were sown in flat polystyrene germination trays (28.5 × 55 cm × 6 cm). The trays contained one of the three different substrates (red lava rock, gravel and electricity cable). The red lava rock (trademark PISUMMA, size 1.5 cm) and gravel (Trademark PISUMMA, white color, size small) were poured into the trays until reaching a uniform depth of 4 cm. To install the cable (Model: Indiana 8.37 mm², 8-AWG), two perforations were made at each end of the trays and the cable was attached between each pair of perforations, providing two fixed cables per tray. Ten randomly-assigned cuttings of *T. recurvata* were sown in each tray. In the case of the trays with cables, five cuttings were attached to each cable equidistantly using covered twist-tie wires. The trays were placed in a greenhouse and the position of each tray was assigned randomly. The relative humidity and temperature of the greenhouse was monitored hourly (Onset HoBo ProSeries, model H08-32-IS) over the course of the experiment (January-March of 2008). The monthly average greenhouse temperatures during the experiment were: January (mean ± SD) 20.91°C ± 0.62°C, February 21.41°C ± 8.97°C and March

22.54 ± 9.14°C. Relative humidity was 42.78% ± 21.75% in January, 34.38% ± 17.26% in February and 33.05% ± 0.99% in March.

According to the treatment in each tray, the plants were sprayed with the cytokinins 6-Benzylaminopurine [(BAP) (0.2 mg/L, Sigma B-3408)] or 6-Furfurylaminopurina [(KIN) (0.2 mg/L, Sigma K-0753)] or with distilled water containing no growth regulator (control). These plant growth regulators were chosen because they promote cellular division and break the dormancy of lateral buds [31]. The cytokinin concentrations were low since the plants were being sprayed every third day. At the end of the experiment, the cuttings were removed from the trays and their length and weight recorded, along with the number of new leaves, live leaves and the number of cuttings that remained alive. Based on the measurements of the cuttings before and after sowing, a relative growth index was obtained for each of the variables, using the following formula:

$$\frac{VA - VP}{VP}$$

Where VA is equal to the value of the variable (e.g. weight) after sowing the cutting and VP is the value of the variable prior to sowing. A positive value of the index indicates growth.

Statistical analysis was conducted using an ANOVA with two factors [32], where one factor was the "growth regulator" with three levels: BAP, KIN and No Regulator (NR), and the other were "substrate type", also with three levels: red lava rock, gravel and cable. The combinations of the levels of each factor gave a total of nine treatments (BAP-red lava rock, BAP-gravel, BAP-cable, KIN-red lava rock, KIN-gravel, KIN-cable, NR-red lava rock, NR-gravel, NR-cable). Within each treatment, there were three trays or units (N=27) with 10 cuttings in each tray (N=270). The response variables were the relative growth indices of cutting length, weight, new leaves, live leaves and survival measured as the number of live cuttings remaining in each tray. All analysis was conducted using the program Stata 13 [33].

Results

Prior to sowing, the length of the cuttings of *Tillandsia recurvata* was 48.84 mm ± 14.77 mm, while the weight was 0.31 g ± 0.13 g and the number of leaves was 8.67 ± 2.27. Relative growth rate in weight of the cuttings did not differ among the different substrate types ($F_{2,18}=0.31$, $P=0.73$; cable=0.08 g ± 0.96 g, gravel=-0.03 g ± 0.09 g, red lava rock=-0.12 g ± 0.04 g), neither the growth regulator ($F_{2,18}=0.85$, $P=0.44$; BAP=-0.14 g ± 0.08 g, KIN=-0.10 g ± 0.06 g, NR=0.17 g ± 0.93 g) nor the interaction between substrate and regulator ($F_{2,18}=0.69$, $P=0.60$; Table 1) affected this variable.

Survival of the cuttings (measured as live cuttings per tray) was also unaffected by substrate type ($F_{2,18}=0.92$, $P=0.041$; cable=0.97 ± 0.04, gravel=0.93 ± 0.08, red lava rock=0.96 ± 0.05), growth regulator

Substrate	Plant Growth Regulator	Survival (Proportion of alive cuttings)	Relative Growth Rate (Mean ± SE)		
			Weight (g)	Number of new leaves	Number of live leaves
Red lava rock	KIN	0.97 ± 0.58	-0.12 ± 0.02	0.38 ± 0.14	0.17 ± 0.09
	BAP	0.97 ± 0.58	-0.11 ± 0.03	0.31 ± 0.10	0.21 ± 0.08
	Control	0.97 ± 0.58	-0.14 ± 0.07	0.54 ± 0.19	0.27 ± 0.14
Gravel	KIN	0.93 ± 0.11	-0.05 ± 0.03	0.24 ± 0.22	0.14 ± 0.10
	BAP	0.93 ± 0.05	-0.08 ± 0.08	0.14 ± 0.07	0.20 ± 0.20
	Control	0.93 ± 0.11	0.02 ± 0.14	0.86 ± 0.81	0.23 ± 0.19
Cable	KIN	0.96 ± 0.05	-0.15 ± 0.07	0.13 ± 0.15	0.41 ± 0.27
	BAP	1.00 ± 0.00	-0.24 ± 0.05	0.33 ± 0.06	0.47 ± 0.10

Table 1: Mean ± SE of the variables measured in cuttings of *Tillandsia recurvata*. Values correspond to the interaction of substrate × plant growth regulator.

($F_{2,18}=0.07$, $P=0.93$; $BAP=0.96 \pm 0.05$, $KIN=0.95 \pm 0.07$, $NR=0.95 \pm 0.07$) or by the interaction between substrate and regulator ($F_{2,18}=0.07$, $P=0.98$; Table 1).

Relative growth rate in the new leaves did not differ among the three different substrates ($F_{2,18}=0.18$, $P=0.82$; $cable=0.63 \pm 1.39$, $gravel=0.41 \pm 0.64$, $red\ lava\ rock=0.41 \pm 0.16$), with the addition of the different plant growth regulators ($F_{2,18}=1.82$, $P=0.19$; $BAP=0.26 \pm 0.11$, $KIN=0.25 \pm 0.18$, $NR=0.94 \pm 1.37$) or with the interaction between substrate and regulator ($F_{2,18}=0.35$, $P=0.83$; Table 1).

Substrate type did influence the relative growth rate in live leaves per cutting ($F_{2,18}=3.90$, $P=0.03$), where those sown on the cable (0.39 ± 0.18) presented a higher quantity of live leaves than those sown in gravel (0.19 ± 0.15), while the relative growth rate of those sown in red lava rock was intermediate (0.21 ± 0.10 , Tukey $P<0.05$). Neither growth regulator ($F_{2,18}=0.24$, $P=0.78$; $BAP=0.29 \pm 0.17$, $KIN=0.24 \pm 0.19$, $NR=0.26 \pm 0.16$) nor the interaction between growth regulator and substrate type affected the growth of the live leaves ($F_{2,18}=0.50$, $P=0.72$; Table 1).

Substrate type also affected the growth rate in terms of cutting length ($F_{2,18}=7.21$, $P=0.005$); cuttings sown in the red lava rock were longer ($0.10\ mm \pm 0.06\ mm$) than those sown on cable ($0.005\ mm \pm 0.12\ mm$), while those sown in gravel actually lost length ($-0.002\ mm \pm 0.04$). The addition of growth regulators also affected the final length of the cuttings ($F_{2,18}=4.02$, $P=0.03$); cuttings with KIN ($0.69\ mm \pm 0.08\ mm$) were longer than those with BAP ($-0.01\ mm \pm 0.08\ mm$), while the control did not differ from either cytokinin treatment ($0.05\ mm \pm 0.09\ mm$). Growth regulator and substrate type interacted marginally in terms of cutting length ($F_{2,18}=3.12$, $P=0.04$; Figure 1). The longest cuttings were those sown in red lava rock with no growth regulator ($0.69\ mm \pm 0.08\ mm$), those sown in red lava rock with added KIN ($0.09\ mm \pm 0.06\ mm$) and those sown on cable with added KIN ($0.12\ mm \pm 0.09\ mm$, Figure 1). Cuttings sown on cable with added BAP presented a negative index of length ($-0.09\ mm \pm 0.08\ mm$; Figure 1).

Discussion

Tillandsia recurvata presents a CAM metabolism [34] that is adapted to stressful conditions. Among other attributes (anemochory, autogamy) [35], this metabolism type confers advantages in terms of colonization of the canopy in arid and semi-arid environments [24,25]. As with other *Tillandsia* species [36], *T. recurvata* can establish itself and grow abundantly on inert substrates, such as suspended cables [23]. In this study, we found that the cuttings of *T. recurvata* transplanted onto cable presented a higher number of live leaves than those sown in red lava rock, but an equal number of leaves as those presented by the cuttings sown in gravel. This may be due to the atmospheric habit of this species [21]; i.e., growing on suspended cable implies that the plant is more exposed in terms of water and nutrient capture, while excess water is eliminated to the atmosphere. The water that *T. recurvata* acquires through its leaves enters its tissues, but if excess water is not eliminated, this can interfere with respiration, increasing leaf mortality. On the cable, the cuttings can generate leaves without increasing stem length, but in the two other inert substrate types there were negative costs to the leaves and increases in stem length. Unlike the cable, red lava rock is a porous substrate that can store water in its cavities [7], while the gravel does not accumulate water in its pores, but only between its conglomerate units. It is possible that higher leaf mortality occurs as a result of not eliminating moisture, generating fewer live leaves and causing lower growth of the plant in these substrates. The cuttings in red lava rock had as many live leaves as those on the cable and presented increased length. This growth pattern could be a strategy adopted by the cuttings in order to escape the effects of excess moisture.

There is evidence that when plants of *T. juncea* are small they grow faster [28]. When our experiment began, the cuttings of *T. recurvata* measured 48.84 mm of height, which is similar to the height of *T. brachycaulus* seedlings (<0.4 cm) or infants (0.5-2.4 cm) [37]. Accordingly, we expected that three months would be enough to detect a conspicuous growth in cuttings of *T. recurvata*; however, growth in *T. recurvata* cutting length was lower than in other atmospheric *Tillandsia* species. It has been reported that *T. butzii* and *T. juncea* have an annual




Figure 1: Length of *Tillandsia recurvata* cuttings sown in three different substrates with addition of two different plant growth regulators and a control with no added regulator (■=BAP, ●=KIN, ◆=control). Different letters indicate significant differences (Tukey test, $P<0.05$). Points represent mean values and bars represent 95% confidence intervals.

growth rate of 2.8 cm \pm 0.28 cm and 3.5 cm \pm 0.28 cm respectively [28]. Without considering the effects of environmental conditions throughout the year, this implies that monthly average growth would be 0.23 cm and 0.29 cm for *T. butzii* and for *T. juncea*, respectively. These values are higher than the maximum growth rate we found for *T. recurvata* (0.15 mm equivalent to 0.015 cm). It is possible that this low growth rate of *T. recurvata* is a result of the stressful conditions of dry tropical forest, while *T. butzii* and *T. juncea* inhabit lower montane cloud forests, where water is not a limiting factor.

The cytokinins BAP and KIN are frequently used in biotechnological protocols since they induce the development of buds and increase leaf growth due to foliar expansion as a consequence of cellular division and lengthening [38]. In this study, we found that the cuttings to which KIN were added were longer at the end of the experiment than those to which BAP were added. However, addition of KIN to the cuttings did not produce results that differed significantly from those of the control. This suggests that, in this case, addition of plant growth regulators is probably unnecessary because addition of these cytokinins by irrigation means that they are absorbed via the leaf stomata, a common procedure among atmospheric epiphytes, while when they are administered to explants cultivated *in vitro*, they are absorbed by the roots.

T. recurvata is a very abundant species in some zones [24-26]; however, there is evidence that the abundance of such species of horticultural and ethnobotanical value has been impacted by human extraction [1]. For this reason, it is important to consider alternatives for propagation within conservation plans that select the most desired phenotypic characteristics of the *T. recurvata* plants.

Conclusion

The results of this study demonstrate that red lava rock was the substrate in which the plants maintained the highest number of live leaves and presented greatest growth. With respect to the addition of plant growth regulators, at the end of the experiment, the cutting length was equal with or without the addition of KIN. It can therefore be concluded that the choice of substrate for sowing cuttings of *Tillandsia recurvata* is a more important factor than the addition of plant growth regulators. It is therefore suggested to use red lava rock for the purposes of propagation of plants of this species. Finally, given that the final length of the cuttings did not differ between the KIN and control treatments, the use of plant growth regulators appears to be unnecessary at least in the case of BAP and KIN.

Acknowledgements

We are grateful to T. Cruz-Fernández, A. Martínez-Ray, and S.E. Flores-Valencia for their help with work in the greenhouse. We appreciate the comments of K. Macmillan to the ms. This study formed part of the Doctorate studies of SVD. We thank the Consejo Nacional de Ciencia y Tecnología for the scholarship (87232) awarded to SVD.

References

- Palacios AF, Díaz SV (2007) Local illegal trade reveals unknown diversity and involves a high species richness of wild vascular epiphytes. *Biol Conserv* 136: 372-387.
- Kingston DGI (2010) Modern natural products drug discovery and its relevance to biodiversity conservation. *J Nat Prod* 74: 496-511.
- Urrell JA, de Albuquerque UP (2012) Is ethnobotany an ecological science? Steps towards a complex ethnobotany. *Ethnobiol Conserv* 1: 4-20.
- Ticktin T (2004) The ecological implications of harvesting non-timber forest products. *J App Eco* 41: 11-21.
- Osorio-Rosales ML, Mata-Rosas M (2005) Micropropagation of endangered mexican species of Ponytail palms. *HortScience* 40: 1481-1484.
- Palevitch D (1996) *Agronomy Applied to Medicinal Plants*. In: Akerele O, Heywood V, Synge H (eds.) Conservation of Medicinal Plants. Cambridge University Press: Cambridge pp: 167-178.
- Anicua-Sánchez R (2008) Physical and micromorphological characterization of organic and inorganic materials for the generation of mixtures of substrates in the production of *Lisianthus (Eustoma grandiflorum)*. Montecillo, Edo. From Mexico: College of Postgraduates. Institute of Teaching and Research in Agricultural Sciences.
- Gaspar T, Kevers C, Penel C, Grepping H, Reid DM, et al. (1996) Plant hormones and plant growth regulators in plant tissue culture. *In Vitro Cell Dev Biol-Plant* 32: 272-289.
- Hornug-Leoni CT (2011) Avances sobre usos Etnobotánicos de las Bromeliaceae en Latinoamérica. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* 10: 297-314.
- Haeckel IB (2008) The "Arco Floral": ethnobotany of *Tillandsia* and *Dasyliirion* spp. in a Mexican religious adornment. *Economic Botany* 62: 90-95.
- Toledo-Aceves T, García-Franco JG, López-Barrera F (2014) Bromeliad rain: An opportunity for cloud forest management. *Forest Ecol Manag* 329: 129-136.
- Flores-Palacios A, Bustamante-Molina A, Corona-López AM, Valencia-Díaz S (2015a). Seed number, germination and longevity in wild dry forest *Tillandsia* species of horticultural value. *Scientia Horticulturae* 187: 72-79.
- Vite-Posadas JA, Brechú-Franco AE, Laguna-Hernández G, Rojas-Bribiesca MG, Osuna-Fernández RH (2011) Morphoanatomical characterization and antimicrobial activity of *Tillandsia imperialis* (Bromeliaceae). *Polibotánica* 31: 21-29.
- Lowe HIC, Watson CT, Badal S, Ateh EN, Toyang NJ, et al. (2014) Unearthing the medicinal properties of *Tillandsia recurvata* (Ball Moss): a mini review. *Eur J Med Plants* 4: 1138-1149.
- Lowe HIC, Watson CT, Badal S, Ateh EN, Toyang NJ, et al. (2012) Anti-angiogenic properties of the Jamaican Ball moss (*Tillandsia recurvata* L.). *Inter Res J Bio Sci* 1: 73-76.
- Valencia-Díaz S, Flores-Palacios A, Rodríguez-López V, Jiménez-Aparicio AR (2012) Effects of *Tillandsia recurvata* extracts on the seed germination of *Tillandsia* spp. *Allelopat J* 29: 125-136.
- Flores-Palacios A, Barbosa-Duchateau CL, Valencia-Díaz S, Capistrán-Barradas A, García-Franco JG (2014) Direct and indirect effects of *Tillandsia recurvata* on *Prosopis laevigata* in the Chihuahua desert scrubland of San Luis Potosi, Mexico. *J Arid Environ* 104: 88-95.
- Valencia-Díaz S, Flores-Palacios A, Rodríguez-López V, Ventura-Zapata E, Jiménez-Aparicio AR (2010) Effect of host-bark extracts on seed germination in *Tillandsia recurvata*, an epiphytic bromeliad. *J Trop Ecol* 26: 571-581.
- Valencia-Díaz S, Corona-López AM, Toledo-Hernández VH, Flores-Palacios A (2014) Is branch damage by xylophages related to the presence of epiphytes? *Arthropod-Plant Interactions* 8: 25-32.
- Flores-Palacios A, Corona-López AM, Ríos MY, Aguilar-Guadarrama B, Toledo-Hernández VH, et al. (2015b) Does allelopathic activity of *Ipomoea murucoides* induced by xilophage damage? *Plos ONE* 10: e0143529.
- Benzing DH, Seeman J, Renfrow A (1978) The foliar epidermis in Tillandsioideae (Bromeliaceae) and its role in habitat selection. *Am J Bot* 65: 359-365.
- Espejo-Serna A, López-Ferrari AR, Ramírez-Morillo I (2005) Bromeliaceae. *Flora de Veracruz*. Fascículo 136. Instituto de Ecología A.C. Xalapa, Veracruz y University of California, Riverside CA. México.
- Puente ME, Basahn Y (1994) The desert epiphyte *Tillandsia recurvata* harbours the nitrogen-fixing bacterium *Pseudomonas stutzeri*. *Can J Bot* 72: 406-409.
- Vergara-Torres CA, Pacheco-Álvarez MC, Flores-Palacios A (2010) Host preference and host limitation of vascular epiphytes in a tropical dry forest of central Mexico. *J Trop Ecol* 26: 563-570.
- Flores-Palacios A, García-Franco JG, Capistrán-Barradas A (2015) Biomass, phorophyte specificity and distribution of *Tillandsia recurvata* (L.) L. in a tropical semi-desert environment (Chihuahuan Desert, Mexico). *Plant Ecol Evol* 148: 68-75.
- Caldiz DO, Fernández LV (1995) The role of the epiphytic weeds *Tillandsia recurvata* and *T. aëranthos* in native rural and urban forest. *Int J Ecol Environ Sci* 21: 177-197.

27. Bernal R, Valverde T, Hernández-Rosas L (2005) Habitat preference of the epiphyte *Tillandsia recurvata* (Bromeliaceae) in a semi-desert environment in Central Mexico. Can J Bot 83: 1238-1247.
28. Hietz P, Ausserer J, Schindler G (2002) Growth, maturation and survival of epiphytic bromeliads in a Mexican humid montane forest. J Trop Ecol 18: 177-191.
29. Flores-Palacios A (2016) Does structural parasitism by epiphytes exist? A case study between *Tillandsia recurvata* and *Parkinsonia praecox*. Plant Biol 18: 463-470.
30. DeVere Burton L (2015) Agriscience: fundamentals and applications. Delmar Thomson Learning, USA p: 841.
31. Hopkins WG (2007) Plant biotechnology. Chelsea House Publishers, NY, USA.
32. Zar JH (2010) Biostatistical analysis (5th edn.). Prentice Hall, NJ, USA.
33. Statacorp (2013) Stata statistical software: release 13. StataCorp LP, College Station, TX, USA.
34. Matiz A, Miotto PT, Yepes A, Freschi L, Mercier H (2013) CAM photosynthesis in bromeliads and agaves: what can we learn from these plants? In: Dubinsky Z (ed.) Photosynthesis. InTech, Rijeka, pp: 91-134.
35. Orozco-Ibarrola OA, Flores-Hernández PS, Victoriano-Romero E, Corona-López AM, Flores-Palacios A (2015) Are breeding system and florivory associated with the abundance of *Tillandsia* species (Bromeliaceae)? Bot J Linnean Soc 177: 50-65.
36. Wester S, Zott G (2010) Growth and survival of *Tillandsia flexulosa* on electrical cables in Panama. J Trop Ecol 26: 123-126.
37. Mondragón D, Durán R, Ramírez I, Valverde T (2004) Temporal variation in the demography of the clonal epiphyte *Tillandsia brachycaulos* (Bromeliaceae) in the Yucatán Peninsula, Mexico. J Trop Ecol 20: 189-200.
38. George EF, Hall MA, Jan de Klerk G (2008) Plant propagation by tissue culture (Vol. 1). Springer p: 479.

Citation: Díaz SV, España PC, Arango ICP, Zapata EV, Aparicio AR, et al. (2017) Cuttings as an Alternative for Propagation of *Tillandsia recurvata*: An Epiphytic Species of Ethnobotanic Value. J Hort 4: 202. doi: [10.4172/2376-0354.1000202](https://doi.org/10.4172/2376-0354.1000202)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>