

Cytotoxic and Anti-inflammatory Activities of *Bursera* species from MexicoMaddiel Acevedo<sup>1</sup>, Pablo Nuñez<sup>1</sup>, Leticia González-Maya<sup>2</sup>, Alexandre Cardoso Taketa<sup>1</sup> and María Luisa Villarreal<sup>1\*</sup><sup>1</sup>Centro de Investigación en Biotecnología, Universidad Autónoma del Estado de Morelos, Av. Universidad 1001, Col. Chamilpa, Cuernavaca Morelos, México<sup>2</sup>Facultad de Farmacia, Universidad Autónoma del Estado de Morelos, Av. Universidad 1001, Col. Chamilpa, Cuernavaca, México**\*Corresponding author:** María Luisa Villarreal, Centro de Investigación en Biotecnología, Universidad Autónoma del Estado de Morelos, Av. Universidad 1001, Col. Chamilpa, Cuernavaca Morelos, México, Tel: +51 777 3297057; Fax: 51 777 3297030; E-mail: [luisav@uaem.mx](mailto:luisav@uaem.mx)**Received date:** Nov 13, 2014; **Accepted date:** Feb 19, 2015; **Published date:** Feb 21, 2015**Copyright:** © 2015, Acevedo M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.**Abstract**

**Objective:** Organic extracts from nine species (eleven populations) of the *Bursera* genus native to Mexico, were investigated for cytotoxic and anti-inflammatory activities. The influence of the cytotoxic plant extracts on cell cycle progression and apoptosis were also analyzed.

**Materials and methods:** Cytotoxic activity from *B. ariensis* (Kunth) McVaugh and Rzed. Kew Bull (two populations), *B. bicolor* Engl., *B. lancifolia* Engl., *B. glabrifolia* Engl. (two populations), *B. fagaroides* (La Llave) Rez., Calderón and Medina, *B. linanoe* (La Llave) Rez., Calderón and Medina, *B. galeottiana* Engl., *B. kerberi* Engl., and *B. excelsa* Engl. were evaluated by the sulforhodamine B protein staining assay against four carcinoma cell lines: KB (nasopharyngeal), HF-6 (colon), MCF-7 (breast), and PC-3 (prostate), as well as on HFS-30 fibroblast normal skin cell line. Chloroform extracts from *B. ariensis* (two populations), *B. kerberi* and *B. galeottiana* were added to PC-3 cells to analyze percentage of cells in G1/S and G2/M phases by flow cytometric analysis, while apoptosis was determined in a fluorescence microscope. *In vivo* anti-inflammatory experiments were conducted through the 12-O-tetradecanoylphorbol-13-acetate (TPA) induced ear edema in mice.

**Results:** With the exception of *B. glabrifolia* and *B. excelsa*, hexane extracts of all the collected plant species displayed selective cytotoxic activity in at least one cell line. For the chloroform extracts, only *B. bicolor*, *B. kerberi*, *B. galeottiana*, and *B. fagaroides* exhibited cytotoxic activity, while for the ethyl acetate extracts *B. fagaroides* was the only plant displaying cytotoxic effect. There was no cytotoxic activity detected with methanol extracts. While *B. galeottiana* showed an important effect in G2/M arrest, induced apoptosis in PC-3 cells was detected with *B. ariensis* (two populations) and *B. galeottiana*. Important anti-inflammatory activity similar to the control indomethacin was displayed by *B. galeottiana*, *B. excelsa*, and *B. schlechtendalii*.

**Conclusion:** Findings demonstrated that the studied species possess pharmacological potential qualities for developing plant medicines.

**Keywords:** Cytotoxic; *Bursera*; Anti-inflammatory; Cell cycle, Apoptosis; Ethnomedical use

**Introduction**

From the genus *Bursera*, approximately 80 of the 84 species growing in Mexico are endemic, and are one of the most abundant plant groups of the tropical dry forest [1]. In traditional Mexican medicine several *burseras* are used in the treatment of cancer, although the term “cancer” includes general conditions consistent with cancer symptomatology, i.e., inflammations, ulcers; and such dermatological conditions as hard swellings, abscesses, calluses, corns, warts and polyps [2]. Since pre-hispanic times “copal”, the bark and resin from *Bursera* trees, has been employed in the treatment of cancer symptoms, as well as against some other conditions [3-5]. Some examples are the following: *B. excelsa* known as *pomo* or *tecomahaca* [6] is used to treat tumors, and muscle spasms [7]; *B. glabrifolia* known as *zomplante* is used to treat fever, inflammation and weakness in Oaxaca [8]; *B. galeottiana* known as *cuajote colorado* is used as an analgesic, healing herb, and to treat abscesses [9]; *B. linanoe* known as *copal* is used in the treatment of gums inflammation [10]; the water extract of *B. fagaroides* known as *copalillo* is believed to have an


anticancer effect [11]; *B. bicolor* known as *ticumaca* is used as a medicine to treat inflammation of the muscles [12,13]; *B. ariensis* known as *copal blanco* is used to treat cold and a disease called ‘sacar el frío’, which is a term related to inflammation [14], and *B. schlechtendalii* is used in the treatment of flu [15].

Results of previously conducted experiments on other *bursera* species demonstrated various biological activities. Against cancer, some of them have displayed cytotoxic action; for example, *B. bipinata*, *B. copallifera* and *B. fagaroides* were toxic against several human cancer cell lines that included nasopharyngeal, colon, breast and prostate carcinomas, while *B. graveolens* and *B. grandifolia* were active against fibrosarcoma and epidermoid carcinoma, respectively [2,16]. Other studies with *B. microphylla*, *B. fagaroides*, *B. schlechtendalii*, *B. klugii*, and *B. morelensis* demonstrated antitumor actions against tumors transplanted in mice [11,17-20].

In the present investigation we studied nine Mexican species of the *Bursera* genus which, as mentioned above, are currently in popular use to treat patients with cancer symptomatology. Using traditional and still currently applied ethnomedical information, we conducted experimental analyses to study cytotoxic and anti-inflammatory activities that tend to validate their popular uses. We also determined

the influence of the cytotoxic plant extracts on cell cycle progression and apoptosis.

To our knowledge, the eleven selected populations belonging to nine species have no previously documented investigations, and this new information opens up the possibility of advancing medicinal approaches from these natural Mexican autochthonous resources.



**Figure 1:** Percentage of PC-3 cells in G0/G1 and G2/M after treatments with the chloroform extracts of *B. ariensis* from the state of Mexico (c) *B. ariensis* from Morelos (d), *B. kerberi* (e), and *B. galeottiana* (f); in comparison with a negative control (vehicle, a) and a positive control (PTOX, b).

## Materials and Methods

### Plant material

Nine Mexican species (eleven plant populations) of the genus *Bursera* were included in this study. Seven populations were collected

in the central part of the Mexican Republic. The species *B. ariensis* (Kunth) McVaugh and Rzed. Kew Bull., *B. bicolor* Engl., *B. glabrifolia* Engl., *B. fagaroides* (La Llave) Rez., Calderón and Medina, and *B. linanoe* (La Llave) Rez., Calderón and Medina were collected in the state of Morelos, while *B. kerberi* Engl. and *B. ariensis* (Kunth) McVaugh and Rzed. Kew Bull. were collected in the state of Mexico. Four populations were collected in the southeast of Mexico: *B. excelsa* Engl. was collected in the state of Chiapas; from Oaxaca *B. glabrifolia* Engl. and *B. galeottiana* Engl. were collected, and *B. schlehtendalii* was collected in Puebla. It is important to point out that for *B. glabrifolia* two plant populations were collected, one in Morelos, and the other in the southeastern state of Oaxaca where it was noticed on an unplanned trip; as also happen in the case of a second collection for *B. ariensis*, with one collected in Morelos while the other was fortuitously found on an unplanned trip to Zacazonapan in the state of Mexico. All plant material was collected between September and November 2010. Voucher specimens have been deposited at the Herbarium of the Centro de Investigación en Biodiversidad y Conservación (CIByC), UAEM and were authenticated by Dr. J. R. Bonilla-Barbosa (Table 1).

### Extraction

All bark of the plant material was air-dried and then pulverized with a grinder. The powdered plant material (10 g) was extracted with 100 mL of n-hexane (3x) in a sonicator for 30 min, and filtered at room temperature. To obtain three other extracts, the same procedure was applied but using first chloroform, then ethyl acetate followed by methanol. All of the extracts were then concentrated under reduced pressure at 40°C and stored at 4°C for later use.

### Reagents

RNase, Propidium iodide (PI) and Acridine orange (AO) were obtained from Molecular Probes, Life Technologies, Sigma-Aldrich, MO, USA. RPMI-1640, Fetal bovine serum (FBS), Trypsin- EDTA and NaHCO<sub>3</sub> were obtained from In vitrogen-GIBCO, Carlsbad, CA, USA.

Botanical name	Local name	Coordinates of collection	Site of collection	Voucher specimen no.
<i>B. ariensis</i>	Copal blanco	18°26'00"N 9°02'00"O	Cd. Ayala, elos	27198
<i>B. bicolor</i>	Ticumaca	18°26'00"N 99°02'00"O	Cd. Ayala, Morelos	27196
<i>B. glabrifolia</i>	Copal	18°26'00"N 99°02'00"O	Cd. Ayala, Morelos	27200
<i>B. fagaroides</i>	Copalillo	18°26'00"N 99°02'00"O	Cd. Ayala, Morelos	27199
<i>B. linanoe</i>	Linaloe, Copal	18°28'00"N 99°06'00"O	Tlaquitenango, Morelos	27197
<i>B. glabrifolia</i>	Jiote blanco	16°55'00"N 96°24'00"O	Mitla, Oaxaca	27206
<i>B. galeottiana</i>	Cuajilote colorado	16°55'00"N 96°24'00"O	Mitla, Oaxaca	27195
<i>B. kerberi</i>	Copal	19°05'00"N 100°15'00"O	Zacazonapan, México	27205
<i>B. ariensis</i>	Copalillo	19°05'00"N 100°15'00"O	Zacazonapan, México	27204
<i>B. excelsa</i>	Pomo, Tecamahaca	19°05'00"N 100°15'00"O	Tuxtla Gutierrez, Chiapas	27203
<i>B. schlehtendalii</i>	Copal	18°45'00"N 100°15'00"O	Teotlaco, Puebla	13297

**Table 1:** Collection data of the investigated Mexican *Bursera* species.

### Cytotoxic assay

Cytotoxicity of plant extracts was measured by the sulforhodamine B (SRB) (MP Biomedicals, LLC) protein staining assay [21] using KB (nasopharyngeal), HF-6 (colon), MCF-7 (breast), and PC-3 (prostate) cancer cell lines from ATCC, along with a normal skin fibroblast cell line (HFS-30), passage number 33. Cell cultures were maintained in RPMI-1640 medium (Sigma) supplemented with 10% fetal bovine serum (FBS), 5000 units/mL penicillin, 5 mg/mL streptomycin, 7.5% NaHCO<sub>3</sub>, and cultured in a 96-well microtiter plate (10<sup>4</sup> cells/mL, 190 µL/well) at 37°C in a 5% CO<sub>2</sub>-air atmosphere (100% humidity). The cells at the log phase of growth were treated in triplicate with various concentrations of the test extracts (0.16, 0.8, 4, and 20 µg/mL) that were dissolved in RPMI medium supplemented with 0.025% DMSO, and incubated for 72 h in the conditions described above. The cell concentration was determined by the National Cancer Institute (NCI)

sulforhodamine method [22]. The optical density was measured at 590 nm with an ELISA-Reader (Molecular Devices, SPECTRA max plus 384). Results were expressed at the concentration that inhibits 50% of control growth after the incubation period (IC<sub>50</sub>). The values were estimated from a semilog plot of the extract concentration (µg/mL) against the percentage of viable cells. Podophyllotoxin (PTOX) (Sigma) and camptothecin (Sigma) were included as positive standards. Extracts with IC<sub>50</sub> ≤ 20 µg/mL were considered active according to the NCI guidelines described in the literature [23].

To determine the specificity of cytotoxic activity, the extracts, were assayed against the normal skin fibroblast cell line (HFS-30), and the selective index (SI) was determined using equation (1):

$$SI = \frac{IC_{50} \text{ normal cell line}}{IC_{50} \text{ cancerous cell line}}$$

Extract	Growth inhibition (IC <sub>50</sub> , µg/mL)*				
	Cell line				
	HF6	MCF7	KB	PC3	HFS-30
<i>B. ariensis</i> Morelos	6.34 ± 0.02	14.22 ± 0.01	>20	>20	4.33 ± 0.02
<i>B. bicolor</i>	7.29 ± 0.05	13.76 ± 0.03	>20	9.69 ± 0.01	>20
<i>B. linanoe</i>	>20	>20	13.31 ± 0.05	>20	>20
<i>B. ariensis</i> México	1.65 ± 0.03	2.49 ± 0.05	4.38 ± 0.03	>20	1.27 ± 0.03
<i>B. glabrifolia</i> Morelos	>20	>20	>20	>20	>20
<i>B. kerberi</i>	6.97 ± 0.03	10.29 ± 0.03	>20	>20	>20
<i>B. glabrifolia</i> Oaxaca	6.96 ± 0.02	7.82 ± 0.01	>20	>20	>20
<i>B. galeottiana</i>	4.67 ± 0.01	4.64 ± 0.02	14.26 ± 0.02	5.94 ± 0.04	9.85 ± 0.01
<i>B. fagaroides</i>	1.61 ± 0.04	2.25 ± 0.01	6.29 ± 0.03	2.13 ± 0.01	0.11 ± 0.03
<i>B. excelsa</i>	>20	>20	>20	>20	>20
<i>B. schlechtendalii</i>	4.58 ± 0.01	9.94 ± 0.02	2.13 ± 0.01	0.39 ± 0.03	7.86 ± 0.01
CTP §	0.0258 ± 0.02	0.0424 ± 0.01	0.7855 ± 0.03	0.0383 ± 0.02	0.5821 ± 0.02
PTOX §	5.3127E-04 ± 0.01	3.7453E-03 ± 0.02	4.3833E-04 ± 0.01	1.5584E-03 ± 0.01	2.82E-02 ± 0.03

\*IC<sub>50</sub> is defined as the concentration that resulted in a 50% decrease in cell number, and the results are means ± standard deviations of three independent replicates. Values greater than 20 µg/mL are considered to be non-cytotoxic. § Positive control substances. The cytotoxic effect was investigated on the human cancer cell lines: nasopharyngeal (KB), colon (HF-6), breast (MCF-7), and prostate (PC-3), as well as on a normal fibroblast cell line (HFS-30)

**Table 2:** Cytotoxicity of hexane extracts of selected *Bursera* species from Mexico.

### Cell cycle analysis

PC-3 cells (1x10<sup>5</sup>) were plated in 6-well plates and allowed to attach overnight at 37°C in 5% CO<sub>2</sub>. Exponentially growing cells were treated for 72 h with a concentration of the extracts calculated in accordance to IC<sub>50</sub> values. Cells from each treatment were trypsinized and collected into single cell suspensions, centrifuged and fixed in cold ethanol (70%) overnight at -20°C. The cells were then treated with RNase (0.01M) and stained with PI (7.5 µg/mL) for 30 min in the dark. PI has the ability to bind to RNA molecules and hence, RNase enzyme was added in order to allow PI to bind directly to DNA. The percentage of cells in G1, S and G2 phases was analyzed with a flow cytometer (Becton, Dickinson, FACS Calibur, San Jose, CA), the

number of cells analyzed for each sample was 10,000. Data obtained from the flow cytometer were analyzed using the FlowJo Software (Tree Star, Inc., Ashland, OR, USA) to generate DNA content frequency histograms, and to quantify the number of cells in the individual cell cycle phases.

### Apoptosis analyses

PC-3 cells were seeded in 6-well plates (7.5 x 10<sup>4</sup> cells/mL) and incubated for 18 h. Exponentially growing cells were treated for 72 h with RPMI medium added with 0.025% of DMSO (control) or in accordance with CI<sub>50</sub> of the extracts. Cells treated with H<sub>2</sub>O<sub>2</sub> [24] were used as apoptotic death control, and for the necrotic control PC-3 cells

were treated with boiling water. Then each cell culture was washed with 100 mL of PBS and stained with 100 µL AO/EB solution (100 µg/mL-1 AO, 100 µg/mL-1 EB), according to the procedures of Yang et al. [25]. The cells were observed using a fluorescence microscope (Olympus Co, Tokyo, Japan, with emission at 521 nm). AO/EB are intercalating nucleic acid-specific fluorochromes, which emit green and orange fluorescence respectively when they are bound to DNA. It is well known that AO can pass through cell membranes, but EB cannot. Under the fluorescence microscope, living cells appear green. Necrotic cells stain red but have a nuclear morphology resembling that of viable cells. Apoptotic cells appear green, and morphological changes such as cell blebbing and formation of apoptotic bodies are observed. The criteria for identification are as follows: (i) viable cells appear to have green nucleus with intact structure; (ii) early apoptosis cells exhibit a bright-green nucleus showing condensation of chromatin; (iii) late apoptosis appear as dense orange areas of chromatin condensation, and (iv) Orange intact nucleus depict secondary necrosis [13].

### Anti-inflammatory activity

In vivo anti-inflammatory experiments were conducted in mice according to the Mexican Guidelines for Animal Welfare NOM-Z00-062-1999 (2013) and were approved by the Ethics and Security

Committee from the Centro de Investigación en Biotecnología, UAEM.

Male BALB/c mice weighting between 20 and 25 g were used. Mice were maintained in adequate living conditions at a temperature of 24°C, 70% of humidity with a 12-h light/dark cycle and water/food *ad libitum*. Animal inflammation was induced using 12-O-tetradecanoylphorbol-13-acetate (TPA from Sigma) by the method previously described by Rao et al. [26]. Under general anesthesia by sodium pentobarbital (3.5 mg/kg, intraperitoneally), 10 µL of TPA in ethanol (0.25 mg/mL) were topically applied on the internal and external surface of the right ear to cause edema. After 10 min of incubation, samples at different doses (from 0.05-1 mg/ear) in ethanol (vehicle) were applied on the same ear. The left ear (negative control) received only 10 µL of ethanol, and 20 µL of vehicle. Indomethacin was used as the anti-inflammatory positive control. After 4 h of incubation with TPA, the animals were sacrificed by cervical dislocation. Circular sections of 7 mm were taken from both ears as samples for further measurement. Increase in the weight of the right ear in respect to the left, was considered edema. The inhibition of edema was calculated by the formula:

$$\% \text{ of inhibition} = [(C-E) / C] \times 100$$

Extract	Growth inhibition (IC <sub>50</sub> , µg/mL) <sup>*</sup>				
	Cell line				
	HF6	MCF7	KB	PC3	HFS-30
<i>B. ariensis</i> Morelos	>20	>20	>20	6.78 ± 0.04	>20
<i>B. bicolor</i>	3.90 ± 0.02	11.51 ± 0.03	>20	0.13 ± 0.03	>20
<i>B. linanoe</i>	>20	>20	>20	>20	>20
<i>B. ariensis</i> México	>20	>20	>20	1.87 ± 0.02	>20
<i>B. glabrifolia</i> Morelos	>20	>20	>20	>20	>20
<i>B. kerberi</i>	0.53 ± 0.01	1.74 ± 0.03	2.93 ± 0.03	0.49 ± 0.05	>20
<i>B. glabrifolia</i> Oaxaca	>20	>20	>20	>20	>20
<i>B. galeottiana</i>	3.14 ± 0.04	7.25 ± 0.02	13.14 ± 0.05	2.26 ± 0.04	>20
<i>B. fagaroides</i>	1.46 ± 0.01	3.71 ± 0.01	0.96 ± 0.03	1.62 ± 0.02	2.74 ± 0.04
<i>B. excelsa</i>	>20	>20	>20	>20	>20
<i>B. schlechtendalii</i>	>20	>20	6.52	>20	>20
CTP §	0.0258 ± 0.02	0.0424 ± 0.03	0.7855 ± 0.01	0.0383 ± 0.01	0.5821 ± 0.01
PTOX §	5.3127E-04 ± 0.01	3.7453E-03 ± 0.02	4.3833E-04 ± 0.02	1.5584E-03 ± 0.01	2.82E-02 ± 0.02

<sup>\*</sup>IC<sub>50</sub> is defined as the concentration that resulted in a 50% decrease in cell number, and the results are means ± standard deviation of three independent replicates. Values greater than 20 µg/mL are considered to be non-cytotoxic. § Positive control substances. The cytotoxic effect was investigated on the human cancer cell lines: nasopharyngeal (KB), colon (HF-6), breast (MCF-7), and prostate (PC-3), as well as on a normal fibroblast cell line (HFS-30).

**Table 3:** Cytotoxicity of chloroform extracts of selected *Bursera* species from Mexico.

Where C=edema of control group (treated with TPA), and E=edema of experimental group (TPA with compound). Each crude extract was assayed in triplicate as well as the positive control indomethacin. Statistical analysis for anti-inflammatory activity was

performed using Prism 6 for Windows version 6.0 (GraphPad Software Inc., La Jolla, CA, USA). Using one-way analysis of variance (ANOVA) followed by Dunnett's multiple comparison test, p=0.05 was considered to be statistically significant. ED50 values were

obtained from regression curves with coefficient factors between  $R^2 = 0.80$  and  $0.98$ .


The determination of  $ED_{50}$  for the more active extracts was obtained through a curve constructed with four concentrations (0.05, 0.1, 0.5 and 1 mg/ear) and five replicates for each concentration.

## Results

### Cytotoxicity of extracts

Cytotoxic activities of organic extracts obtained from the nine plant species (eleven populations) are shown in Tables 2 and 3. Results in Table 2 show the cytotoxic values of hexane extracts and reveals that, except for *B. glabrifolia* (Morelos) and *B. excelsa*, all the plants exhibited cytotoxic activity ( $IC_{50} < 20 \mu\text{g/mL}$ ) in at least one cancer cell line, with values ranging from 0.39 to  $14.26 \mu\text{g/mL}$ . From the cancer cell lines used to carry out the evaluation, HF-6 and MCF-7 were the most sensitive ones, showing a 72% response for the tested extracts. The least responsive cancer cell line was PC-3, since only four extracts (*B. bicolor*, *B. galeottiana*, *B. schlechtendalii* and *B. fagaroides*) showed cytotoxic action against this carcinoma. *B. linanoe* showed a selective cytotoxic effect only toward KB carcinoma ( $IC_{50} = 13.31 \mu\text{g/mL}$ ). Both *B. kerberi* and *B. glabrifolia* from Oaxaca, showed cytotoxic action against HF-6 ( $IC_{50} = 6.97 \mu\text{g/mL}$ ), and also toward MCF-7 ( $IC_{50} = 10.29$  and  $7.82 \mu\text{g/mL}$ , respectively) being selective against these two cancer cell lines. *B. fagaroides* was the most active plant against HF-6 ( $IC_{50} = 1.61 \mu\text{g/mL}$ ) and MCF-7 ( $IC_{50} = 2.25 \mu\text{g/mL}$ ) carcinomas and *B. schlechtendalii* was the one most active against KB ( $IC_{50} = 2.13 \mu\text{g/mL}$ ) and PC-3 ( $IC_{50} = 0.39 \mu\text{g/mL}$ ), although it also exhibited activity against the other two cancer cell lines, as well as toward normal fibroblasts. Even though hexane extracts of *B. bicolor*, *B. lancifolia*, *B. linanoe*, *B. kerberi* and *B. glabrifolia* from Oaxaca exhibited toxic effects in at least one cancer cell line, they did not show cytotoxicity against the normal human skin fibroblast cell line HFS-30. On the contrary, *B. ariensis* from Morelos and México, *B. galeottiana*, *B. schlechtendalii* and *B. fagaroides* showed toxic action against normal fibroblasts.


Results in Table 3 show the cytotoxic evaluation of the chloroform extracts from the studied species revealing that *B. bicolor*, *B. kerberi*, *B. galeottiana*, *B. schlechtendalii*, *B. fagaroides*, and *B. ariensis* extracts possess cytotoxic activity ( $IC_{50} < 20 \mu\text{g/mL}$ ) with values ranging from 0.13 to  $13.14 \mu\text{g/mL}$ . *B. fagaroides* showed a generalized cytotoxic action against all carcinomas and also toward normal fibroblasts. *B. schlechtendalii* showed cytotoxic action only against KB ( $IC_{50} = 6.52 \mu\text{g/mL}$ ) and no action against normal fibroblasts. *B. bicolor* was the most active one against PC-3 ( $IC_{50} = 0.13 \mu\text{g/mL}$ ), and presented an outstanding SI (Selective Index) ( $SI = 153.85$ ), also showing cytotoxicity toward the other two carcinomas (HF-6 and MCF-7), but with no action against normal fibroblasts. Similarly *B. kerberi* and *B. galeottiana* extracts were active against the four carcinomas, with no effects against normal fibroblasts; *B. kerberi* presented an important SI for HF-6 and PC-3 cell lines (37.74 and 40.82, respectively). For the ethyl acetate extracts, results of the cytotoxic evaluation revealed that only *B. fagaroides* was active, and this activity was exerted against the four carcinomas HF-6 ( $IC_{50} = 3.46$ ), MCF-7 ( $IC_{50} = 4.06$ ), KB ( $IC_{50} = 2.67$ ) and PC-3 ( $IC_{50} = 4.04 \mu\text{g/mL}$ ). Notably, the methanolic extracts obtained from all the studied species did not exhibit cytotoxicity in any of the studied carcinoma cell lines (data not shown).



**Figure 2:** Phase and fluorescent micrographs of acridine orange and propidium iodide double-stained PC-3 cells. Cells were treated at  $IC_{50}$  values of extracts from *B. ariensis* from the state of Mexico and Morelos (e and f respectively), *B. galeottiana* (g) and *B. kerberi* (h) in a time-dependent manner. Cells were cultured in RPMI 1640 media maintained at  $37^{\circ}\text{C}$  and 5%  $\text{CO}_2$ . Untreated cells after 72 h showed normal structures without prominent apoptosis and necrosis (a). Blebbing and orange color representing the hallmark of late apoptosis were noticed by g and h, early apoptosis features were represented by intercalated acridine orange (bright green) amongst the fragmented DNA (b, c and e-h), and red bright colored necrosis was observed by  $\text{H}_2\text{O}_2$  treatment (d). Images are representatives of one of three similar experiments. Fluorescence and phase contrast images were captured at the field. Fluorescence images of samples were captured at 10x and 40x magnification.

### Cell cycle arrest

To assess the growth inhibitory effect on normal cell cycle progression on the basis of cytotoxic assay results and availability of samples, we conducted a cell cycle analysis measuring intracellular DNA content through flow cytometry in *B. ariensis* from Mexico and Morelos, *B. galeottiana* and *B. kerberi*. The status of the cell cycle of PC-3 cells treated with the four chloroform extracts for 72 h was analyzed, and PTOX was included as a positive control. As shown in Figure 1, exposure of PC-3 cells to chloroform extracts caused the appearance of a population in the sub-G1 region of the profile where apoptotic cells are found. Importantly, the treatment of PC-3 cells with extracts of *B. galeottiana* (Figure 1e) increased the G2/M populations, and correspondingly decreased G0/G1 phase in relation to the negative control (Figure 1a); whereas the treatment with *B. ariensis* from Morelos and from the state of Mexico (Figures 1c and 1d) presented similar apoptotic cells in the sub G1 phase to those of the positive control PTOX (Figure 1b). Only *B. kerberi* did not show a clear change in any phase of the cell cycle (Figure 1f). The above results necessitated performing apoptosis assay



**Figure 3:** Anti-inflammatory activities of selected species of the *Bursera* genus. 1. Indomethacin, 2. *B. ariensis* from Morelos, 3 *B. bicolor*, 4. *B. linanoë*, 5. *B. ariensis* from Mexico state, 6. *B. glabrifolia* from Morelos, 7. *B. kerberi*, 8. *B. glabrifolia* from Oaxaca, 9. *B. galeottiana*, 10. *B. fagaroides*, 11. *B. excelsa*, 12. *B. schlechtendalii*.

#### Apoptosis assay by AO/EB staining method

The cytotoxic assay and cell cycle analyses indicated apoptosis when using the chloroform extracts from *B. ariensis* from Mexico and Morelos, *B. galeottiana* and *B. kerberi*. These results were confirmed by fluorescence microscopy analysis. The cells were scored under the fluorescence microscope in order to quantify viable cells. The analysis was repeated in triplicate and the images were captured in fluorescence and phase contrast at the same field, and in 10 and 40 x magnification. Fluorescence microscopic analysis showed that the untreated PC-3 cells were stained with uniform green fluorescence (Figures 2a and 2c); PC-3 cells in early apoptosis were stained with bright spots in green fluorescence (Figures 2b PTOX; 2e *B. ariensis* from Morelos; 2f *B. ariensis* from the state of Mexico; 2g *B. galeottiana* and 2h *B. kerberi*), in late apoptosis PC-3 cells were stained in orange fluorescence with condensed and fragmented cell nucleus (Figures 2g *B. galeottiana* and 2h *B. kerberi*), and in necrosis PC3 cells were stained in orange fluorescence, but nucleus morphology appeared without condensation of chromatin (Figure 2d, H<sub>2</sub>O<sub>2</sub>).

#### Anti-inflammatory assay on ear edema induced by TPA in mice

The anti-inflammatory effects of the *burseras* extracts are shown in Figure 3, and expressed as percentage of edema inhibition (% EI). The data indicated an important loss of weight by the extracts of *B. galeottiana* (89.92%), *B. excelsa* (90.79%) and *B. schlechtendalii* (91.60%). The Dunnett's test indicated that these three species were the most active extracts and statistically comparable to the control indomethacin. The best extracts with anti-inflammatory potential were selected to determine the average effective dose (ED<sub>50</sub>). *B. galeottiana* was the most active extract (ED<sub>50</sub>=0.23 ± 0.02 mg/ear) when compared to the positive control indomethacin (ED<sub>50</sub>=0.19 ± 0.02 mg /ear) (Table 4).

#### Discussion

Plants have a long history of application in the treatment of cancer, but many of the claims for the efficacy of such treatment should be viewed with some skepticism because cancer, as a specific disease entity, is likely to be poorly defined in terms of folklore and traditional medicine [27]. In our case, besides the claimed popular use of these plants, the chemotaxonomic approach proved to be effective, since it was possible to identify *Bursera* species with cytotoxic activity indicating or affirming its usefulness as a method of selection.

Sample	CI50 (mg/mL)
<i>B. excelsa</i>	0.26 ± 0.01
<i>B. galeottiana</i>	0.23 ± 0.02
<i>B. schlechtendalii</i>	0.25 ± 0.02
Indomethacine	0.19 ± 0.02

**Table 4:** Anti-inflammatory activity of chloroform extracts from selected *burseras*.

With the exception of *B. glabrifolia* from Morelos and *B. excelsa*, all the studied *bursera* species exhibited cytotoxic activity. *B. fagaroides* proved to be the most active plant studied in this investigation since all three extracts (hexane, chloroform and ethyl acetate) displayed important cytotoxic effect against the four carcinoma cell lines HF-6, MCF-7, KB and PC-3. The obtained results are in accordance with those previously reported by other authors who studied this plant but from a population collected in the state of Michoacán [11,16]. *B. schlechtendalii* showed important cytotoxic activity, and this finding is in accordance with McDonield and Cole [18] study, since they identified two lignans with anti tumor activity in this species. In the case of *B. ariensis* notable cytotoxic effects from the hexane extract against all the cancer cell lines were obtained, and although there are no previous reports that evaluate the cytotoxic activity of this species, a lignan, denominated 'ariensin', was isolated from *B. ariensis* growing in the state of Michoacán [27,28], and was then synthesized by Burke and Stevenson [29], as well as later identified in *Acanthopanax koreanum* [30].

Koulman in 2003 [31] did not perform cytotoxic studies in *B. excelsa* collected in the state of Mexico but he did isolate three compounds: iso-bursehernin, 3,4-dimethoxy-3',4'-methylenedioxy lignano-9,9'-epoxy lignan-9'-ol and guayadequiol, that he proposed as substrates for the synthesis of cytotoxic lignans. Some recent studies using NMR spectra of *B. ariensis* and *B. fagaroides* conducted by our research group indicated the presence of characteristic signals of aryltetralin lignans, whose structures are oxidized at C9 and C9' (signals situated at 4.56 ppm in the 1H NMR spectra). Since lignans are important compounds with attributed cytotoxic activity, the isolation and identification of these metabolites from the two mentioned species is an important accomplishment.

In our study, the hexane extract of *B. kerberi* showed cytotoxic effect against HF-6, MCF-7 and, the chloroform extract showed cytotoxic effect against HF-6, MCF-7, KB and PC-3. Hernandez et al. in 2005 [32] reported the isolation of verticillane derivatives from *B. kerberi* collected in Jalisco; and although there are no cytotoxic data from these diterpenes, they were proposed as biogenetic precursors of the cytotoxic taxanes. Several reports in the literature do show cytotoxic activity displayed by triterpene compounds obtained from

different species such as *Glochidion eriocarpum*, *Bridelia cambodiana* and species of *Protium* genus [33-35]. Some phytochemical studies report the presence of triterpenes in the *Bursera* genus [36-39], but only one investigation of the anti-tumoral activity of two triterpenes known as 'sapeline' A and B, isolated from *B. klugii*, showed activity against the P-388 lymphocytic leukemia test system (3PS), as well as toward the human epidermoid carcinoma of the nasopharynx (KB) [19]. In ongoing investigations conducted by our research group on the *Bursera* species selected for this study; *B. bicolor*, *B. lancifolia*, *B. linanoe*, *B. kerberi*, *B. excelsa*, *B. galeottiana* and *B. glabrifolia* showed <sup>1</sup>H NMR spectra with characteristic triterpene signals profiling from 0.5 to 2.5 ppm. Extracts from these species also gave a positive Liebermann-Buchard test, which is characteristic of triterpenoids.

The lack of cytotoxic activity from *B. bicolor*, *B. kerberi* and *B. galeottiana* extracts on normal fibroblasts HFS-30 merits further studies in search of potential remedies with therapeutic value, since they exert high cytotoxic activity against cancer cell lines. *B. bicolor* and *B. kerberi* chloroform extracts showed outstanding selective cytotoxicity against PC-3 cancer cell line (SI=153.85 and 40.82, respectively), and only *B. kerberi* showed important selective cytotoxicity against HF-6 (SI=37.74).

It is important to note that while *B. glabrifolia* collected in Oaxaca was toxic against HF-6 and MCF-7 cell lines, the population of this species collected in Morelos was non-toxic against the tested cancer cell lines. This variation indicates different chemical profiles between both populations, possibly related to geographical, climatic or soil conditions.

In relation to cell cycle analysis of their the DNA content, cells can be classified into three categories: cells in G1 do not present DNA duplication, cells in G2/M exhibit duplicated DNA, and cells in S phase present intermediate DNA content. In each phase of the cell cycle, important control mechanisms exists that involve checkpoints which ensure the proper execution of cell cycle events. The p53 protein checkpoint blocks the entry of cells to mitosis when DNA is damaged in G2/M phase. This protein (p53) can activate the transcription of several apoptosis associated genes to program cell death in response to genotoxic stresses [40,41]. In our study, the results show that with exception of *B. kerberi*; the chloroform extracts can effectively induce apoptosis in PC-3 cells. These findings support those results obtained with cell cycle and cytotoxic experiments. *B. galeottiana* chloroform extract showed an important inhibitory effect related to G2/M phase arrest, and it remains to be demonstrated if this is due to the inhibition of the p53 checkpoint. At this phase, cells undergo apoptosis since they cannot be repaired

The observation of apoptosis induced by the plant extracts could be a key factor in determining their efficacy, in as much as most tumor cells remain sensitive to some apoptotic stimuli from anticancer drugs.

An inflammation-cancer relation has been proposed many years ago, and the interaction between them has become more and more evident. Inflammation is an immediate host defense mechanism of the body to tissue injury caused by noxious stimuli, and is characterized by the release of mediators (prostaglandins, cytokines, chemotactic molecules and vasoactive peptides), which affect the cellular infiltration and vascular permeability. For example, colorectal cancer is linked to colitis, which increases the risk of colorectal cancer by 10-fold, and the treatment of patients with anti-inflammatory therapy reduces this risk [42]. The role of inflammation in epigenetic, and

changing genetic events associated with cancer is estimated in more than 25% of all cancers [43].

Two of the studied species (*B. galeottiana*, and *B. schlechtendalii*) showed important anti-inflammatory activity comparable to the control indomethacin, as well as cytotoxic effects against cancer cell lines. As mentioned in the introduction, several reports in the *Bursera* genus had identified triterpenes as the involved compounds the in anti-inflammatory response. The triterpenoids of the oleanane and ursane series displayed anti-inflammatory activity [44] and were isolated from *B. simaruba* leaves, *B. graveolens* and *B. lancifolia* [44-46].

The overall results of this investigation postulate the studied species of the *Bursera* genus as candidates to investigate for their therapeutic potential in the treatment of cancer diseases and inflammation. Clearly, our results are an important contribution to the understanding on the potentialities of the *Bursera* genus as a source of new medical approaches.

This is the first report that focuses a systematic study on the cytotoxic and anti-inflammatory activities of nine species (eleven populations) of the *Bursera* genus growing in Mexico. The correlation of this information with ethnomedical data will be of great benefit for the rational use of these plant species.

## References

1. Becerra JX, Venable DL (2008) Sources and sinks of diversification and conservation priorities for the Mexican tropical dry forest. PLoS One 3: e3436.
2. Alonso AJ, Villarreal ML, Salazar LA, Gómez M, Domínguez F, et al. (2011) Mexican medicinal plants used for cancer treatment: Pharmacological, phytochemical and ethnobotanical studies. J Ethnopharmacol 133: 945-72.
3. Lona NV (2012) Objects made of copal resin: a radiological analysis. Bol Soc Geol Mex 64: 207-213.
4. García-Silberman S (2002) An explanatory model of behavior toward mental illness. Salud Publica Mex 44: 289-296.
5. Sahagún B. de (2000) Historia general de las cosas de Nueva España. 3 vols. Estudio introductorio, paleografía, glosario y notas Alfredo López Austin y Josefina García Quintana: México, Cien de México/CONACULTA. p 1450.
6. Montemayor C, Frischmann D (2005) Words of the true peoples/ Palabras de los seres verdaderos: Anthology of Contemporary Mexican Indigenous-Language Writers/Antología de Escritores Actuales en Lenguas Indígenas de México: Volume Two/Tomo Dos: Poetry/Poesía. University of Texas Press. p 220.
7. Galeote M (1997) Nombres indígenas de plantas americanas en los tratados científicos de Fray Agustín Farfán. Universidad de Granada. p 161.
8. <http://www.medicinatradicionalmexicana.unam.mx/monografia.php> 3: 7989.
9. Loeza-Corte JM, Díaz-López E, Campos-Pastelín JM, Orlando-Guerrero JI (2013) Efecto de lignificación de estacas sobre enraizamiento de *Bursera morelensis* Ram. y *Bursera galeottiana* Engl. en la Universidad de la Cañada en Teotitlán de Flores Magón, Oaxaca, México. Ciencia Ergo Sum 20: 222-226.
10. Queiroga CL, Duarte MC, Ribeiro BB, de Magalhães PM (2007) Linalool production from the leaves of *Bursera aloexylon* and its antimicrobial activity. Fitoterapia 78: 327-328.
11. Puebla AM, Huacuja L, Rodríguez G, Villaseñor MM, Miranda ML, et al. (1998) Cytotoxic and antitumor activity from *Bursera fagaroides* ethanol extract in mice with L5178Y lymphoma. Phytoter Res 12: 545-548.

12. Elton A (1998) Documento técnico justificativo para la creación de la Reserva de la Biosfera Sierra Huautla-Cerro Frío Morelos. Instituto Nacional de Ecología, Centro de Investigación Ambiental e Investigación Sierra de Huautla, p 155.
13. Mohan S, Bustamam A, Ibrahim S, Al-Zubairi AS, Aspollah M, et al. (2011) In vitro ultramorphological assessment of apoptosis on CEMs induced by linoleic acid-rich fraction from *Typhonium flagelliforme* tuber. *Evidence-Based Complementary and Alternative Medicine*.
14. Maldonado B, Ortiz A, Dorado O (2004) Preparados galénicos e imágenes de plantas medicinales. CEAMISH-UAEM, México 7-11.
15. Téllez-Valdés O, Reyes-Castillo M, Dávila-Aranda P, Gutiérrez-García K, Téllez-Poo O, et al. (2009) Guía ecoturística. Las plantas del valle de Tehuacán-Cuicatlán. UNAM. p 41.
16. Rojas-Sepúlveda AM, Mendieta-Serrano M, Mojica MY, Salas-Vidal E, Marquina S, et al. (2012) Cytotoxic podophyllotoxin type-lignans from the steam bark of *Bursera fagaroides* var. *fagaroides*. *Molecules* 17: 9506-9519.
17. Cole JR, Bianchi E, Trumbull ER (1969) Antitumor agents from *Bursera microphylla* (Burseraceae). II. Isolation of a new lignan-burseran. *J Pharm Sci* 58: 175-176.
18. McDoniel PB, Cole JR (1972) Antitumor activity of *Bursera schlechtendalii* (burseraceae): isolation and structure determination of two new lignans. *J Pharm Sci* 61: 1992-1994.
19. Jolad SD, Wiedhopf RM, Cole JR (1977) Cytotoxic agents from *Bursera klugii* (Burseraceae) I: isolation of sapelins A and B. *J Pharm Sci* 66: 889-890.
20. Jolad SD, Wiedhopf RM, Cole JR (1977) Cytotoxic agents from *Bursera morelensis* (Burseraceae): deoxypodophyllotoxin and a new lignan, 5'-desmethoxydeoxypodophyllotoxin. *J Pharm Sci* 66: 892-893.
21. Houghton P, Fang R, Techatanawat I, Steventon G, Hylands PJ, et al. (2007) The sulphorhodamine (SRB) assay and other approaches to testing plant extracts and derived compounds for activities related to reputed anticancer activity. *Methods* 42: 377-387.
22. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, et al. (1990) New colorimetric cytotoxicity assay for anticancer-drug screening. *J Natl Cancer Inst* 82: 1107-1112.
23. Suffness M, Pezzuto J (1991) Assays related to cancer drug discovery. In: Dey P, Harborne J, (eds.) *Methods in Plant Biochemistry*. London: AC. 71-133.
24. Kasibhatla S, Amarante-Mendes GP, Finucane D, Brunner T, Bossy-Wetzel E, et al. (2006) Acridine Orange/Ethidium Bromide (AO/EB) Staining to Detect Apoptosis. *CSH Protoc* 2006.
25. Yang GY, Liao J, Kim K, Yurkow EJ, Yang CS (1998) Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis* 19: 611-616.
26. Rao TS, Currie JL, Shaffer AF, Isakson PC (1993) Comparative evaluation of arachidonic acid (AA)- and tetradecanoylphorbol acetate (TPA)-induced dermal inflammation. *Inflammation* 17: 723-741.
27. Cragg GM, Newman DJ (2008) Anticancer drug discovery and development from natural products. In: Colegate S, Molyneux R, (eds.) *Bioactive Natural Products, Detection, Isolation and Structural Determination*. Boca Raton: CRC Press. 323-370.
28. Hernández JD, Román LU, Espiñeira J, Joseph-Nathan P (1983) Ariensin, a new lignan from *Bursera ariensis*. *Planta Med* 47: 215-217.
29. Burke JM, Stevenson R (1985) Synthesis of (+/-)-ariensin. *Planta Med* 51: 450-452.
30. Rao V (2012) Phytochemicals and their pharmacological aspects of *Acanthopanax koreanum*. In: Kim YH, Kim JA, Nhiem NX (eds.) *Phytochemicals Rijeka, Croatia*: In Tech 451-466.
31. Koulman A (2003) Evolution of lignan biosynthesis in the genus *Bursera*. In: Koulman A, Quax WJ, Judith X, Becerra JX, (eds.), *Podophyllotoxin*. Netherlands: Stichting Regenboog Drukkerij Press 141-156.
32. Hernández-Hernández JD, Román-Marín LU, Cerda-García-Rojas CM, Joseph-Nathan P (2005) Verticillane derivatives from *Bursera suntui* and *Bursera kerberi*. *J Nat Prod* 68: 1598-1602.
33. Khiev P, Cai XF, Chin YW, Ahn KS, Lee HK, et al. (2009) Cytotoxic terpenoids from the methanolic extract of *Bridelia cambodiana*. *J Korean Soc Appl Biol Chem* 52: 626-631.
34. Nhiem NX, Thu VK, Van Kiem P, Van Minh C, Tai BH, et al. (2012) Cytotoxic oleane-type triterpene saponins from *Glochidion eriocarpum*. *Arch Pharm Res* 35: 19-26.
35. Barros FW, Bandeira PN, Lima DJ, Meira AS, de Farias SS, et al. (2011) Amyrin esters induce cell death by apoptosis in HL-60 leukemia cells. *Bioorg Med Chem* 19: 1268-1276.
36. Ara K, Rahman MS, Rahman AHMM, Hassan CM, Rashid MA (2009) Terpenoids and coumarin from *Bursera serrata*. *J Pharm Sci* 8: 107-110.
37. Moreno J, Rojas LB, Aparicio R, Marcó LM, Usbillaga A (2010) Chemical composition of the essential oil from the bark of *Bursera tomentosa* (Jacq) Tr and Planch. *Bol Latinoam Caribe Plant Med Aromát.* 9: 491-494.
38. Cárdenas R, Reguera JJ, Llanos E, Aguirre E, Herrera J, et al. (2012) Effects of organic extracts of *Bursera copallifera* and *B. lancifolia* leaves in the development of *Spodoptera frugiperda*. *J Entomol* 9: 115-122.
39. Ionescu F, Jolad SD, Cole JR (1977) The structure of benulin, a new pentacyclic triterpene hemiketal isolated from *Bursera arida* (Burseraceae). *J Org Chem* 42: 1627-1629.
40. Luk SC, Siu SW, Lai CK, Wu YJ, Pang SF (2005) Cell Cycle Arrest by a Natural Product via G2/M Checkpoint. *Int J Med Sci* 2: 64-69.
41. Li H, Wang P, Liu Q, Cheng X, Zhou Y, et al. (2012) Cell cycle arrest and cell apoptosis induced by *Equisetum hyemale* extract in murine leukemia L1210 cells. *J Ethnopharmacol* 144: 322-327.
42. Reuter S, Gupta SC, Chaturvedi MM, Aggarwal BB (2010) Oxidative stress, inflammation, and cancer: how are they linked? *Free Radic Biol Med* 49: 1603-1616.
43. Kundu JK, Surh YJ (2012) Emerging avenues linking inflammation and cancer. *Free Radic Biol Med* 52: 2013-2037.
44. Carretero ME, López-Pérez JL, Abad MJ, Bermejo P, Tillet S, et al. (2008) Preliminary study of the anti-inflammatory activity of hexane extract and fractions from *Bursera simaruba* (Linneo) Sarg. (Burseraceae) leaves. *J Ethnopharmacol* 116: 11-15.
45. Robles J, Torrenegra R, Gray RI, Piñeros C, Ortiz L et al. (2005) Triterpenos aislados de la corteza de *Bursera graveolens* (Burseraceae) y su actividad biológica. *Braz J Pharmacog* 15: 283-286.
46. Zúñiga B, Guevara-Fefer P, Herrera J, Contreras JL, Velasco L, et al. (2005) Chemical composition and anti-inflammatory activity of the volatile fractions from the bark of eight Mexican *Bursera* species. *Planta Med* 71: 825-828.