

## Dairy Ruminant Nutrient Intake Orchestration: A Novel Chronophysiological Discipline

Akbar Nikkhah\*

Chief Highly Distinguished Professor, Department of Animal Sciences, Faculty of Agricultural Sciences, University of Zanjan, Zanjan, Iran

\*Corresponding author: Akbar Nikkhah, Chief Highly Distinguished Professor, Department of Animal Sciences, Faculty of Agricultural Sciences, University of Zanjan, Zanjan 313-45195, Iran, Tel: +98-241-5152801; Fax: +98-241-5283202; E-mail: [anikkha@yahoo.com](mailto:anikkha@yahoo.com)

Rec date: Jul 19, 2014; Acc date: Jul 21, 2014; Pub date: Jul 23, 2014

Copyright: © 2014 Nikkhah A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Editorial

Chronophysiology as an evolutionary multiscience has enabled ruminants to better cope with the changing environment from the mother nature towards modernity. With the recent discoveries on timing of feeding orchestrating circadian and postprandial nutrient intake and metabolic rhythms, time has already come to contemplate chronophysiology as a foundation of voluntary feed intake (VFI) regulation in dairy ruminants as workable models for humans.

This editorial describes evolving chronophysiological mediations of VFI in ruminants of especially dairy cows. Chronophysiology has been a key missing interscience in realistic modeling and understanding of VFI regulation in high-producing ruminants. The shortcoming gains increasing thoughts given that accurate and reliable nutrient intake predictions are essential to consistent meeting of nutrient requirements and, thus, to profitable human food production. With the recent chronophysiological discoveries on dairy farm management strategies, time has already come to realize that evolutionary circadian rhythms in grazing, chewing, rumen kinetics, and peripheral substrate assimilation can help develop innovative strategies to improve VFI prediction accuracy and dairy ruminant efficiency and health [1].

Inspired by the nature, ruminants have evolved to ruminate habitually overnight and graze during day, particularly just about sunrise and sunset. Accordingly, rumen fermentation, postrumen assimilation and peripheral metabolism have developed specialized circadian rhythms [2,3]. These evolutionary rhythms in feeding behavior and metabolism have generated circannual, seasonal, and circadian rhythms in endocrinology. The drastic changes in production systems due to modernization have altered such natural eating, ruminating and rumen fermentation rhythms. Therefore, it has become vital to preserve synchronies between external cues and dairy ruminant internal state [2,4].

Most recent findings suggest altered postfeeding VFI rhythms of nongrazing dairy cows by shifting feeding timing [1,5]. Eating rate and VFI, especially shortly postfeeding, have increased by night instead of day feeding. As a result, rhythms of circadian and postprandial rumen

fermentation and intermediary metabolism have changed [6,7]. Timing of feeding/eating as a key external manager, thus, determines the rate and extent of VFI, rumination, fermentation and splanchnoperipheral metabolism in dairy cows.

The impacts of chronophysiological management strategies (e.g., eating timing) should be quantified for inclusion in dairy cow nutrient intake prediction and production estimation models to improve accuracy and practicality under varying physiological and management states. This will help optimize ruminant efficiency, food safety and security, and environmental quality interconnectedly. Such practical ruminant models greatly serve human health initiatives, particularly on overcoming challenges of obesity, diabetes, and cardiovascular diseases.

### Acknowledgments

The Ministry of Science, Research and Technology and University of Zanjan are thankfully acknowledged for supporting the author's global programs of optimizing the new millennium science edification.

### References

1. Nikkhah A (2013) Chronophysiology of ruminant feeding behavior and metabolism: an evolutionary review. *Biol Rhythm Res* 44: 197-218.
2. Nikkhah A (2011) Bioscience of ruminant intake evolution: feeding time models. *Adv Biosci Biotechnol* 2: 271-274.
3. Taweel HZ, Tas BM, Dijkstra J, Tamminga S (2004) Intake regulation and grazing behavior of dairy cows under continuous stocking. *J Dairy Sci* 87: 3417-3427.
4. Nikkhah A (2011) Ruminant chronophysiological management: an emerging bioscience. *Open Access Anim Physiol* 3: 9-12.
5. Nikkhah A (2011) Science of eating time: A novel chronophysiological approach to optimize glucose-insulin dynamics and health. *J Diab Mellit* 2: 8-11.
6. Nikkhah A, Furedi C, Kennedy A, Wittenberg K, Plaizier JC (2010) Feed delivery at 2100 h vs. 0900 h for lactating dairy cows. *Can J Anim Sci* 91: 113-122.
7. Nikkhah A (2012) Timing of feed presentation entrains periprandial rhythms of energy metabolism indicators in once-daily fed lactating cows. *Biol Rhythm Res* 43: 651-661.