

DAT Conformation Does Not Predict the Ability of Atypical Dopamine Uptake Inhibitors to Substitute for Cocaine

Takato Hiranita*

Division of Neurotoxicology, National Center for Toxicological Research (NCTR), U.S. Food and Drug Administration (FDA), Jefferson, USA

*Corresponding author: Takato Hiranita, Division of Neurotoxicology, National Center for Toxicological Research (NCTR), U.S. Food and Drug Administration (FDA), 3900 NCTR Road, Jefferson, AR 72079-9501, USA, Tel: 870-543-6823; Fax: 870-543-7745; E-mail: takato.hiranita@fda.hhs.gov

Received date: August 02, 2016; Accepted date: August 03, 2016; Published date: August 08, 2016

Copyright: © 2016 Hiranita T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

The reuptake inhibition of dopamine (DA) through the DA transporter (DAT) is thought to be the primary mechanism underlying the reinforcing effects of cocaine [1,2]. However, there are several DA reuptake inhibitors that do not produce appreciable cocaine-like effects *in vivo* [2-4]. Previously, there were several relatively viable hypotheses related to the mechanisms underlying the atypical effects of such agents, but new studies have decreased this number [3,5]. The new findings may contribute to the development of a better understanding of what constitutes cocaine-like behavioral effects.

It is well known that the DAT plays an important role in cocaine reinforcement. For example, cocaine has a relatively high affinity for the DAT [6]. Further, various DA reuptake inhibitors can maintain self-administration responding above vehicle levels when substituted for cocaine and potentiate cocaine self-administration when used as a pretreatment. These include standard DA reuptake inhibitors including WIN 35,428, methylphenidate and nomifensine [2,3,7-9]. However, several DA reuptake inhibitors are not reinforcing when substituted for cocaine and antagonize cocaine self-administration when used as a pretreatment. These include the atypical DA uptake inhibitors, JHW 007 and RTI-371 [2,3,7]. The attributes that contribute to their atypical effects as DA reuptake inhibitors are unknown but there are several relatively viable hypotheses that address this issue [10,11]. One of such hypotheses is that conformational differences may affect DAT binding.

Studies evaluating the accessibility of the sulfhydryl-reactive reagent [2-(trimethylammonium)ethyl]-methanethiosulfonate to an internal cysteine residue of the DAT (I159C), which is only accessible when the extracellular DAT gate is open, have indicated that cocaine and its analogue WIN 35,428 bind to an open DAT conformation at synaptic clefts (the 'outward-facing' conformation), whereas several atypical DA

reuptake inhibitors (e.g. JHW 007) bind to the 'closed' or 'inward-facing' conformation [10,12]. Thus, binding to the inward-facing conformation appears to be important for the atypical effects of these agents. In contrast to cocaine, JHW 007 was shown to be inactive in locomotor activity assessments and cocaine discrimination and self-administration procedures (Table 1). Both RTI-371 and RTI-336 bind to the outward-facing conformation of the DAT (Table 1), but only RTI-336 substituted for cocaine using cocaine self-administration procedures (Table 1). A more recent study assessed whether preference for the DAT conformation is predictive of the cocaine-like behavioral effects of various novel DA reuptake inhibitors using locomotor activity assessments and cocaine discrimination procedures (Table 1). Among a series of novel compounds [LX-10, -11, -12, 13, -16, -19, -21, -22, -23 and -24 in Table 1, assumedly synthesized by Lifeng Xu (LX), one of the authors] that have been shown to preferentially bind to the outward-facing conformations of the DAT, only LX-10 was cocaine-like in both behavioral assessments. In contrast, six LX compounds (LX-11, -12, -21, -22, -23 and -24) were inactive in both assessments. In addition, LX-13, -16 and -19 stimulated locomotor activity but did not fully substitute for cocaine using cocaine discrimination procedures. Thus, the conformation of the DAT is not a viable determinant of the production of cocaine-like behavioral outcomes.

In summary, new findings suggest that the conformation of the DAT is not predictive of the behavioral effects of DA reuptake inhibitors. Differences in kinetic variables and DAT/ σ -receptor dual inhibition are two remaining relative viable hypotheses for addressing the mechanisms underlying the reduced cocaine-like effects of atypical DA reuptake inhibitors [10,11,13]. Future studies to explore these two hypotheses should facilitate the development of effective medication(s) for cocaine abuse.

Compound [reference]	DAT conformation (μ M)	Locomotor activity (μ mol/kg, I.P.)	Substitution for cocaine in cocaine discrimination (μ mol/kg, I.P.)	Substitution for cocaine self-administration (mg/kg/injection, I.V.)
Cocaine [3,5]	Outward-facing (100)	Increased (30-100) Increased (20 or 40 mg/kg, I.P.)	Substituted (30)	Substituted (0.1 and 0.32)
JHW 007 [2,3]	Inward-facing (1)	Inactive (0.3-3.0 mg/kg, I.P.)	Not substituted (0.17-5.6 mg/kg, I.P.)	Not substituted (0.032-1.0)
RTI-336 [3]	Outward-facing (1)	Increased (10-56 mg/kg, I.P.)	Substituted (56 mg/kg, I.P.)	Substituted (0.1 and 0.32)
RTI-337 [3]	Outward-facing (1)	Increased (100 mg/kg, I.P.)	Partially substituted (10-100 mg/kg, I.P.)	Not substituted (0.1-3.2)
LX-10 [5]	Outward-facing (10)	Increased (3 and 10)	Substituted (10)	N.T.

LX-11 [5]	Outward-facing (20)	Inactive (3-56)	Not substituted (17)	N.T.
LX-12 [5]	Outward-facing (20)	Inactive (3-300)	Not substituted (0.3-10)	N.T.
LX-13 [5]	Outward-facing (10)	Increased (10 and 30)	Not substituted (3-17)	N.T.
LX-15 [5]	No change (20)	Inactive (3-100)	Not substituted (0.3-10)	N.T.
LX-16 [5]	Outward-facing (10)	Increased (3-100)	Partially substituted (0.3-3)	N.T.
LX-19 [5]	Outward-facing (10)	Increased (30-100)	Partially substituted (0.3-10)	N.T.
LX-20 [5]	No change (20)	Inactive (30)	Not substituted (0.3-3)	N.T.
LX-21 [5]	Outward-facing (10)	Inactive (0.56-170)	Not substituted (0.3-3)	N.T.
LX-22 [5]	Outward-facing (20)	Inactive (0.56-300)	Not substituted (0.3-10)	N.T.
LX-23 [5]	Outward-facing (10)	Inactive (0.56-56)	Not substituted (0.3-5.6)	N.T.
LX-24 [5]	Outward-facing (20)	Inactive (1.7-56)	Not substituted (0.3-5.6)	N.T.
NT: Not Tested				

Table 1: Preference of the DAT conformation, effects on locomotor activity and substitution for cocaine under cocaine discrimination or self-administration procedures.

Acknowledgment

The present work was supported by the Division of Neurotoxicology/NCTR/U.S. FDA (Dr. Merle G. Paule). The information in this article is not a formal dissemination of information by the FDA and does not represent agency position or policy. The author thanks Drs. Merle G. Paule, David A. Thorn, Tylor Ingle and Mallikarjuna Basavarajappa at NCTR for comments during the preparation of the manuscript.

References

- Ritz MC, Lamb RJ, Kuhar MJ (1987) Cocaine receptors on dopamine transporters are related to self-administration of cocaine. *Science* 237: 1219-1223.
- Hiranita T, Soto PL, Newman AH, Katz JL (2009) Assessment of reinforcing effects of benzotropine analogs and their effects on cocaine self-administration in rats: comparisons with monoamine uptake inhibitors. *J Pharmacol Exp Ther* 329: 677-686.
- Hiranita T, Wilkinson DS, Hong WC, Zou MF, Kopajtic TA, et al. (2014) 2-isoxazol-3-phenyltropane derivatives of cocaine: molecular and atypical system effects at the dopamine transporter. *J Pharmacol Exp Ther* 349: 297-309.
- Katz JL, Kopajtic TA, Agoston GE, Newman AH (2004) Effects of N-substituted analogs of benzotropine: diminished cocaine-like effects in dopamine transporter ligands. *J Pharmacol Exp Ther* 309: 650-660.
- Hong WC, Kopajtic TA, Xu L, Lomenzo SA, Jean B, et al. (2016) 2-Substituted 3 β -Aryltropane Cocaine Analogs Produce Atypical Effects without Inducing Inward-Facing Dopamine Transporter Conformations. *J Pharmacol Exp Ther* 356: 624-634.
- Garcés-Ramírez L, Green JL, Hiranita T, Kopajtic TA, Mereu M, et al. (2011) Sigma receptor agonists: Receptor binding and effects on mesolimbic dopamine neurotransmission assessed by microdialysis in rats. *Biol Psychiatry* 69: 208-217.
- Hiranita T, Soto PL, Kohut SJ, Kopajtic T, Cao J, et al. (2011) Decreases in cocaine self-administration with dual inhibition of the dopamine transporter and sigma receptors. *J Pharmacol Exp Ther* 339: 662-677.
- Barrett AC, Miller JR, Dohrmann JM, Caine SB (2004) Effects of dopamine indirect agonists and selective D1-like and D2-like agonists and antagonists on cocaine self-administration and food maintained responding in rats. *Neuropharmacology* 47: 256-273.
- Schenk S (2002) Effects of GBR 12909, WIN 35,428 and indatraline on cocaine self-administration and cocaine seeking in rats. *Psychopharmacology (Berl)* 160: 263-270.
- Reith ME, Blough BE, Hong WC, Jones KT, Schmitt KC, et al. (2015) Behavioral, biological, and chemical perspectives on atypical agents targeting the dopamine transporter. *Drug Alcohol Depend* 147: 1-19.
- Hiranita T (2015) Cocaine Antagonists; Studies on Cocaine Self-Administration. *J Alcohol Drug Depend* 3: e125.
- Loland CJ, Desai RI, Zou MF, Cao J, Grundt P, et al. (2008) Relationship between conformational changes in the dopamine transporter and cocaine-like subjective effects of uptake inhibitors. *Mol Pharmacol* 73: 813-823.
- Katz JL, Hiranita T, Kopajtic TA, Rice KC, Mesangeau C, et al. (2016) Blockade of Cocaine or sigma Receptor Agonist Self Administration by Subtype-Selective sigma Receptor Antagonists. *J Pharmacol Exp Ther* 358: 109-124.