

Development and Quality Evaluation of Tamarind Plum Blended Squash During Storage

Ibrahim Khan^{1*}, Rehman AU¹, Khan SH², Qazi IM¹, Arsalan Khan², Shah FN² and Rehman TU⁴

¹The University of Agriculture Peshawar, Khyber Pakhtunkhwa, Pakistan

²Agricultural Research Institute ARI Tamab Peshawar, Khyber Pakhtunkhwa, Pakistan

³Gomal University of D.I. Khan, Pakistan

⁴Abdul Wali Khan University, Mardan, Pakistan

Abstract

The achievement was done to study the combination of tamarind plum blended squash for 90 days' interval at room temperature. Tamarind and plum was added at a combination of 750: 0, 650: 100, 550: 200, 450: 300, 350:400, 250:500, 150:600 and 50:700 represent each treatment. The prepared tamarind plum blended squash was analyzed physio-chemically for TSS, Ascorbic acid, acidity, sugar acid ration, pH, reducing and non-reducing sugar, organoleptically for taste, color, texture and overall acceptability for a total period of 90 days. The result of the statistical analysis showed that treatment and storage interval shows a significant ($P < 0.05$) effect both physio-chemical and organoleptic evaluation. Results also revealed that the decrease occurred in ascorbic acid content from (39.49 mg/100 gm to 27.40 mg/100 gm), titratable acidity (1.09% to 0.98%), non-reducing sugar (44.36% to 21.97%), and sensory evaluation included taste (6.85 to 5.83), color (6.33 to 5.36), flavor (7.54 to 5.75) and overall acceptability (8.03 to 6.14) while increased was found in total soluble solid (48.98°brix to 49.61°brix), sugar-acid ratio (44.94 to 50.79), pH (2.77 to 2.84), reducing sugar (17.21% to 31.23%) during storage. The maximum mean values were observed for TSS is TPS_7 (51.64°brix), ascorbic acid TPS_7 (37.87 mg/ 100 gm), titratable acidity TPS_1 (2.31%), sugar acid ratio TPS_0 (50.55), pH TPS_7 (2.93), reducing sugar TPS_0 (25.32%), non-reducing sugar TPS_4 (37.64%), color TPS_5 (6.70), flavour TPS_5 (7.54), taste TPS_5 (7.00) and overall acceptability TPS_5 (7.76). Among all the treatment TPS_5 was found to be the best. The result revealed that significant ($P < 0.05$) decreased was found in physio-chemical and organoleptic parameter of treatment TPS_5 .

Keywords: Plum; Tamarind; Vitamin C; Sodium benzoate; Acidity

Introduction

Tamarind (*Tamarind indica L.*) belongs to *Caesalpiniaceae* family. It is mostly grow in tropical Africa but has become naturalized in North and South America from Florida to Brazil, also grown in subtropical China, India, Pakistan, China, Thailand, Philippines, Indonesia and Spain. Tamarind fruit can be used for many purposes such as digestive, carminative, laxative, expectorant and tonic blood [1]. Tamarind pulp has medicinal purposes also and continues to be used by many people in Africa, Asia and America [2]. Tamarind juice have certain disadvantages such as unappetizing color, loss of fresh taste and spoiled easily [3] and hypoglycemic activity [4]. Tamarind pulp is mainly used for souring food products like chutneys, sambar, curries and sauces. Tamarind pulp is also used in preparations of jams, jellies, ice-creams, wine like beverages, canned tamarind juice and syrup. It is also enjoyed in the form of refreshing drinks and beverages. Fruit are commonly processed into juices, nectars, fruit punch, concentrates, glazed and crystallized fruit. The pulp can be used with original flavor after thermal processing [2]. Tamarind fruit contain low water content and is difficult to extract pulp from the fruit. With the advancing of technologies pulp of tamarind can be extracted by conventional processing techniques like soaking, maceration and straining. With the use of such techniques we can easily extract pulp [5]. The pulp of tamarind contains tartaric acid, reducing sugars, pectin, proteins, fiber, and cellulosic materials. The acid and sugar contents differ from sample to sample; for example, tartaric acid: 8%-18%, reducing sugars 25% to 45%, pectin 2% to 3.5% and proteins 2% to 3% [6]. Tamarind pulp has rich aroma and pleasant acidic taste which is widely used as a chief souring agent for curries, sauces, and certain beverages. The pulp also used as a raw material for the preparation of wine like beverages [7].

Plum (*Prunus domestica L.*) is highly perishable climacteric stone fruit and has short shelf-life at optimal temperatures. Decay of plum fruit may be due to mold growth and rapid ripening during storage.

Shelf life of plum can be extend through proper handling, transportation and marketing chain and also to kept in low temperature storage to extent postharvest quality of the fruit [8]. Plum also called as stones fruits consist of a solid covering with seed enclosed. The enclosed seed of plum is richest in proteins, lipids thus, they maybe a cheap source of different substances that could be useful for food, cosmetic, and pharmaceutical industries. The lipid content of plum seeds has already been explored. Plums contain red flesh and peel and are very exciting fruit due to their high content on bioactive compounds, such as the anthocyanins and other polyphenolic compounds with a high antioxidant capacity [9]. These natural substances found in plum acts to prevent diseases such as diabetes and cancer [10]. Concentrated soft drinks are used for refreshing purpose and are very popular drink contains certain proportion of juice. The summer season of Pakistan is long there for mostly people uses such type of beverages. Such type of activities like production, preservation and sale of these beverages provide commercial importance to our country [11]. Fruit beverages are a combination of products containing pulp, juice and water as well as sweetener, coloring, flavoring, and preservatives. Although fruit ingredient present in beverages has a dominant role of providing flavor and overall character, such types of products differ from fruit

***Corresponding author:** Ibrahim Khan, Food Science and Technology, Rahat Abad, House No 41, Peshawar, Khyber Pakhtoon Khwa-25000, Pakistan, Tel: +923003585403; E-mail: ibrahimfst339@gmail.com

Received February 10, 2017; **Accepted** March 06, 2017; **Published** March 13, 2017

Citation: Khan I, Rehman AU, Khan SH, Qazi IM, Khan A, et al. (2017) Development and Quality Evaluation of Tamarind Plum Blended Squash During Storage. J Food Process Technol 8: 662. doi: [10.4172/2157-7110.1000662](https://doi.org/10.4172/2157-7110.1000662)

Copyright: © 2017 Khan I, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Treatments	Tamarind juice (ml)	Plum juice (ml)	CMC (g)	Sugar (kg)	Water (ml)	potassium meta-bi-sulphite (%)
TPS ₀	750	–	2	1	250	0.1
TPS ₁	650	100	2	1	250	0.1
TPS ₂	550	200	2	1	250	0.1
TPS ₃	450	300	2	1	250	0.1
TPS ₄	350	400	2	1	250	0.1
TPS ₅	250	500	2	1	250	0.1
TPS ₆	150	600	2	1	250	0.1
TPS ₇	50	700	2	1	250	0.1

Table 1: Proposed plan of study for research.

juices and are labeled accordingly [12]. Keeping in view the importance of tamarind plum fruit; the plum and tamarind blended squash is developed.

Objectives

- To produce value added beverage from blends of tamarind plum.
- To develop suitable combination of tamarind plum blended squash.
- To analyze tamarind plum blended squash for physicochemical and sensory characteristic during storage.

Materials and Methods

Selection of fruits

Tamarind and Plum fruit at optimum maturity were purchased from the local market of Peshawar and was brought to the laboratory of Food Technology section, ARI Tarnab, Peshawar, for preparation of tamarind plum blended squash.

Pretreatment of blended squash

Tamarind and Plum fruit were carefully sorted to discard diseased, damaged, bruised and immature fruits. Then sorted fruits were thoroughly washed with tap water and the water was drained off. The unwanted portion was removed by trimming. The pulp was extracted by using pulping machine (Model.35027, Rochdale England).

Preparation of blended Squash

Tamarind and plum fruit blended squash were prepared following the method of Archana and Laxman [13], showed in (Table 1). The materials were added following the ratio 4:3:1 of sugar, pulp and water respectively.

Packaging and storage

The prepared squash was packed in PET bottles and was stored at room temperature for 3 months and was study for physico-chemical characteristics and sensory attributes at 15 days of intervals.

Physicochemical analysis

The prepared squash was examining for pH, TSS, Titratable acidity, Vit C, reducing and non-reducing sugar, sugar acid ratio was calculated from the data of TSS and titratable acidity and was measured by method of AOAC [14].

Total soluble solids

TSS (°brix) were find out by the standard method of AOAC [14], method no, 932.14 and 932.12. TSS (°brix) of the blended squash was finding out using hand refractometer. The instrument was calibrated and takes the reading accurately by putting a minute quantity of tamarind plum blended squash.

Titratable acidity

Preparations of standard solution 0.1 N NaOH: Take 6.30 g of oxalic acid and 4.5 g of NaOH in a volumetric flask and add distilled water in it to make a volume of 1 liter separately. Take 10 ml of 0.1 N solution of NaOH and titrate against 0.1 N solution of oxalic acid. Add 3 drops of phenolphthalein (indicator). Repeat the experiment for three times. Taken the reading till pink color is appears.

Titration of sample: Take 10 ml of squash sample, dissolved in distilled water to make a volume of 100 ml. Then take 10 ml of sample solution and add two drops of phenolphthalein and titrate along 0.1N NaOH solution. Repeat the experiment for 3 times to reduce error. Take the reading when pink color is appears.

$$\text{Acidity}(\%) = \frac{\text{CF} \times \text{N} \times \text{T} \times \text{D} \times 100}{\text{V} \times \text{S}}$$

Where:

C.F = Correction Factor for acidity.

N = Normality of sodium hydroxide used.

T = ml of sodium hydroxide used.

D = Dilution Factor for sample.

V = Sample taken for dilution.

S = Sample taken for titration.

Sugar acid ratio

Sugar acid ratio for tamarind plum blended squash was calculated using the formula.

$$\text{Sugar acid ratio} = \frac{\text{Total Soluble Solids (TSS)}}{\text{Titratable acidity}(\%)}$$

pH

The pH; is hydrogen ion concentration and it ranges from 1 to 14 that shows acidity and alkalinity of the sample, while the pH with 7 is neutral that is pure water indication. To find out pH of the sample, proper method of AOAC [14], 2005.02 was applied. Switch on the pH meter and standardized with the buffer solution of pH 4 and pH 7, respectively. Take 10 ml of tamarind plum sample in a beaker and put the electrode in it and note the result.

Reducing sugar

To analyze/ reducing sugar of tamarind plum blended squash standard method' of AOAC [14], 920.183 was' applied.

Reagents

Fehling A: Dissolved. 34.65g of CuSO₄·5H₂O in 500 ml of distilled, water.

Fehling B: Take 173 g of potassium/ titrate and 50gg of NaOH in beaker, dissolve it in 10 ml of water. The prepared solution was; taken and put into 500 ml conical flask and volume was prepared up to the mark by means of distillation water.

Methylene blue: Methylene blue is an indicator. Take 0.2 g of methylene; blue in 100 ml of volumetric flask and dissolve it in 150 ml of distilled water and the level was]made up to the spot, through further addition distilled water.

Procedure: Take 10 ml of tamarind plum blended squash sample and add distilled water to make the exact volume of 100 ml. Then take 5 ml of Fehling A and 5 ml of Fehling B, with 10 ml of distilled water was taken in conical flask. Heat was given to the flask till boiling start. Add the solution from the burette drop by drop till color becomes bricks red. 2 drops of methylene blue was added in a boiling solution. If color changes from red to blue the reaction needed to add extra tamarind plum solution till brick red color persists.

Calculation: Amount of Fehling A is 5 ml + % ml of Fehling B = X ml of the 10% of sample solution is equal of 0.05 g of reducing sugar × 100 ml of 10 % sample solution will contain.

$$100 \text{ ml of } 10 \% \text{ solution will contain} = \frac{0.05 \times 100}{X \text{ ml}} = Y \text{ g of reducing sugar}$$

$$\text{Reducing sugar} (\%) = \frac{Y \times 100}{10}$$

Non-reducing sugars

To investigate non-reducing sugar of tamarind plum blended squash standard method of AOAC [14], 920.184 was applied.

Procedure: 10 ml of sample was taken in volumetric flask and volume was made 100 ml with distill water. 20 ml of solution was taken and dilute with 10 ml of 1 N HCl. Mixture was heated till boiling, 10 ml of 1 N of NaOH was added after cooling and volume was made 250 ml. Take 5 ml Fehling A and B solution and dilute with 10 ml distilled water. Heat the solution to boiling and add tamarind plum blended diluted solution drop by drop till red brick color appears. Add 2 drops of methylene blue to check either the reaction is completed or not. For determination of non-reducing sugar the following formula was applied.

Calculations: Solution is equal to X ml = 0.05 g of reducing sugars

250 ml of sample contains = $250 \times 0.05 / \text{ml} = Y \text{ g of reducing sugars}$

This 250 ml of sample solution was prepared from 20 ml of 10%.

Sample solution contains $Y \times 100 / 20 = P \text{ g reducing sugar.}$

10 ml of sample solution contain = P g of reducing sugar.

100 ml of sample solution contain = $P \times 100/10 = Q \text{ g of total reducing sugar.}$

Q g of reducing sugar = inverted sugar + free reducing sugar.

Formula for non-reducing sugar is = total reducing sugar – free reducing sugar.

Ascorbic acid

Preparation of standard solutions: 42 mg of sodium bicarbonates (NaHCO_3) and 50 mg of 2,6 dichlorophenol indophenols dye to make the volume of 250 ml with distilled water. To prepare standard solution of Vitamin C take 50 mg of ascorbic acid and poured in 50 ml 0.4%

of oxalic acid solution. Keep the solution for 24 hours. Take 5 ml of ascorbic acid solution and titrate along dye till pink color appears and persists for one minute. Formula used to find out dye factor.

$$\text{Dye factor (F)} = \frac{\text{vitamin C solution taken in ml}}{\text{Volume of used dye}}$$

Titration of sample: Take 10 ml of tamarind plum blended squash and make a volume of 100 ml with 0.4% oxalic acid solution. 10 ml of sample solution were taken in a flask and titrate along dye to appear pink color and persist for 15 sec. Formula for Vitamin C content is:

$$\text{Ascorbic acid (mg/100 g)} = \frac{F \times T \times 100}{S \times D}$$

Where,

F = Standardization factor = ml of ascorbic acid / ml of pigment used.

T = ml of pigment used for sample.

S = ml of diluted sample taken for titration.

D = ml of sample taken for dilution.

Sensory evaluation

The samples of tamarind plum blended squash were sensory evaluated for color, texture, flavor and overall acceptability by 10 trained judge's panel. Organoleptic study was carried out for about 3 month. The evaluations were done using 9 points hedonic scale of Larmand [15].

Statistical analysis

All the data concerning treatments and storage interval were statistically analyzed by means of complete Randomized Design (CRD) 2 Factorial as recommended by Hicks [12] and the means were find out using least significant difference (LSD) Test at 5% possibility level.

Result and Discussion

Total soluble solids (°brix)

According to Table 2 the sample of tamarind plum blended squash were studied for TSS (°brix). The data of the samples shows significant ($P < 0.05$) increase during keeping time of storage. The sample of the tamarind plum blended squash were in the range of 47.45 (TPS_2) to 51.35 (TPS_7). The TSS of all the samples was gradually rises from 48.29 (TPS_0) to 51.94 (TPS_7) during 90 days of storage. Table 2 also showed that minimum mean TSS value was recorded for TPS_2 (47.78) while TPS_7 had maximum mean value of (51.64). Similarly, this is also observed from the data that maximum percent increase in TSS was found in TPS_2 (1.76), while a sample TPS_5 (1.05) had a minimum percent increase. The data showed a significant change in blended squash of tamarind plum during storage. Similar observations were recorded by Kotecha and Kadam [16] in tamarind syrup and Nath et al. [17] in ginger blended with mandarin squash that because of hydrolysis of polysaccharides like starch and pectic substances into simpler substances during processing increases in TSS. Gillani [18] investigated increase in TSS in different mango cultivar. With the use of chemical preservative TSS of apple pulp increases Kinsh et al. [19]. It is concluded that TSS of tamarind plum blended squash increased with storage and treatment.

Ascorbic acid (Vitamin C)

The data from Table 3 shows significant ($p < 0.05$) effect on

Treatment	Storage interval							% Inc	Means
	Initial day	15	30	45	60	75	90		
TPS ₀	47.55	47.69	47.78	47.88	47.92	48.12	48.29	1.53	47.89g
TPS ₁	47.65	47.74	47.84	47.96	48.11	48.23	48.33	1.41	47.98f
TPS ₂	47.45	47.54	47.65	47.74	47.84	47.93	48.03	1.76	47.78h
TPS ₃	48.35	48.44	48.56	48.63	48.72	48.81	48.92	1.17	48.63e
TPS ₄	49.25	49.34	49.43	49.52	49.62	49.73	49.82	1.14	49.53d
TPS ₅	49.95	50.04	50.11	50.2	50.29	50.39	50.48	1.05	50.21c
TPS ₆	50.25	50.34	50.43	50.54	50.62	50.73	50.81	1.10	50.53b
TPS ₇	51.35	51.45	51.54	51.63	51.74	51.85	51.94	1.14	51.64a
Mean	48.98g	49.07f	49.17e	49.26d	49.36c	49.47b	49.61a	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 2: TSS (°brix) of squash prepared from blending of tamarind and plum juice at different levels.

Treatments	Storage interval							% Dec	Means
	Initial day	15	30	45	60	75	90		
TPS ₀	35.79	33.35	31.09	29.78	27.87	25.35	23.76	33.61	29.69g
TPS ₁	36.23	35.98	33.98	31.09	26.98	25.18	23.93	33.95	30.60f
TPS ₂	37.98	36.01	34.98	31.35	28.64	26.98	25.75	32.20	31.68e
TPS ₃	38.09	37.98	33.09	32.29	30.01	28.87	26.75	31.23	32.68d
TPS ₄	39.91	37.45	35.67	33.91	31.25	29.65	27.85	30.22	33.67c
TPS ₅	41.25	39.28	38.01	36.61	34.44	32.11	30.01	27.03	35.98b
TPS ₆	41.99	39.24	37.35	36.12	33.42	31.81	29.19	30.48	35.59b
TPS ₇	43.86	41.23	39.09	37.54	36.15	34.54	31.87	27.34	37.87a
Mean	39.49a	37.58b	35.72c	33.69d	31.11e	29.31f	27.40g	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 3: Ascorbic acid of squash prepared from blending of tamarind and plum juice at different levels.

Treatments	Storage interval							% Dec	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	1.00	0.98	0.96	0.95	0.93	0.92	0.90	10.00	2.08h
TPS ₁	1.03	1.01	0.99	0.97	0.95	0.93	0.91	11.65	2.31g
TPS ₂	1.05	1.03	1.01	0.99	0.97	0.95	0.93	11.43	2.29f
TPS ₃	1.08	1.06	1.04	1.03	1.01	0.99	0.97	10.19	2.17e
TPS ₄	1.11	1.09	1.07	1.05	1.03	1.02	1.00	9.91	2.16d
TPS ₅	1.15	1.13	1.11	1.09	1.08	1.06	1.04	9.57	2.15b
TPS ₆	1.13	1.11	1.09	1.07	1.05	1.04	1.02	9.73	2.16c
TPS ₇	1.18	1.16	1.15	1.13	1.11	1.08	1.06	10.17	2.25a
Mean	1.09a	1.07b	1.05c	1.04d	1.02e	1.00f	0.98g	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 4: Titratable acidity of squash prepared from blending of tamarind and plum juice at different levels.

storage and treatment of blended squash of tamarind plum. There shows a significant ($p < 0.05$) decrease in vitamin C. The ascorbic acid of tamarind plum squash was in the zero days from 35.79 (TPS₀) to 43.86 (TPS₇) which is then gradually decrease from 23.76 (TPS₀) to 31.87 (TPS₀) during storage period of 90 days. Mean value of ascorbic acid was recorded 39.49 at zero-day interval, while 27.40 at for the period of 90 days. According to Table 3, TPS₇ had a highest mean value (37.87), while TPS₀ had minimum (29.69) mean value. Sample TPS₁ (33.95) shows highest percent decrease, while sample TPS₅ (27.03) has lowest. The above results are in agreement with Kinh et al. [19], studied lower percent of ascorbic acid found in apple pulp affected by both temperature and light. Saleem et al. [20] studied that time interval also decreases ascorbic acid value. Bezman et al. [21] also concluded that ascorbic acid of grape juice also decreased during time of storage in room temperature. Storage interval, oxygen, light and heat treatment decrease the effect of ascorbic acid by both enzymatic and non-enzymatic catalyst [22]. In most liable nutrients, Vitamin C is very important because its degradation is used as an indicator of quality.

Titratable acidity

In Table 4 samples of tamarind plum squash shows significant (P<0.05) difference during time period of storage. The % acidity of the squash samples was in the range of 1 (TPS₀) to 1.18 (TPS₇) at initial day, while showed a decreasing trend of 0.9 (TPS₀) to 1.06 (TPS₇) correspondingly during 90 days of interval. Mean value at initial day was 1.09, decreases to 0.98 at 90 days intervals. The sample TPS₁ (2.31) shows high value of mean while sample TPS₅ (2.15) shows minimum. TPS₁ (11.65) had a maximum % decrease in acidity, while TPS₅ (9.57) showed the minimum decrease in percent acidity. Increase of acidity is because of storage condition and pectic substance break down [23]. Hye et al. [24] found increasing trend in acidity, while pH decrease of fruit juices during processing and storage time. Analogous result was reported by Gajanana [25] that hydrolysis of polysaccharides and non-reducing sugars reduces acid of amla juice, where the acid is converting to hexose sugars or complexes in the presence of metal ions. Lakshmi et al. [26] and Nidhi et al. [27] also observed reduction in acidity during the storage period of the tamarind RTS and RTS bael-guava beverages

Treatment	Storage interval							% Inc	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	47.55	48.66	49.77	50.40	51.53	52.30	53.66	11.38	50.55a
TPS ₁	46.26	47.27	48.32	49.44	50.64	51.86	53.11	12.89	49.56b
TPS ₂	45.19	46.16	47.18	48.22	49.32	50.45	51.94	12.99	48.35c
TPS ₃	44.77	45.70	46.69	47.21	48.24	49.30	50.43	11.23	47.48d
TPS ₄	44.37	45.27	46.20	47.16	48.17	48.75	49.82	10.94	47.11e
TPS ₅	43.70	44.55	45.43	46.37	46.87	47.86	48.86	10.56	46.23f
TPS ₆	44.20	45.08	45.97	46.92	47.90	48.45	49.49	10.68	46.86e
TPS ₇	43.52	44.35	44.82	45.69	46.61	48.01	49.00	11.19	46.00f
Mean	44.94g	45.88f	46.80e	47.68d	48.66c	49.62b	50.79a	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 5: Sugar acid ratio of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Means
	Initial day	15	30	45	60	75	90		
TPS ₀	2.68	2.69	2.71	2.72	2.73	2.74	2.76	2.90	2.72h
TPS ₁	2.69	2.07	2.72	2.73	2.75	2.76	2.77	2.89	2.73g
TPS ₂	2.72	2.73	2.74	2.75	2.76	2.78	2.79	2.51	2.75f
TPS ₃	2.72	2.74	2.75	2.76	2.78	2.79	2.08	2.86	2.76e
TPS ₄	2.75	2.76	2.77	2.79	2.81	2.82	2.83	2.83	2.79d
TPS ₅	2.86	2.88	2.89	2.91	2.92	2.92	2.93	2.39	2.90b
TPS ₆	2.81	2.83	2.84	2.85	2.86	2.87	2.88	2.43	2.85c
TPS ₇	2.89	2.9	2.92	2.93	2.94	2.95	2.97	2.69	2.93a
Mean	2.77g	2.78f	2.79e	2.81d	2.82c	2.83b	2.84a	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 6: pH of squash prepared from blending of tamarind and plum juice at different levels.

respectively. It's released from the data that the titratable acidity decreases with storage and treatment.

Sugar acid ratio

Table 5 shows a significant (P<0.05) effect on both the treatment effect and storage effect on blended squash of tamarind plum. The ratio sugar acid of squash samples was in the range of 43.52 (TPS₇) to 47.55 (TPS₀) at initial day, while showed an increasing trend of 48.86 (TPS₆) to 53.66 (TPS₁) correspondingly during 90 days storage interval. Initial day storage mean was 44.94, which increase to 50.79. Sample TPS₀ (50.55) show high mean value while the sample TPS₇ (46.00) with lowest value of mean. Sample TPS₀ showed % increase of maximum (11.38), while TPS₅ (10.56) showed the minimum increase in percent sugar acid ratio. According to Chyau et al. [28] substances like pectin, reducing sugar, total sugar and acidity of guava fruit decreases at ripe stage while the sugar/acid ratio of the fruit guava increased. It is concluded from the data that sugar acid ratio increased with time by storage and treatment.

pH

The data of tamarind plum blended squash shows a decreasing trend during the period of storage intervals. Tamarind plum squash pH was in between 2.68 (TPS₀) to 2.89 (TPS₇) at zero days of interval which gradually increases from 2.76 (TPS₀) to 2.97 (TPS₇) during 90 days of storage time. Mean of the data at 1st day was 2.77, and then decreased to 3.73 during keeping time of storage. Sample TPS₇ of tamarind plum squash shows high mean 2.93, while sample TPS₀ of tamarind plum has a lowest mean 2.72. Sample TPS₀ has high percent decrease of (2.90) in case of pH. However, squash sample TPS₅ (2.39) of the minimum pH with percent decrease found. There found a significant (P<0.05) effect of tamarind plum blended squash in case of time and treatment. Nath et al. [17] investigate same results for kinnow (mandarin) ginger

squash. According to Jitareerat et al. [29] pH of fruits and vegetables changes because of heat treatment on biochemical substances, decrease of respiration and metabolic process. Cecilia and Maia [30] studied a decreasing trend in pH of apple juice during keeping time. With the increase of acidity and pectin hydrolysis pH of the juice decline [31]. Thus, concluded that pH increases with treatment and storage effects on tamarind plum blended squash.

Reducing sugar

Table 6 shows effect of time interval and treatment on blended squash of tamarind plum. Reducing sugars of tamarind plum squash was in between 17.10 (TPS₇) to 17.32 (TPS₁) at initial day. There shows an increasing trend of 28.10 (TPS₀) to 33.23 (TPS₃) during 90 days of storage time period. Initial day mean of tamarind plum was 17.21, which shows gradual increase of 31.23 during the storage time period. Tamarind plum sample TPS₀ (25.32) showed the maximum mean value, however sample TPS₅ (23.08) had minimum mean value. Squash sample TPS₃ (48.18) found with maximum percent increase, while the sample TPS₅ (39.07) with lowest percent increase in reducing sugar. There found a significant (P<0.05) effect on tamarind plum blended squash during treatments and storage intervals of times. The above results show similarity with the report of Kotecha and Kadam [16] and Sahu et al. [32] on tamarind syrup and mango lemongrass beverage respectively reported an increase trend in total and reducing sugars. Both acidity and temperature has caused positive effect on reducing sugar (convert sucrose to glucose and fructose) [33]. Reducing sugar of fruits increases because of sucrose reduction. It is concluded that the reducing sugars of the treatment increases with time interval.

Non-reducing sugar

In the Table 7 tamarind plum blended squash data are significantly (P<0.05) reduced during storage and treatment intervals. The data of

squash samples was in the range of 40.20 (TPS₀) to 47.1 (TPS₄) at initial day. While during storage period the non-reducing sugar content decrease gradually from 19.35 (TPS₀) to 25.76 (TPS₃) at 90 days of interval. Initial mean data was 44.36, which shows a reducing trend of 21.97. Tamarind plum blended sample TPS₄ (37.64) with maximum mean, while squash sample TPS₇ (30.87) with minimum mean. High percent decrease (54.95), for sample TPS₂. However, TPS₅ (44.54) showed the minimum % decrease. Kotecha and Kadam [17] and Sahu et al. [32] reported same results of increasing total sugar as well as reducing sugar, while decreasing of non-reducing sugar for tamarind syrup and mango lemongrass beverage respectively, during storage. Main cause of reducing sugar conversion to non-reducing sugar is glycogenesis, also change of vitamins, sugar and organic acid change during storage intervals in carrot pulp. Thus, concluded that the non-reducing sugar decreases with treatment and storage condition.

Taste

According to the data of Table 8, statistically shows a reducing trend significantly (P<0.05) of tamarind plum blended squash during

treatment and storage condition. The sensory score for taste of tamarind plum blended squash were in the range of 6.6 (TPS₀, TPS₇) to 7.4 (TPS₃) at zero days of interval, there found a gradual decrease of 5.5 (TPS₁, TPS₂) to 6.5 (TPS₃) during the storage period of 90 days. Mean data for initial day was 6.85, which gradually down to 5.83. The squash of tamarind plum sample TPS₅ (7.00) shows highest mean, while with lowest score of sample TPS₇ (6.04). Sample (TPS₁) with maximum decrease of 17.91%, while minimum decrease of 12.16% was observed by TPS₅. The data above had a significant effect on taste of tamarind plum blended squash during storage and treatment time intervals. During RTS beverages light effects acids and ascorbic acid (Vitamin C) present in orange squashes [34]. The RTS of tamarind shows same results according to Kotecha and Kadam [17]. The depletion of taste is effected by acid, pH fluctuation [35].

Color

There shows a decreasing effect significantly (P<0.05) on color of tamarind plum squash during period of time interval. At zero day interval, the sensory score for color of tamarind plum squash samples

Treatment	Storage interval							% Inc	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	17.25	20.25	23.76	25.48	28.15	30.35	31.98	46.06	25.32a
TPS ₁	17.32	19.01	21.24	25.31	28.54	30.35	32.31	46.39	24.88ab
TPS ₂	17.28	18.09	21.29	24.32	26.45	28.54	31.46	45.07	24.03bc
TPS ₃	17.22	20.02	22.25	25.32	28.45	30.21	33.23	48.18	25.27a
TPS ₄	17.24	19.25	22.87	24.54	26.93	28.54	30.65	43.75	24.29bc
TPS ₅	17.12	20.21	22.23	23.35	24.43	26.15	28.01	39.07	23.08d
TPS ₆	17.15	19.01	22.98	24.84	26.89	29.45	31.98	46.37	24.63abc
TPS ₇	17.01	19.19	21.09	23.65	26.98	28.09	30.15	43.28	23.98c
Mean	17.21g	19.53f	22.32e	24.60d	27.10c	29.06b	31.23a	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 7: Reducing sugar of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	40.02	39.56	35.43	31.87	27.85	22.96	19.35	51.87	31.03d
TPS ₁	42.35	40.12	35.24	31.01	27.43	24.45	20.25	52.18	31.55d
TPS ₂	44.95	42.01	38.65	34.21	30.24	23.46	20.25	54.95	33.41c
TPS ₃	45.98	43.25	40.13	35.65	31.35	25.78	21.98	52.20	34.87b
TPS ₄	47.01	45.24	41.21	39.19	34.99	30.01	25.65	45.54	37.64a
TPS ₅	46.45	43.87	40.24	35.87	30.87	28.31	25.76	44.54	35.91b
TPS ₆	46.01	45.01	40.95	37.45	30.14	25.09	22.45	51.30	35.44b
TPS ₇	41.76	39.35	34.76	30.12	27.09	22.12	20.01	51.87	30.87d
Mean	44.36a	42.32b	38.33c	34.42d	30.10e	25.39f	21.97g	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 8: Non-reducing sugar of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	6.6	6.4	6.3	6.2	6.0	5.8	5.7	13.64	6.14de
TPS ₁	6.7	6.6	6.4	6.1	6.0	5.8	5.5	17.91	6.16cd
TPS ₂	6.8	6.6	6.5	6.2	6.0	5.9	5.7	16.18	6.24cd
TPS ₃	6.9	6.7	6.5	6.4	6.3	6.2	6.0	13.04	6.43b
TPS ₄	7.0	6.9	6.7	6.5	6.3	6.1	6.0	14.29	6.50b
TPS ₅	7.4	7.3	7.2	7.0	6.9	6.7	6.5	12.16	7.00a
TPS ₆	6.8	6.6	6.5	6.2	6.1	5.9	5.7	16.18	6.26c
TPS ₇	6.6	6.3	6.2	6.0	5.9	5.8	5.5	16.67	6.04e
Mean	6.85a	6.68b	6.54c	6.33d	6.19e	6.03f	5.83g	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 9: Taste of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Mean
	0	15	30	45	60	75	90		
TPS ₀	6.5	6.4	6.2	6.1	5.9	5.6	5.5	15.38	6.03b
TPS ₁	6.3	6.2	6.0	5.9	5.6	5.5	5.3	15.87	5.83c
TPS ₂	6.1	6.0	5.8	5.6	5.5	5.3	5.1	16.39	5.63e
TPS ₃	6.0	5.9	5.7	5.4	5.3	5.2	5.0	16.67	5.50f
TPS ₄	6.2	6.0	5.9	5.8	5.7	5.4	5.3	14.52	5.76d
TPS ₅	7.1	7.0	6.9	6.7	6.6	6.4	6.2	12.68	6.70a
TPS ₆	6.2	6.0	5.9	5.6	5.4	5.3	5.2	16.13	5.66e
TPS ₇	6.2	6.0	5.9	5.7	5.6	5.5	5.3	14.52	5.74d
Mean	6.33a	6.19b	6.04c	5.85d	5.70e	5.53f	5.36g	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 10: Color of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	7.1	6.6	6.0	5.4	4.4	3.3	2.3	67.61	5.01c
TPS ₁	7.8	7.5	7.1	6.7	6.2	5.6	4.9	37.18	6.54b
TPS ₂	7.6	7.5	7.3	7.2	7.0	6.6	6.3	17.11	7.07ab
TPS ₃	7.5	7.4	7.2	7.0	6.9	6.8	6.7	10.67	7.07ab
TPS ₄	7.7	7.4	7.2	6.8	6.6	6.3	6.0	22.08	6.86b
TPS ₅	7.9	7.8	7.7	7.6	7.4	7.3	7.1	10.13	7.54a
TPS ₆	7.4	7.3	7.1	7.0	6.8	6.7	6.3	14.86	6.94b
TPS ₇	7.3	7.2	7.1	7.0	6.9	6.8	6.4	12.33	6.96b
Mean	7.54a	7.34ab	7.09ab	6.84bc	6.53cd	6.18de	5.75e	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 11: Flavor of squash prepared from blending of tamarind and plum juice at different levels.

Treatment	Storage interval							% Dec	Mean
	Initial day	15	30	45	60	75	90		
TPS ₀	7.8	7.3	6.8	6.1	5.3	4.1	3.6	53.85	5.86d
TPS ₁	8.0	7.7	7.4	7.0	6.4	5.8	5.0	37.5	6.76c
TPS ₂	8.2	7.9	7.5	7.1	6.6	6.0	5.4	34.15	6.96bc
TPS ₃	8.0	7.8	7.6	7.5	7.4	7.1	6.9	13.75	7.47ab
TPS ₄	8.1	8.0	7.8	7.5	7.4	7.1	7.0	13.58	7.56a
TPS ₅	8.2	8.0	7.9	7.8	7.7	7.4	7.3	10.98	7.76a
TPS ₆	8.0	7.9	7.5	7.4	7.3	7.1	6.9	13.75	7.44ab
TPS ₇	7.9	7.7	7.6	7.5	7.3	7.1	7.0	11.39	7.44ab
Mean	8.03a	7.79ab	7.51bc	7.24cd	6.93de	6.46ef	6.14f	--	--

^{a-g} Values of different alphabetic letter shows significant (P<0.05) difference from each other.

Table 12: Overall acceptability of squash prepared from blending of tamarind and plum juice at different levels.

from 6 (TPS₃) to 7.1 (TPS₅) which decreased gradually from 5 (TPS₅) to 6.3 (TPS₅) through 90 days of intervals. Initial day mean was 6.33, which decreases to 5.36. The sample TPS₅ with maximum mean of 6.70 were found, while there found lowest mean of 5.50 for sample TPS₃. Decrease of 16.67 % was observed at sample TPS₃ while the minimum % decrease was noted at TPS₅ (12.68). There found a significant (P<0.05) effect on color of tamarind plum blended squash during storage interval. The result was in favor of Jain et al. [36], reported a decreasing trend in color during 90 days storage of squash. Color of the beverages decreases because of presence of 2 Methyl 3 furanthiol and methanol gives rotten flavors in stored orange juices [21]. Brennder et al. [37] studied that presence of SO₂ decreases fruits and vegetables browning.

Flavor

Table 9 shows data of tamarind plum blended squash. The mean sensory scores for flavor of squash decreased significantly (P<0.05) on both treatments and storage time intervals. The judges panel scores for flavor of tamarind plum blended squash from 7.1 (TPS₀) to 7.9 (TPS₅) during zero days of intervals. However, during storage interval of 90

days' flavor of the squash samples decreased gradually from 2.3 (TPS₀) to 7.1 (TPS₅). Mean flavor was found 7.54, which decreased to 5.75 throughout the storage period of time intervals. TPS₅ was found to be high mean (7.54), while the low score mean (5.01) was obtained for TPS₀. The maximum percent decrease in flavor of the squash was recorded in TPS₀ (67.61), while minimum decrease of 10.13% was observed at TPS₅. The tamarind plum squash was significantly (P<0.05) differ in case of treatment and time interval. Results of physicochemical, sensory properties of orange drink shows similarity were reported by Jain et al. [35]. According to Martin [38] results on pasteurized orange juice shows depletion of organoleptic quality kept in glass bottles. Similar with these results of Paracha [39], that loss of flavor of guava squashes during storage of 3 months of storage interval. A slight difference in flavor may be due to storage conditions and storage time.

Overall acceptability

Table 10 shows the effect of both the treatments and storage interval on overall quality of tamarind plum blended squash. The acceptability of overall quality of the blended squash reduces

considerably ($P < 0.05$) on both treatments and storage time interval. The overall acceptance score of tamarind plum squash at initial days ranges from 7.8 (TPS₀) to 8.2 (TPS₅, TPS₂), which fall gradually from 3.6 (TPS₀) to 7.3 (TPS₅) during the 90 days of storage period of time. Mean value for over-all acceptance was 8.03, which decrease down to 6.14 during the storage period. The highest score of mean (7.76) was observed at TPS₅, while minimum score of mean (5.86) was observed at TPS₀. The highest percent decrease of 53.85 was recorded at TPS₀, while minimum percent decrease of 10.98 was observed at TPS₅. The overall acceptability of tamarind plum blended squash is significantly ($P < 0.05$) influenced by treatments and storage interval (Tables 11 and 12). Rosario [34] observed that with the increasing of days' storage overall quality of acceptance decreases. Loss of overall quality were affected by processing like, temperature and storage time [24].

Conclusion and Recommendations

Conclusion

Present work of tamarind plum blended squash was carried out with different proportions. Chemical preservatives were used to inhibit the growth of microbial activity in tamarind plum blended squash. Prepared squash was packed in plastic bottles and stored at room temperature for 90 days of storage. Prepared squash was then evaluated for physicochemical and sensory properties during 90 days of storage. Some physicochemical and sensory analysis was examined to be changed but not affected overall quality of the squash. On the basis of above results it was concluded that sample TPS₅ show best in keeping quality during storage time intervals. Hence, the results of sample TPS₅ of tamarind plum blended squash is more recommended in terms of commercial use and for large scale industrial production. Squash prepared from tamarind and plum are more acceptable to consumers because of sour test, need commercialization.

Recommendations

1. Different proportion of tamarind pulp can also be used with other fruit pulp.
2. It is suggested to study the influence of storage condition and packaging materials on tamarind plum blended squash.
3. This is recommended to carry a research on non-caloric tamarind plum blended squash.

References

1. Komutarin T, Azadi S, Butterworth L, Keil D, Chitsomboon B, et al. (2004) Extract of the seed coat of *Tamarindus indica* inhibits nitric oxide production by murine macrophages *in vitro* and *in vivo*. Food Chem Toxicol 42: 649-658.
2. Siddiq KE, Gunasena HP, Prasad BA, Pushpakumar DK, Ramana KV, et al. (2006) Tamarind monograph. Southampton centre for underutilized crops, Southampton, UK pp: 1-198.
3. Martinello F, Soares SM, Franco JJ, Santos AJ, Sugohara A, et al. (2006) Hypolipemic and antioxidant activities from *Tamarindus indica* L. Pulp fruit extract in hypercholesterolemic hamsters. Food Chem Toxicol 44: 810-818.
4. Maiti R, Das UK, Ghosh D (2005) Attenuation of hyperglycemia and hyperlipidemia in streptozotocin-induced diabetic rats by aqueous extract of seed of *Tamarindus indica*. Bio Pharm Bulletin 28: 1172-1176.
5. Joshi AA, Kshirsagar RB, Sawate AR (2012) Studies on standardization of enzyme concentration and process for extraction of tamarind pulp, variety Ajanta. J Food Process Technol 3: 1-3.
6. Shankaracharya NB (1998) Tamarind chemistry, technology and uses: A critical appraisal. J Food Sci Technol 35: 193-208.
7. Sanchez PC (1985) Tropical fruit wines. A lucrative business Research at Los Banos 3: 10-13.
8. Wang CY (1993) Approaches to reduce chilling injury of fruits and vegetables. Hort Rev 15: 63-95.
9. Diaz MHM, Zapata PJ, Guillen F, Romero DM, Castillo S (2009) Changes in hydrophilic and lipophilic antioxidant activity and related bioactive compounds during postharvest storage of yellow and purple plum cultivars. Postharvest Bio Technol 51: 354-363.
10. Seeram NP, Adams LS, Zhang YJ, Lee R, Sand D (2006) Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells *in vitro*. J Agri Food Chem 54: 9329-9339.
11. Ismail S, Rehman S (1995) Beverages: Science and Technology.
12. Hicks D (1990) Production and packaging of non-carbonated fruit juices and fruit beverages. Van Nostrand Reinhold, New York.
13. Archana P, Laxman K (2014) Studies on preparation and storage of tamarind squash. J Spice Aromatic Crop 24: 254-261.
14. AOAC (2012) Official method of analysis. (19th edn.) The Association for Official Analysis in Chemistry, Rockville, USA.
15. Larmand E (1977) Laboratory method for sensory evaluation of food. Pub. Canada Department of Agriculture, Ottawa.
16. Kotecha PM, Kadam SS (2003) Preparation of ready to serve beverage, syrup and concentrate from tamarind. J Food Sci Tech 40: 76-79.
17. Nath A, Yadav DS, Sarma P, Dey B (2005) Standardization of ginger-kinnow squash and its storage. J Food Sci Tech 42: 520-522.
18. Gillani SSN (2002) Development of mango squash from four different cultivars of mango. Department of Food Science and Technology, NWFP Agricultural University, Peshawar.
19. Kinh AE, Shearer CP, Dunne S, Hoover DG (2001) Preparation and preservation of apple pulp with chemical preservatives and mild heat. J Food Process 28: 111-114.
20. Saleem N, Kamran M, Shaikh SA, Tarar OM, Jamil K (2011) Studies on processing and preparation of peach squash. Pak J Biochem Mol Biol 44: 12-17.
21. Bezman Y, Russell L, Rouseff D, Naim M (2001) 2-Methyl-3-furanthiol and methional are possible off-flavors in stored orange juice. J Agric Food Chem 49: 425-432.
22. Mapson LW (1970) Vitamins in fruits. In: Hulme AC (Ed.) The Biochemistry of Fruits and their Products. Academic Press, London pp: 369-384.
23. Hashmi MS, Alam S, Riaz A, Shah AS (2007) Studies on microbial and sensory quality of mango pulp storage with chemical preservatives. Pak J Nutr 6: 85-88.
24. Hye WY, Streaker CB, Zhang QH, Min DB (2000) Effect of pasteurized electric field on the quality of orange juice and comparison with heat pasteurization. J Agri Fd Chem 48: 4597-4605.
25. Gajanana K (2002) Processing of aonla (*Emblica officinalis Gaertn.*) fruits. University of Agriculture Sciences, Dharwad, India.
26. Lakshmi K, Kumar KAKV, Rao LJ, Naidu MM (2005) Quality evaluation of flavored RTS beverage and beverage concentrate. J Food Sci Tech 42: 411-414.
27. Nidhi R, Gehlot R, Singh, Rana MK (2008) Changes in chemical components of RTS bael-guava blended beverages during storage. J Food Sci Tech 45: 378-380.
28. Chayu CC, Wu SY, Chen CM (1992) Differences of volatile and nonvolatile constituents between mature and ripe guava fruit. J Agric and Food Chem 40: 846-849.
29. Jitareerat P, Paumchai S, Kanlayanarat S (2007) Effect of chitosan on ripening enzymatic activity and disease development in papaya (*Carica papaya*) fruit. New Zealand J Crop Hort Sci 35: 211-218.
30. Cecilia E, Maia GA (2002) Storage stability of cashew apple juice preserved by hot fill and aseptic process. University of Ceara, Brazil.
31. Imran A, Rafiullah K, Muhammad A (2000) Effect of added sugar at various concentration on storage stability of guava pulp. Sarhad J of Agric 7: 35-39.
32. Sahu C, Choudhary PL, Patel L, Sahu R (2006) Physico-chemical and sensory characteristics of whey based mango herbal (lemon grass) beverage. Ind Food Packer 60: 127-132.

33. Singh S, Shivhare US, Ahmed J, Raghavan GSV (1999) Osmotic concentration kinetics and quality of carrot preserve. J Food Res Int 32 : 509-514.
34. Muhammad R, Ahmed M, Chaudhry MA, Hussain B, Khan I (1987) Ascorbic acid quality retention in orange squashes as related to exposure to light and container type. J Pak Sci Ind Res 30: 480-483.
35. Rosario MJG (1996) Formulation of ready to drink blends from fruits and vegetables juices. J Philippines 9: 201-209.
36. Jain S, Sankhla APK, Dashora A, Sankhla AK (2003) Physiochemical and sensory properties of orange drink. J Food Sci Tech Ind 40: 656-659.
37. Brenndor K, Oswin CO, Trim DS, Mrema GC, Werek GC (1985) Solar driers and their role in post-harvest processing. Common wealth Sci council 2: 78-83.
38. Martin JJ, Solances E, Bota E, Sancho J (1995) Chemical and organoleptic changes in pasteurized orange juice. Alimentaria 216: 59-63.
39. Paracha GM (2004) Development and storage stability of low caloric guava squash. Agricultural University, Peshawar, Pakistan.

Citation: Khan I, Rehman AU, Khan SH, Qazi IM, Khan A, et al. (2017) Development and Quality Evaluation of Tamarind Plum Blended Squash During Storage. J Food Process Technol 8: 662. doi: [10.4172/2157-7110.1000662](https://doi.org/10.4172/2157-7110.1000662)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid peer review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled for better prominence and citations
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Best discounts for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>