



Development and Validation of an Instrument to Enhance the Community Pharmacy Practitioner's Knowledge towards Handling of High Risk/Alert Medications

Balakeshwa Ramaiah*, Sejal Sharma, Saroj Poudel and Raju Koneri

Department of Pharmacy Practice, Karnataka College of Pharmacy, Bengaluru, India

Abstract

Aim: Medication errors may not occur more often with high-risk drugs, but the consequences or impacts from them could be more dangerous for the patients. This study was aimed to prepare and validate an instrument as learning modules to boost community pharmacists' knowledge on High-risk Medications (HRM), which will help in minimizing serious consequences arising due to mishandling of HRM.

Methods: The instrument (videos) included chapters "introduction to HRM," "Look Alike and Sound Alike (LASA) drugs" and "storage and labeling of HRM." The instrument was ensured to be imperative, relevant, reactive and appropriate with the help of content and face validation which was then confirmed to be sensitive enough to distinguish knowledge levels of community pharmacists. The split-half reliability test by Kuder-Richardson formula 20 (KR 20) to obtain a homogenous reliability index value ($r_{KR20} = (k / k - 1) / (1 - \sum pq / \sigma^2)$) ensured internal consistency of the instrument.

Results: The five-points Likert scale showed an average score of above four points with Content Validity Index (CVI) for I-CVI as 0.913 and for S-CVI as 0.916. This indicated appropriateness, conciseness, and importance of the training materials. During face validation, the approval of the design of the learning modules was done, and the importance of the issues to the community pharmacy profession was thus emphasized. The KR 20 index values homogeneously reached 0.937 for introduction, 0.8424 for LASA and 0.8195 for storage and labelling chapters, suggesting that the learning modules were reliable, operational, feasible and attractive.

Conclusion: The validated instrument considered as a vital tool to improve the community pharmacist's knowledge and handling of HRM (s), thus contributing to better patient care.

Keywords: High-risk medications; Community pharmacists; Learning modules; Content and face validation; Reliability

Introduction

Medications have a significant role in diseases management and their prevention. Drugs are manufactured and marketed with the potential of a wide range of safety. However, a rare class or group of medicines, called as High-Risk or High-Alert Medications (HRM), are known to have a threat in causing significant patient harm, disability or death if they are unintentionally misused or improperly administered. The term "high-risk" medications were coined initially by the Institute for Safe Medication Practices (ISMP) in 1998 for those drugs which are linked or related to most dangerous Preventable Adverse Drug Events (PADEs). Medication errors may not occur more often with high-risk drugs, but the consequences or impacts from them could be more dangerous for the patients. Therefore, various risks or hampers that could take place while prescribing, storing, dispensing, and finally administering a high-risk drug should be supervised carefully at each phase of the medication management process [1-3].

According to the American Pharmaceutical Association, eight categories were listed as high-risk medications that include high concentration electrolytes, chemotherapeutic agents, opiates, anticoagulants, narcotics, neuromuscular blocking agents, benzodiazepines and cardiovascular drugs. The process of drug dispensing or administration to patients at a hospital involves multifarious phases that in turn is based on a series of inter related actions and decisions overcoming daily obstacles. Nonetheless, this management process may not be satisfactorily safe every time, due to which the faults arising may or may not cause damage to the patient. These faults or mistakes, typically said as medication errors, arising in the administration pathway can be considered as preventable

adverse events [4]. In 2003, ISMP performed a study for assessment of knowledge on high risk medications for distinguishing variances between pharmacy and nursing perspectives, most of the participants responded their agreement on which medications were considered high risk. This survey was repeated by ISMP in 2007 and 2012. In all the three surveys, it was noted that the pharmacists were not able to identify medications as high risk, as often as nurses did.

It is projected that a hospitalized patient is identified to be exposed to at least one error per day related to drug. According to ISMP, an estimate of as low as 450,000 medication errors result in injury to patients in the United States per year, with around 25% of these errors meant to be avoidable. In addition, 7,000 deaths each year are recognized to be because of medication errors. In the field of community pharmacy, a few studies have been found to report the occurrence of injury to the patient caused by medication errors that are preventable. Ghandi et al. [5] mentioned that adverse events that were preventable occurred in 5% of ambulatory patients with medications that were dispensed from

***Corresponding author:** Balakeshwa Ramaiah, Associate Professor, Department of Pharmacy Practice, Karnataka College of Pharmacy, Bengaluru, India, Tel: +919845516090; E-mail: balupharmacy@gmail.com

Received: February 21, 2019; **Accepted:** March 07, 2019; **Published:** March 17, 2019

Citation: Ramaiah B, Sharma S, Poudel S, Koneri R (2019) Development and Validation of an Instrument to Enhance the Community Pharmacy Practitioner's Knowledge towards Handling of High Risk/Alert Medications. J Biomed Pharm Sci 2: 118

Copyright: © 2019 Ramaiah B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

community pharmacies. Also, Gurwitz et al. [6] identified that one-half of life-threatening, serious or fatal adverse drug events resulted from medications dispensed from pharmacies that were preventable.

The study was thus aimed to prepare and develop instruments prior in labelling, handling, storage and dispensing of HRM for the pharmacists who would be further implemented with the important process of validation and reliability. As documented in several studies, validation has always been an important factor as the measurement of accuracy and consistency in research instruments. However, in various health and social science research taking place in developing countries, validation of instruments is not being commonly performed. This has been linked to the shortage of information on how validation should be carried out to certain degree of conclusion. As per the review article from Nigerian researcher Bolarinwa [7], highlighted that the literary and technical meanings of instruments were both reflected by validation and reliability making them an important procedures to be done in research works. They elaborated numerous forms and methods of analyzing validation and reliability of an instrument, the main goal of which was to improve knowledge of these tests among young researchers in developing countries [8,9].

According to an international literature published by Sampaio and Sequeira [10], an instrument that was proven to be valid and reliable was developed for assessment of knowledge in nurses regarding HRM. The validity consisted of content, construct and face validity whereas the reliability of the instrument was measured through internal consistency using Kuder-Richardson reliability 20 (KR 20) formula. In the same manner, considering the importance of assessment of knowledge of HRM to the community pharmacist, in this research, the instrument as educative materials, has been developed and validated by deep and vigorous study from various experts. Finally, the same was done with measurement of reliability with KR 20 for its internal consistency. This study was aimed to prepare and validate an instrument as learning modules to boost community pharmacists' knowledge on High Risk Medications (HRM), which will help in minimizing serious consequences arising due to mishandling of HRM.

Methods

The study was a prospective interventional methodological program. This study attempted on methods of preparation of an instrument in the form of suitable educative video materials, following with organization and analysis of data collected for the main purpose of validation of the research instruments and techniques. The summarized study methodology is represented in Figure 1.

Preparation and development of an instrument

For collection of data, the setting of HRM management in the particular area was needed to be known. So, a visit was made to various pharmacies for the same. With the respect to Indian pharmacy practice environment, training materials were suitably prepared on the information from global guidelines and practice for HRM management. Three informative and revealing chapters on management of HRM were prepared from various sources and literatures. The materials were prepared in such a way that they become easily understandable and comprehensive. The materials were both accessible in high resolution PC formats and size compressed mobile formats such that the acceptability of the material by the participants would highly be favoured. It was




Figure 1: The overall study procedure along with reliability scores.

reaffirmed that the training materials would be helpful as a knowledge material in the Indian setup. The training material in the form of hard copy was validated (already accomplished) out of the objective of the study. After which, the hard copy materials were converted to scripts in the form of narrations for the purpose of recording it and preparing as convenient video materials. The language of the script was ascertained for easy understandability.

Development of an instrument in the form of video materials

A suitable female artist was chosen for the recording the script; considering factors like voice, tone, clarity of pronunciation, speech flow and finally delivery of the speech. The whole process was carried out in a studio environment under the supervision of a technical team having hands-on experience in recording and editing such videos. The processing of the video materials involved the following five crucial steps viz

- (i) Recording the scripts using appropriate voice software into individual sound tracks and then joining up the same into one single audio file.
- (ii) Collaborating the slides of power point presentations with their specific audio files to produce a video file.
- (iii) Adjustment of time intervals.
- (iv) Addition of suitable background tracks to the collaborated file.
- (v) Converting the file into high resolution PC format as well as in compressed mobile format. The software such as Audio Recorder by Green Apple Studio (Version 1.9.45), Audacity. The free, Cross-Platform Sound editor by Audacity Development Team (Version 2.1.3) and Corel Video Studio Ultimate X1 was utilized for the purpose of recording the scripts and collaborating it with the slides of the power point materials. The videos were finally ensured to be checked for synchronization and clarity.

Validation of the instrument

Various literature evidences implicated the importance of the

validation as a degree to which a measurement measures what it purports to measure. A validation technique can be either logical or rational. Validation illustrates the estimate of how much a measure or a dimension represents each and every single component of a hypothesis. The content validity (8 experts involving senior consultants and community pharmacist) and face validity (45 community pharmacy practitioners) were performed in this study. The prepared instrument (Learning modules) was ensured to be important, relevant, reactive and appropriate with the help of content and face validation and was then confirmed to be sensitive enough to distinguish knowledge levels of community pharmacists.

A total of 45 pharmacists (15 hospital pharmacists and 30 community pharmacists) were involved for the process of reliability. For the collection of data from the participant's responses regarding the training material, a form containing scoring columns for various aspects related to the material such as contents, clarity of the video and audio as well as various diagrammatic illustrations was distributed. The final scoring and feedbacks were evaluated and the appropriateness and reliability of the material was finally measured with KR 20.

Content validity: The percentage of agreement among specialists as to the instrument assessment and its item was obtained by means of the calculation of Content Validity Index (CVI). This index permits for the analysis of each item individually, and subsequently, the instrument as a whole. Lynn [11] through rigorous research illustrated that the researchers follow CVIs of two types based on the agreement of experts on the content of the instrument. It involves the Item-Content Validity Index (I-CVI) and Scale level-Content Validity Index (S-CVI). For I-CVI, the settlement among reviewers concerning each item of the instrument was measured by means of a Likert scale, with scores that range from score 1 to 4 (where, 1=irrelevant, 2=slightly relevant, 3=fairly relevant and 4=extremely relevant) Item that obtained scores of 1 or 2 were reviewed or eliminated. The calculation of the I CVI for each item consisted of the division between the numbers of answers that were fairly and extremely relevant by the total number of answers. The study also recommended an I-CVI > 0.78 for analyses of instrument by six or more judges.

The S-CVI involves the mean proportion of items rated as fairly and extremely relevant across various experts. This description of the CVI for scales was referred as S-CVI/average as for the purpose of convenience. This was interpreted as the combination of the number of items that were rated fairly and extremely relevant by all experts and to then which the total number of all the ratings is divided. It is also theorized that the S-CVI/ average is the mean or average I-CVI value

because it happens to concentrate on mean or average item quality rather than on average enactment by the experts [12,13]. Waltz and Strickland [14] stated that for mean congruity, the standard value to be considered is 0.90.

Face validity: Face validation consists of the subject experts observing thoroughly at the items in the instrument (learning modules) and approving that the test is a valid measure of the conception which is being evaluated just on the face of it. In simple words, they are assessing each aspect of the measuring items if they really match with the theoretical domain of the model. The approval of the design of the learning modules was strongly highlighted when face validation was performed and the importance of the issues to community pharmacy profession was thus emphasized as the experts understood all the components of the training material providing them with a secure atmosphere.

Reliability: The demand of reliability for measurement of internal consistency of a test is that it is needed to be estimated after only one test administration which therefore helps to escape the issues associated with testing over multiple time periods. By KR-20 formula, an index score for reliability was calculated as shown in the Formula 1.

$$r_{KR20} = \left(\frac{k}{k-1} \right) \left(1 - \frac{\sum pq}{\sigma^2} \right) \quad (1)$$

Where, r_{KR20} is the Kuder-Richardson formula 20; k is the total number of test items; \sum indicates to sum; p is the proportion of the test takers who pass an item; q is the proportion of test takers who fail an item; σ^2 is the variation of the entire test.

Results

As illustrated in Table 1, the mean I-CVI was figured out to be 0.913. Lynn [11] also suggested that when there is participation of five or fewer experts, there should be a universal agreement on the content validity for their rating to be said as an equitable representation. As per definition, the S-CVI/average is the combination of number of items rated either extremely or fairly relevant by all experts (Y), divided by the total number of all the possible ratings. Therefore, the S-CVI/average were found to be 0.916 (Table 1). The overall reliability for all three chapters is shown in Table 2. In this study the overall sample observations used for the reliability (n=45) was documented in excel sheet and documented the correct score per slide for the respective chapters as shown in Tables 3-5 respectively. The mean sum of a product of the proportion passed and proportion failed was calculated to apply standard deviation for each slide of all the three chapters,

Chapters	Expert 1	Expert 2	Expert 3	Expert 4	Expert 5	Expert 6	Expert 7	Expert 8	Number in Agreement	Item CVI (I- CVI)
Introduction to HRM	Y	Y	Y	Y	Y	Y	Y	N	7	0.87
LASA	N	Y	Y	Y	Y	Y	Y	Y	7	0.87
Storage and labelling	Y	Y	Y	Y	Y	Y	Y	Y	8	1.00
Mean I-CVI= 0.913 Mean S-CVI= 0.916										
*I-CVI: Item-level Content Validity Index; *S-CVI : Average scale- level content validity; *Y: Agreement on the content by the expert; *N: Disagreement on the content by the expert.										

Table 1: Content Validation by six experts using content validity index.

Chapter Name	Index	Reliability
Introduction to High risk medications	0.937*	Homogenous
Look-alike and sound-alike medications	0.8424*	Homogenous
Storage and labelling	0.8195*	Homogenous
*If the index value is >0.50 the sample is having good reliability		

Table 2: Representation of index value and reliability for the three learning video modules.

Slide Number	Correct Score	Proportion Passed	Proportion Failed	p*q
1	26	0.57	0.43	0.245
2	32	0.7	0.3	0.21
3	26	0.57	0.43	0.24
4	31	0.68	0.32	0.2176
5	15	0.33	0.77	0.2541
6	30	0.66	0.34	0.2244
7	21	0.46	0.54	0.2484
8	11	0.24	0.76	0.1824
9	23	0.51	0.49	0.2499
10	25	0.55	0.45	0.2475
11	27	0.6	0.4	0.24
12	30	0.66	0.34	0.2244
13	29	0.64	0.36	0.2304
14	36	0.8	0.2	0.16
15	15	0.33	0.77	0.2541
16	35	0.77	0.33	0.2541
17	30	0.66	0.44	0.4224
18	25	0.55	0.45	0.2475
19	28	0.62	0.38	0.2356
20	30	0.66	0.34	0.2244

Mean Sum of p*q: 5.0663; Standard deviation squared: 46.0648

$$\text{Index value } r_{KR20} = \left(\frac{k}{k-1} \right) \left(1 - \frac{\sum pq}{\sigma^2} \right) : 0.937$$

Table 3: Reliability table for the chapter "Introduction of HRM" by KR 20 Formula.

Slide Number	Correct Score	Proportion Passed	Proportion Failed	p*q
1	35	0.77	0.33	0.2541
2	32	0.71	0.29	0.2059
3	33	0.73	0.27	0.1971
4	34	0.75	0.25	0.1875
5	26	0.57	0.43	0.2451
6	29	0.64	0.36	0.2304
7	24	0.53	0.47	0.2491
8	35	0.77	0.33	0.2541
9	32	0.71	0.29	0.2059
10	33	0.73	0.27	0.1971
11	34	0.75	0.25	0.1875
12	28	0.62	0.38	0.2356
13	22	0.48	0.52	0.2496
14	29	0.64	0.36	0.2304
15	26	0.57	0.43	0.2451
16	29	0.64	0.36	0.2304
17	33	0.73	0.27	0.1971
18	19	0.42	0.58	0.2436
19	20	0.44	0.56	0.2464
20	22	0.48	0.52	0.2496
21	29	0.64	0.36	0.2304

Mean Sum of p*q: 4.8962; Standard deviation squared: 21.8146

$$\text{Index value } r_{KR20} = \left(\frac{k}{k-1} \right) \left(1 - \frac{\sum pq}{\sigma^2} \right) : 0.8424$$

Table 4: Reliability table for the chapter "Look-alike and sound-alike medications" by KR 20 Formula.

and finally, KR-20 reliability formulae were applied. The individual lessons assessed for their reliability, from which the index was obtained

Slide Number	Correct Score	Proportion Passed	Proportion Failed	p*q
1	27	0.6	0.4	0.24
2	25	0.55	0.45	0.2475
3	23	0.51	0.49	0.2499
4	30	0.66	0.34	0.2244
5	24	0.53	0.47	0.2491
6	28	0.62	0.38	0.2356
7	34	0.75	0.25	0.1875
8	36	0.8	0.2	0.16
9	33	0.73	0.27	0.1971
10	30	0.66	0.34	0.2244
11	30	0.66	0.34	0.2244
12	31	0.68	0.32	0.2176
13	34	0.75	0.25	0.1875
14	34	0.75	0.25	0.1875
15	33	0.73	0.27	0.1971
16	30	0.66	0.34	0.2304
17	30	0.66	0.34	0.2304
18	31	0.68	0.32	0.2172
19	24	0.53	0.47	0.2491
20	34	0.75	0.25	0.1875

Mean Sum of p*q: 4.3442; Standard deviation squared: 19.6144

$$\text{Index value } r_{KR20} = \left(\frac{k}{k-1} \right) \left(1 - \frac{\sum pq}{\sigma^2} \right) : 0.8195$$

Table 5: Reliability table for chapter "Storage and labeling" by KR 20 Formula.

as 0.937, 0.8424 and 0.8195 for chapter 1, chapter 2 and chapter 3 respectively.

Discussion

Content validity

The content validity index was reported only in methodological studies because it has focused merely on explaining the process of content validations.

Item-level content validity index (I-CVI)

I-CVI is used commonly by researchers to obtain information on guiding themselves in reviewing, erasing, or replacing items. However, the researchers do not generally provide information about I-CVI values in their reports, as I-CVIs are meant only to report in procedural research which mainly concentrates on clarifications of the overall content validity process. I-CVI expressed as the total number of experts giving a fair or extreme rating of either 3 or 4 (thus dichotomizing the standard scale into either relevant or not relevant), divided by the total number of experts. In simple words, the I-CVI should be closely 1.00 when there are five or fewer experts giving their ratings. The standard value could be a little relaxed when there are six or more raters, but I-CVIs usually must not show less than 0.78. For example, there could be one "not relevant" rating (I-CVI ¼, i.e., 0.83) with six raters and there could be two not relevant ratings with nine raters. Thus, the mean I-CVI which we obtained could measure as an ideal value.

Scale-level content validity index (S-CVI)

The S-CVI average is persistently identical to the Average Congruency Percentage (ACP). Rubio et al. [15] demonstrated their content validity procedure while evolving the Caregiver Well-Being Scale, in which they used the averaging approach for the S-CVI based

on ratings of relevance by six judges. This method for the calculation of S-CVI used in particular with concern that while performing universal approach with more than six raters the content validity index would be slightly depressed because wide-reaching approach demands agreement among all experts. Similarly, Waltz and Strickland [14] stated a recommendation on the standard value for the acceptability of S-CVI as 0.90 but not 0.80.

Reliability

While instituting the quality of a settled instrument wholly, Kuder and Richardson developed a formula known as KR-20. In estimating the regularity of a test based on internal consistency, also called a reliability coefficient, KR-20 has been the most widely used formula. It requires only a single administration of an assessment. The internal consistency by KR-20 obtained by evaluating the reliability of the material within a test based on the total number of items in the assessment as a whole, the proportion of participants giving correct answers for each question and the standard deviation of total score obtained. The value could range from 0 to 1. The closer the score is to 1, the more reliable the test. A study [16] stated when KR-20 formula is in use, internal consistency estimates the range from 0.75 as an acceptable mark to an excellent 0.97 mark [12].

A KR20 value range of 0.86 to 0.94 was reported by Lin et al. [17], for the analysis of internal consistency while doing item analysis of a multiple choice test questionnaire was used in licensure examination for registered nurses. Also, Sampaio and Sequeira [10] used KR reliability for their accurate and false tests in development of a valid instrument to assess nurses' knowledge of High-risk medication in a tertiary care hospital and got a value of 0.74, which indicated acceptable reliability. While, Priscila et al. [18] did the Brazilian transformation of the work done by Taiwanese researchers [10], where they too computed KR 20 formulae for their instrument to assess nurses' knowledge and obtained a value of 0.74 respectively. Similarly, Farhan et al. [19] designed an instrument for students to observe tests involved in the research of a string ensemble course for music students and while examining its reliability, the KR 20 formula showed a value as 0.717. Also, a study developed an Instructor-Mediated Performance Assessment Test, for which they did reliability and attained an index of 0.95 [20].

Conclusion

This study was an effort to prepare suitable informative materials in the form of videos that would benefit community pharmacists on their perspective about HRM and its management. The insufficiency of basic knowledge on this topic in the Indian setting of community pharmacy was highly reflected when pharmacists were evaluated previously with a set of a questionnaire regarding various aspects of HRM management, out of the objective. With this concern, the instrument in the form of learning modules consisting of demanding information on diverse aspects of high-risk medication management was prepared and finally validated. The validation done by content validity showed ideal values with I-CVI of 0.913 and S-CVI of 0.916. Similarly reliability followed with KR 20 formula analysis. The reliability index was 0.937, 0.8424 and 0.8195 for chapter 1, chapter 2 and chapter 3, respectively as parts of the learning modules. The prepared instrument thus can be concluded as a valid and reliable source to benefit community pharmacists for management of HRM and finally for better patient care.

Funding

This work was supported by Rajiv Gandhi University of Health Sciences: ORDER NO. RGU: Adv. Res. Proposal-P-147:2015-16 DATE:06-01- 2016.

Acknowledgement

The author sincerely thanks to Dr K Ramesh, Director, Karnataka College of Pharmacy, Bengaluru for his valuable contribution and consistent encouragement.

References

1. Australian Commission on Safety and Quality in Health Care.
2. Graham S, Clopp MP, Kostek NE, Crawford B (2008) Implementation of a High-Alert Medication Program. *Perm J* 12: 15-22.
3. Bohomol E (2014) Medication errors: descriptive study of medication classes and high-alert medication. *Esc Anna Nery* 2014; 18:311-316.
4. Engels MJ, Ciarkowski SL (2015) Nursing, pharmacy, and prescriber knowledge and perceptions of high-alert medications in a large, academic medical hospital. *Hosp Pharm* 50: 287-300.
5. Gandhi TK, Borus J, Weingart SN, Seger AC (2003) Adverse drug events: ambulatory care setting. *Clin Alert* 41: 7-7.
6. Gurwitz J, Field T, Avorn J, McCormick D, Jain S, et al. (2000) Incidence and preventability of adverse drug events in nursing homes. *Am J Med* 109: 87-94.
7. Bolarinwa O (2015) Principles and methods of validity and reliability testing of questionnaires used in social and health science researches. *Niger Postgrad Med J* 22: 195.
8. Hsaio G, Chen I, Yu S, Wei I, Fang Y, et al. (2009) Nurses' knowledge of high-alert medications: instrument development and validation. *J Adv Nurs* 66: 177-190.
9. Peterson G, Wu M, Bergin J (1999) Pharmacist's attitudes towards dispensing errors: their causes and prevention. *J Clin Pharm Ther* 24: 57-71.
10. Sampaio F, Sequeira C (2014) Nurses' knowledge and practices in cases of acute and chronic confusion: A questionnaire survey. *Perspect Psychiatr Care* 51: 98-105.
11. Lynn MR (1986) Determination and quantification of content validity. *Nurs Res* 35: 382-385.
12. Polit D, Beck C (2006) The content validity index: Are you sure you know what's being reported? critique and recommendations. *Res Nurs Health* 29: 489-497.
13. Zohrabi M (2013). Mixed method research: instruments, validity, reliability and reporting findings. *Theory and practice in language studies* 3: 254-262.
14. Waltz CF, Strickland OL (1989) Measurement of nursing outcomes. *Res Nurs Health* 12: 399.
15. Rubio D, Berg-Weger M, Tebb S, Lee E, Rauch S (2003) Objectifying content validity: Conducting a content validity study in social work research. *Soc Work Res* 27: 94-104.
16. Measurement error studies at the national center for education statistics (1997).
17. Lin LC, Tseng HM, Wu SC (1999) Item analysis of the registered nurse licensure exam taken by nurse candidates from vocational nursing high schools in Taiwan. *Proc Natl Sci Counc* 9: 24-31.
18. Priscila PA, Roberta CMR, Juliana BDS, Silvia RS, Maria HDML (2015) Translation, adaptation and practicability of nurses' knowledge of high alert medications to the Brazilian culture. *Rev Eletr Enf* 17: 1-12.
19. Farhan B, Soltani T, Do R, Perez C, Choi H, et al. (2018) Face, content, and construct validations of endoscopic needle injection simulator for transurethral bulking agent in treatment of stress urinary incontinence. *J Surg Educ* 75: 1673-1678.
20. Rutherford-Hemming T (2015) Determining content validity and reporting a content validity index for simulation scenarios. *Nursing Educ Perspect* 36: 389-393.