

Open Access

Diabetes and Metabolic Syndrome Can Contribute to Recurrent Vascular Events in Patients with Lacunar Stroke?

Roever L* and Resende ES

Department of Clinical Research, Federal University of Uberlândia, Brazil

Lacunar stroke corresponds to 15-25% of all ischemic strokes, and has a more favorable prognosis and their survival rates at 30 days and 1 year are about 90-100%, and about 70-80% patients are functionally independent at 1 year, compared with less than 50% of patients with non-lacunar stroke. Patients with a lacunar stroke usually present with a classical lacunar syndrome (sensorimotor stroke, pure sensory syndrome, ataxic hemiparesis or dysarthria-clumsy hand. and an atypical lacunar syndrome). Lacunar stroke may involve the thalamus, basal ganglia and pons [1-7].

Zhu and colleagues reported a study with 2,999 participants (SPS3 trial), 25% had metabolic syndrome (MetS) only, 6% had diabetes mellitus (DM) only, 32% had both conditions. In 3.8 years of follow-up, there were 274 recurrent strokes (240 ischemic, 34 hemorrhagic) and 74 myocardial infarctions (MI); among the 240 ischemic strokes, 134 (56%) were lacunar. The rates of any recurrent stroke (HR 1.7, 95% [CI] 1.3-2.3) or lacunar stroke (HR 2.4, 95% [CI] 1.5-3.7) were significantly higher for those with concurrent METS and DM. Risk of incident MI was higher in participants with DM (HR 2.8, 95% CI 1.1-7.0) or concurrent DM and MetS (HR 2.6, 95% CI 1.4-4.9).

MetS and DM were associated with stroke recurrence.

Lacunar stroke reduction is to control vascular risk factors such as hypertension, DM, MetS, dyslipidemia, visceral obesity and smoking cessation. Changes in lifestyle like a healthy diet, exercise, reducing stress levels are also recommended strategies.

References

- 1. Arboix A, Martí Vilalta JL (2009) Lacunar stroke. Expert Rev Neurother 9: 179-196.
- Esenwa C, Gutierrez J (2015) Secondary stroke prevention: challenges and solutions. Vasc Health Risk Manag 11:437-450.
- Arboix A (2015) Cardiovascular risk factors for acute stroke: Risk profiles in the different subtypes of ischemic stroke. World J Clin Cases 3: 418-429.
- 4. 4. Caplan LR (2015) Lacunar infarction and small vessel disease: pathology and pathophysiology. J Stroke 17: 2-6.
- Mok V, Kim JS (2015) Prevention and Management of Cerebral Small Vessel Disease. J Stroke 17: 111-122.
- Tsai CF, Anderson N, Thomas B, Sudlow CL (2015) Risk factors for ischemic stroke and its subtypes in Chinese vs. Caucasians: Systematic review and meta-analysis. Int J Stroke 10: 485-493.
- Zhu S, McClure LA, Lau H, Romero JR, White CL, et al. (2015) Recurrent vascular events in lacunar stroke patients with metabolic syndrome and/or diabetes. Neurology 85: 935-941.

*Corresponding author: Leonardo Roever, Department of Clinical Research, Federal University of Uberlândia, Av. Pará, P.O Box: 1720, Bairro Umuarama, Brazil, Tel: 553488039878; E-mail: leonardoroever@hotmail.com

Received September 08, 2015; Accepted September 22, 2015; Published September 25, 2015

Citation: Roever L, Resende ES (2015) Diabetes and Metabolic Syndrome Can Contribute to Recurrent Vascular Events in Patients with Lacunar Stroke? J Neurol Disord S1: e101. doi:10.4172/2329-6895.S1-e101

Copyright: © 2015 Roever L, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.