

## Drug-Mediated Image-Guided Radiotherapy—Mining for the Mother Lode

Charles A. Kunos\*

Department of Radiation Oncology, Summa Health System, Akron, OH, USA

The therapeutic nugget desired of anticancer agents is not to restore normal biologic processes in cancer cells but rather is to kill them. Non-selective cytotoxic agents inherently strike against cancer cells, but also make collateral adverse effects in healthy cells a major health concern. In this way, non-selective cytotoxic agents might be considered 'fools gold' to some. However, targeted anticancer agents have made the prospect of striking against specific biomolecular differences between cancer cells and healthy cells an attractive therapeutic strategy. For instance, a new potent ribonucleotide reductase inhibitor has been mined for its claim of accentuating radiation and chemotherapy effects with clinical benefit [1,2]. What if this or other novel biological therapeutic could serve as imaging agents used to discriminate cancer cells from healthy cells in a better way?

Here, the radio chemotherapeutic management of advanced stage cervical cancer provides a meaningful illustrative example. Cervical cancers over express ribonucleotide reductase [3], a ubiquitous rate-limiting enzyme responsible for the de novo generation of the building blocks of DNA. Healthy cervix cells have less expression of ribonucleotide reductase [4]. Envision a biologic anti cancer agent that nestles around an iron-stabilized radical in the ribonucleotide reductase enzyme and disrupts the vital proton-coupled electron transfer of its radical to its active site [5,6]. When radiation and chemotherapy damage DNA, and thus increase the demand for DNA building blocks, a blocked ribonucleotide reductase cannot provide the demanded supply of DNA precursors and cells die [1]. Clinically, the targeting of ribonucleotide reductase by such a potent inhibitor during radio chemotherapy has found success [2]. The eureka! Moment might be just that there is more to learn from this interesting drug-mediated therapeutic pharmacology.

There is a rich history of cancer-directed radiation therapy guided by 2-[18F] fluoro-2-deoxy-D-glucose Positron Emission Tomography (PET) images superimposed upon computed tomography radiation planning images [7-9]. But, PET images do not necessarily distinguish between cancer cells or immune cells rapidly dividing. Could an anticancer agent like a ribonucleotide reductase inhibitor be a more discriminatory nuclear imaging agent? APET-compatible, labeled ribonucleotide reductase inhibitor, that has desirable properties of target specificity and of short circulation time, could fill this role. The

overarching idea is to mark cancer cell bearing tissue using a labeled ribonucleotide reductase inhibitor, overlay mined drug-mediated PET images and radiation therapy planning images, irradiate cancer-bearing tissue precisely, and thus eradicate cancer cells.

New cancer research strategies have focused on the clinical development of targeted biologic agents that aim to accentuate cancer cell kill. Clinical development of labeled targeted biological agents might also prove 'golden' in the setting of drug-mediated image-guided radiotherapy. Future study is warranted.

### References

1. Kunos CA, Radivoyevitch T, Pink J, Chiu SM, Stefan T, et al. (2010) Ribonucleotide reductase inhibition enhances chemoradiosensitivity of human cervical cancers. *Radiat Res* 174: 574-581.
2. Kunos CA, Radivoyevitch T, Waggoner S, Debernardo R, Zanotti K, et al. (2013) Radiochemotherapy plus 3-aminopyridine-2-carboxaldehyde thiosemicarbazone (3-AP, NSC #663249) in advanced-stage cervical and vaginal cancers. *Gynecol Oncol* 130: 75-80.
3. Kunos CA, Winter K, Dicker AP, Small W Jr, Abdul-Karim FW, et al. (2013) Ribonucleotide reductase expression in cervical cancer: a radiation therapy oncology group translational science analysis. *Int J Gynecol Cancer* 23: 615-621.
4. Kunos CA, Radivoyevitch T, Kresak A, Dawson D, Jacobberger J, et al. (2012) Elevated ribonucleotide reductase levels associate with suppressed radiochemotherapy response in human cervical cancers. *Int J Gynecol Cancer* 22: 1463-1469.
5. Popovic-Bijelic A, Kowol CR, Lind ME et al (2011) Ribonucleotide reductase inhibition by metal complexes of Triapine (3-aminopyridine-2-carboxaldehyde thiosemicarbazone): a combined experimental and theoretical study. *J Inorg Biochem* 105(11):1422-1431.
6. Boal AK, Cotruvo JA Jr, Stubbe J, Rosenzweig AC (2010) Structural basis for activation of class Ib ribonucleotide reductase. *Science* 329: 1526-1530.
7. Lin LL, Mutic S, Malyapa RS, Low DA, Miller TR, et al. (2005) Sequential FDG-PET brachytherapy treatment planning in carcinoma of the cervix. *Int J Radiat Oncol Biol Phys* 63: 1494-1501.
8. Lin LL, Yang Z, Mutic S, Miller TR, Grigsby PW (2006) FDG-PET imaging for the assessment of physiologic volume response during radiotherapy in cervix cancer. *Int J Radiat Oncol Biol Phys* 65: 177-181.
9. Kunos C, Radivoyevitch T, Abdul-Karim FW et al (2011) 18F-fluoro-2-deoxy-D-glucose positron emission tomography standard uptake value ratio as an indicator of cervical cancer chemoradiation therapeutic response. *Int J Gynecol Cancer* 21(6):1117-1123.

\*Corresponding author: Charles Kunos, Summa Comprehensive Cancer Institute, Department of Radiation Oncology, 161 North Forge Street, USA, Tel: 330-375-3557; Fax: 330-375-3072; E-mail: [kunos@summahealth.org](mailto:kunos@summahealth.org)

Received October 16, 2013; Accepted October 18, 2013; Published October 21, 2013

Citation: Kunos CA (2013) Drug-Mediated Image-Guided Radiotherapy—Mining for the Mother Lode. *OMICS J Radiology* 2: e121. doi:10.4172/2167-7964.1000e121

Copyright: © 2013 Kunos CA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.