

Dynamics Phytoremediation of Zn and Diesel Fuel in Co-contaminated Soil using Biowastes

Agamuthu P and Dadrasnia A*

Institute of Biological Sciences, Faculty of Science, University of Malaya, Malaysia

Abstract

In phytoremediation of co-contaminated soil, developing strategies is a major challenge for simultaneous and efficient remediation of multiple pollutants. A lab-scale experiment was set up to investigate the efficiency of adding 5% (w/w) individually of three different organic waste amendments [tea leaves, soycake and potato skin] to enhance the phytoaccumulation of Zinc and diesel fuel co-contaminated soil by *Dracaena reflexa* for a period of 180 days. Soil contaminated with Zn (80 mg kg⁻¹DW) was spiked with an initial concentration of diesel fuel (25000 mg kg⁻¹DW). Application of biowastes demonstrates significant degradation of DF compared to control soil. The highest rate of oil degradation was recorded in co-contaminated soil planted with *D. reflexa* and amended with SC (85%) followed by PS and TL (60 and 53.3%). Although the plants did not accumulate any hydrocarbon from the contaminated soil but there was significant bioaccumulation of Zn in the root and stem of *Dracaena* plant which was observed. At the end of 180 days, 16.5 mg/kg and 12.2 mg/kg of Zn in the root and the stem of *D. reflexa* was recorded. Results indicated that the *D. reflexa* could effectively extract Zn from Zn-contaminated soils even in the presence of diesel fuel concentration.

Keywords: Biowastes; Bioaccumulation; Diesel oil; Phytodegradation; Zinc

Abbreviations: TL: Tea Leaf; SC: Soy Cake; PS: Potato Skin; TPHs: Total Petroleum Hydrocarbons; DF: Diesel Fuel; DW: Dry Weight; MHs: Heavy Metals; °C: Degree Celsius; GC/MS: Gas Chromatography/Mass Spectrometry; EPA: Environmental Protection Agency; BCF: Bioconcentration Factor; TF: Translocation Factor

Introduction

The two major sources of soil pollution are toxic metals and petroleum products. Millions of cubic meters of soil are polluted by these sources [1]. Total petroleum hydrocarbon (TPH) and Zn are persistent concerns due to their insolubility and presence in the environment. US Environmental Protection Agency (USEPA) (1999) reported that more than 40% of the national priority list sites are co-contaminated with both PAHs and heavy metals which are environmental concerns and must be reduced to acceptable levels [2]. Food web contamination is a one of the important pathways through which toxic pollutants enter into the human body. Consequently, heavy metals are often found to coexist in with PAHs, due to the same pollution sources. Metal-PAHs combination has been found in industrial and agricultural places where a strong relation between heavy metal and PAHs was observed [3]. Chemical and physical methods, as well as, a combination of them have been used to remediate co-contaminated soils. Information regarding the mechanisms, translocation, accumulation and the combined uptake of heavy metals and PAHs present in the soil, sediment and contamination are still under study. Due to rapid industrialization and urbanization in the world, soil pollution by PAHs and Cd has been considered during the last decade [4].

Phytoremediation technology has been considered for restoration of co-contaminated soil by organic pollutants and toxic metal [5,6]. The main challenge to phytodegradation of co-contaminated lands is the concurrent removal of various pollutants. For phytodegradation of hydrocarbon compounds or metals, many hyper accumulating plants have been widely studied, as plant root exudates lead to increase in the population of native soil microbes [4,7]. Several researchers reported that some plants could forbid the loss of organic compounds in contaminated soil [8,9]. For instance, *Sedum alfredii* is a native plant in

China and it has been demonstrated an efficient transport of Cd in the presence of pyrene or phenanthrene from Cd-contaminated soil into roots to shoots compared to other hyperaccumulators [10]. However, additional techniques are required to carry out simultaneous removal of PAHs and heavy metal from co-contaminated soils. The application of animal manures and organic fertilizers is a common approach to extract and enhance the PAHs and/or heavy metal from contaminated soil. It must be noted however this research is focused on organic waste (biowastes) as the supplements which are available cheap and easy to apply for remediation of soil and also replace fertilizers which are expensive. This could be an option for waste management especially in those areas that generate a high rate of food, vegetables and fruit wastes. However, the effects of biowastes on phytoextraction of heavy metals and PAHs in co-contaminated soil is not comprehensively studied, the objective of this study is to determine the feasibility of biowastes to remedy co-contaminated soil.

Materials and Methods

Collection of organic wastes, soil and diesel fuel

The garden soil was taken from a farm in Subang Jaya, Selangor, Malaysia. The organic wastes used in this study were collected from different locations; tea leaf (TL) and potato skin (PS) was collected from the IGS (Institute of Graduate Studies) canteen, University of Malaya, while the soycake (SC) was made in the laboratory. Physicochemical properties of organic wastes and soil employed were determined using standard methods.

*Corresponding author: Dadrasnia A, Institute of Biological Sciences, Faculty of Science, University of Malaya, Malaysia, Tel: +603 79674631; Fax: +603 79674631; E-mail: are.dadrasnia@gmail.com

Received November 23, 2013; Accepted December 30, 2014; Published January 05, 2014

Citation: Agamuthu P, Dadrasnia A (2013) Dynamics Phytoremediation of Zn and Diesel Fuel in Co-contaminated Soil using Biowastes. J Bioremed Biodeg S4: 006. doi:10.4172/2155-6199.S4-006

Copyright: © 2014 Agamuthu P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Preparation of co-contaminated soil

According to the critical level of heavy metals pollution in Malaysian soil [11], it was decided to use Zn (80 mg kg⁻¹). Plastic bags were tilled with 2 kg of soil and the desired amount of Zn was spiked with an aqueous solution of ZnCl₂ as a source of Zn in 200 ml of distilled water. It was sprayed layer by layer, and the soil was completely mixed. Then, with the purpose of creating a balance between the different fractions of elements in the soil, the soil samples were put through an incubation period of one month. During this period, the moisture of the bags was kept at about (70% ± 10). At the end of the incubation period, soil samples were co-contaminated with 2.5 % w/w of diesel fuel and thoroughly mixed well. 5% (w/w) of different biowastes (TL, SC and PS) were also mixed individually with the oil contaminated soil. Then plants transfer to treatments. The experiments were carried out in three replicates at room temperature (30 ± 2°C). The plants were moderately watered with ordinary tap water every three days to prevent leaching from the plastic bags. The design of the experiment is shown in Table 1.

Sample collection

The pot experiment was monitored under laboratory conditions (Solid waste laboratory, University of Malaya) at a humidity of 75-85% and temperature 30-32°C. Fifteen grams of soil sample were collected from the phytoremediation experiments monthly and analyzed immediately across six months study duration. The soil samples were taken from the rhizosphere zone of the plants from each plastic bag monthly, and were homogenized for the analysis.

The total amount of diesel fuel loss in soil was determined by suspending ten grams of soil in 20 ml of acetone: hexane (1:1 v/v) in a 250 ml Erlenmeyer flask. After shaking for one hour in an orbital shaker, the oil-solvent mixture was filtered into a beaker of known weight and the solvent was completely evaporated by rotary evaporator (A1000S). The new weight of the beaker was recorded. Percentage degradation of oil was estimated using the follow formula [10];

$$\% \text{ biodegradation} = [\text{TPH control} - \text{TPH treatment} / \text{TPH control}] \times 100 \quad (1)$$

Determination of biomass

The biomass of plants growing in clean soil and under diesel fuel and heavy metal stress (the control) was measured [12]. Prior to the analysis, plant samples were carefully washed with tap water and thoroughly rinsed with deionized water to remove any soil particles attached to the plant surfaces and wiped with filter paper to remove excess liquid before determining the fresh weight. Stem height and root length and also measured. Then the samples were oven-dried at 70°C until the weight was constant and the dry weight was recorded [13].

Analysis of Zn and Diesel fuel

Plants were uprooted at the end of the experiment (180 days). The root tissue was extracted with a 1:1 hexane/acetone in a Soxhlet

Treatment	Details of Treatment
A	2 Kg soil+2.5 % oil+5% TL+80 ppm Zn+ <i>Dracaena reflexa</i>
B	2 Kg soil+2.5 % oil+5% SC+80 ppm Zn+ <i>Dracaena reflexa</i>
C	2 Kg soil+2.5 % oil+5% PS+80 ppm Zn+ <i>Dracaena reflexa</i>
D	2 Kg soil+2.5 % oil+80 ppm Zn+ <i>Dracaena reflexa</i>
E	2 Kg soil+80 ppm Zn++ <i>Dracaena reflexa</i>
F	2 Kg soil+80 ppm Zn+2.5 % oil
G	2 Kg autoclaved soil+2.5 % oil+80 ppm Zn+0.5% NaN ₃

Table 1: Experimental Design for phytoremediation of co-contaminated soil.

extractor for 10 hours to determine if the plant tissues had absorbed the hydrocarbon from the soil. To assess hydrocarbon content removal, the extracts were analyzed for hydrocarbons using GC (gas chromatography) with a mass-selective detector (QP2010A). The GC was equipped with cross-linked 5% phenyl methyl siloxane capillary column. Helium was used as carrier gas. The temperature was set at 40°C and raised by 10°C/min until 300°C, which was held for eight minutes. The amount of heavy metal in plant tissues was determined by hot plate wet digestion method [14]. The Zn concentration in the soil was determined by the EPA method 3050B (acid extraction method). Briefly, soil was dried at 40°C and grounds with a laboratory blender (Waring model). 1 g samples were placed in a 250 ml flask for digestion. Then the samples were heated at the 95°C with 10 ml of 50% HNO₃ without boiling. This was followed by the addition of 65% HNO₃ until no brown fumes were given up by the samples. Then, gradually, 10 ml of 30% H₂O₂ and 37% HCL were added at the 95°C for 15 minutes. The digested samples obtained were filtered (0.45 µm filter paper) and diluted to 100 ml with deionized water and analyzed by Inductively Coupled Plasma-Optical Emission Spectroscopy (ICP- OES).

Calculations of Translocation factor and Bioconcentration factor

The Translocation Factor (TF) index is described as the capability of heavy metals from the soil and transfer to the edible part of vegetables l

$$TF = \frac{\text{Zn concentration inedible part of plant}}{\text{Zn concentration in root of plant}} \quad (2)$$

For plants, the Bioconcentration factor (BCF) is as a measure of the metal accumulation efficiency. A BCF value of greater than one is an implication of plants potential to phytoextraction. BCF was calculated using the following formula,

$$BCF = \frac{\text{Average Zn conc. in the whole plant tissue (mg kg}^{-1}\text{)}}{\text{Zn in the soil (mg kg}^{-1}\text{)}} \quad (3)$$

Rate of metal uptake by plants

The rate of uptake of Zn by *Dracaena* was calculated using the first order kinetic model as follows:

$$k = -1/t (\ln M / Mo) \quad (\text{Eq. 4})$$

Where; k=rate of metal uptake per month, t=time in month, M=mass of residual metal in the soil (ppm), Mo=initial mass of metal in the soil (ppm).

The data were analyzed for significant differences (p<0.05) between treatments using analyses of variance (ANOVA) with SPSS 18.0, and Duncan test comparison was used to determine the difference between treatments [15,16].

Results and Discussion

The physicochemical properties of the investigated soils and biowastes used are presented in Table 2. The native soil had a natural pH (~7) with low N amount (0.24 %) and P content (0.08 %) (Table 2).

Loss of diesel fuel in co-contaminated soil

The percentage of diesel degradation in all the amendments ranged from 9 to 85% in co-contaminated soil planted with *D. reflexa* (Figure 1). The percentage of biodegradation in contaminated soil containing *D. reflexa* without organic wastes recorded 50%, while in the control soil without the plant recorded a 15% oil loss during the period of study.

Parameters	Organic Wastes			
	Soil	TL	sc	PS
Total nitrogen (%)	0.8 ± 0.1	1.02 ± 0.08	1.3 ± 0.1	1.10 ± 0.04
Phosphorus (%)	0.6 ± 0.5	0.7 ± 0.6	0.9 ± 0.9	0.7 ± 0.1
Moisture content (%)	10.2 ± 0.8	34.3 ± 0.5	75.9 ± 1.6	62.1 ± 2.0
Organic C (%)	13.1 ± 1.3	55.6 ± 1.2	72.2 ± 0.9	66.3 ± 1.1
pH (1/ 2.5 soil/ water, w/v)	7.03 ± 1.5	6.5 ± 1.2	6.8 ± 1.2	6.9 ± 0.5

TL: Tea Leaf, SC: Soy Cake, PS: Potato Skin, Each value represents the mean (n=3)

Table 2: Characteristics of soil and organic wastes used for the phytoremediation.




Figure 1: Biodegradation of diesel fuel and Zinc from soil. Bars indicate standard error (n = 3). TL: tea leaf, SC: soycake, PS: potato skin.

Treatments	Roots	Stems	Leaves
Soil+2.5% oil+TL	15.1	10.9	ND
Soil+2.5% oil+SC	16.5	12.2	6.3
Soil+2.5% oil+PS	12.2	8.2	4.2
Soil+2.5% oil	3.4	1.8	ND
Soil without oil	1.4	ND	0.08

ND: Not detected

Table 3: Zn (mg/Kg) contents in plants in soil contaminated with 80 ppm Zn.

About 8% oil loss was recorded in autoclaved control soil without plant and polluted with Zn and 2.5% diesel fuel. The degradation of oil in autoclaved control soil (without plant) could be a result of some non-biological factors such as volatilization or photodegradation which was recorded in poisoned treatments amended with oil and sodium azide. In addition, the highest rate of degradation was recorded in the soil amended with SC which is probably due to a higher amount of N and P, compared to other organic waste amendments. It was also reported by work that soil amended with SC, showed a higher rate of degradation [17,18]. Similarly, Dominguez-Rosado and Pichtel [19] recorded 67% engine oil loss in contaminated soil planted with mustard and sunflower plants. ANOVA result shows a significant difference between soil treated with different biowastes, unplanted contaminated soil and contaminated soil planted with *D. reflexa* at both concentrations ($P < 0.05$). This is similar to the finding of Vouillamoz and Milke [20] who indicated that addition of compost allowed diesel loss down from 1200 to 200 mg TPH kg⁻¹ in contaminated soil planted with yre

grass. Kim et al. [21] also recorded a significant reduction during the phytodegradation of diesel-contaminated soil at the end of 120 days.

Bioaccumulation of Zn by *D. reflexa*

Diesel fuel contained the lowest concentration of Zn compared to unplanted polluted soil. However, soil used for phytoremediation had 18.1 (mg/kg) Zn. After the soil was artificially polluted with 2.5% diesel oil and 80 (mg/kg) Zn, the amount of Zn reached 74.52 (mg/kg). Zn accumulation in treating soil amended with organic wastes recorded a range of between 32.5 mg/Kg and 41 mg/Kg, whereas in polluted and unamended treatment there was a higher quantity of Zn. At the end of 180 days, appreciable quantities of Zn (16.5, 12.2 and 6.3 ppm) were detected at *D. reflexa* roots, stem and leaf (Table 3). The result is in conflict with the findings of Palmroth et al. [22] which demonstrated that contaminated soil with weathered hydrocarbons and amended with fertilizers (NPK) and biowaste composts; there is no accumulation of heavy metals in plant contextures. In general, from the results obtained and described above, *D. reflexa* obviously has the ability to accumulate heavy metals in tissues. Hence, the accumulation of Zn in *D. reflexa* plant amended with organic wastes was higher than with unamended control soil and treated with the *D. reflexa* without organic wastes. Organic waste amendments might provide a suitable condition to increase the bioavailability of metals in hydrocarbon polluted soil by enhancing the capability of the plant to uptake these metals in different plant tissues. The result is in line with the finding of Tan et al. [23] who reported that *D. reflexa* has the ability to accumulate Zn and Cd in different tissues in soil polluted with Zn and Cd, under greenhouse conditions. However, the result is in contrast to Clemente et al. [24] who reported that the mobility of heavy metals could be reduced in the case of fresh contamination by adding organic amendments and vegetation.

Plant uptake of diesel fuel

Gas chromatography mass spectrometry (GC/MS) analysis of the plant extract did not show the accumulation of hydrocarbons in any of the treatments. This is in contrast with the results of Palmroth et al. [25] who reported uptake of diesel oil by the root of the grass. The differences might be due to the different plants used in the studied; it might also be due to differences in the weather conditions as study by Palmroth et al. [25]. The result suggests that the mechanism of hydrocarbon removal by plants might be via phytovolatilization or rhizodegradation which has been well documented [26,27]. This is supported by the findings of different authors, who stated that flavonoids and other compounds released by roots can stimulate growth and activity of hydrocarbon degrading bacteria [25,28].

Translocation, Bioconcentration factors and rate of Zn uptake by *D. reflexa*

The Bioconcentration factor was higher in those treatments amended with organic wastes compared with the unamended control soil. This might be because of available nutrients for the plant growth that produced high plant biomass, thereby causing bioaccumulation of the metals in the plant tissues more than those of the unamended control treatments. Statistical analysis shows that there is a significant difference between TF in stems and leaves ($P < 0.05$). The highest TF in leaves and steam was observed in the soil amended with SC (Table 4).

The highest BCF was observed in SC amended soil, while the lowest was, in the soil without any contamination (0.01) and plant. In terms of TF in stems and leaves, statistical analysis does not show significant differences between different treatments at $P < 0.05$ (Table 4).

Treatments	TF (Stems)	TF (Leaves)	BCF	K (Monthly)
Soil+2.5% oil+TL	0.72	0.00	0.34	0.13
Soil+2.5% oil+SC	0.73	0.38	0.47	0.12
Soil+2.5% oil+PS	0.67	0.33	0.32	0.09
Soil+2.5% oil	0.52	0.00	0.06	0.04
Soil without oil	0.00	0.05	0.01	0.09

Table 4: Bioconcentration, translocation factors and rate uptake of Zinc (mg/Kg) with *D. reflexa* remediated soil.




Figure 2: Root and shoot biomass of *D. reflexa* after 180 days of growth. The same letter represents no significant difference ($p > 0.05$).

The result is differed that of Adesodun et al. [29] who reported that TF of Zn in soil remediated with sunflower and contaminated with Zn was more than one. The differences in the results recorded in the two studies might be because of the use of different plants for phytoremediation.

Plant growth and biomass

After six months of growth, no plant death was recorded in co-contaminated soil; however, some of the plants showed signs of phytotoxicity such as stunted growth and yellowing of leaves compared with the control. The highest *D. reflexa* longitudinal growth was observed in the amendment with SC contaminated soils. The plant growth was significantly improved by application of organic wastes (Figure 2).

Conclusion

Within the 180 day period of study soils amended with biowastes recorded the highest rate of metal uptake than the unamended soil. However, soil amended with SC recorded a higher rate of uptake of Zn contamination, followed by the soil which was treated with PS and TL, but in the Zn contaminated soil the rate of uptake was as follows SC > TL > PS. This demonstrates that the potential of organic waste amendments to enhance phytoremediation of the oil and at the same time bioaccumulation of metals. The result is in line with the finding of the uptake rate in soil with *D. reflexa* plant.

Acknowledgement

This study was supported by the University of Malaya IPPP Grants FP014/2010A, PV055/2011B and PS300/ 2010B. The authors would like to thank Dr. Babak Motesharezadeh for his technical assistance.

References

- Zukauskaite A, Jakubauskaite V, Belous O, Ambrazaitiene D, Stasiskiene Z (2008) Impact of heavy metals on the oil products biodegradation process. Waste Management & Research 26 : 500 -507.

- Reddy K, Al-Hamdan A, Ala P (2011) Enhanced Soil Flushing for Simultaneous Removal of PAHs and Heavy Metals from Industrial Contaminated Soil. Journal of Hazardous, Toxic and Radioactive Waste 15:166-174
- Li Hua W, Liu X, Tientchen CM, Chen YX (2008) Heavy Metals and PAHs in Sewage Sludge from Twelve Wastewater Treatment Plants in Zhejiang Province. Biomedical and Environmental Sciences 21: 345 - 352
- Sun M, Fu D, Teng Y, Shen Y, Luo Y, et al. (2011) In situ phytoremediation of PAH-contaminated soil by intercropping alfalfa (*Medicago sativa* L.) with tall fescue (*Festuca arundinacea* Schreb.) and associated soil microbial activity. Journal of Soils and Sediments 11: 980-989.
- Zhang G, Ling JY, Sun H, Luo J, Fan Y, et al. (2009) Isolation and characterization of a newly isolated polycyclic aromatic hydrocarbons-degrading *Janibacter anophelis* strain JY11. Journal of Hazardous Materials 172: 580-586
- Lin X, Xiaojun L, Li P, Li F, Zhang L, et al. (2008) Evaluation of Plant-Microorganism Synergy for the Remediation of Diesel Fuel Contaminated Soil. Bulletin of Environmental Contamination and Toxicology. 81 (1): 19-24
- Liu PG, Whang L, Yang M, Cheng S (2008) Biodegradation of diesel-contaminated soil: A soil column study. Journal of the Chinese Institute of Chemical Engineers 39: 419-428
- Smith KE, Schwab AP, Banks MK (2008) Dissipation of PAHs in saturated, dredged sediments: A field trial. Chemosphere 72: 1614-1619
- Cofield N, Banks MK, Schwab AP (2007) Evaluation of hydrophobicity in PAH-contaminated soils during phytoremediation. Environmental Pollution 145: 60-67
- Wang K, Zhang J, Zhu Z, Huang H, Li Ti, et al. (2012) Pig manure vermicompost (PMVC) can improve phytoremediation of Cd and PAHs co-contaminated soil by *Sedum affredii*. Journal of Soils and Sediments 12: 1089-1099
- Ibrahim R B (2009) Contaminated Land Management and Control Guidelines No. 1: Malaysian Recommended Site Screening Levels for Contaminated Land. Kuala Lumpur, Malaysia
- Phillips LA, Greer CW, Germida JJ (2006) Culture-based and culture-independent assessment of the impact of mixed and single plant treatments on rhizosphere microbial communities in hydrocarbon contaminated flare-pit soil. Soil Biology & Biochemistry 38: 2823-2833
- Merkel N, Kraft RS, Arias M (2006) Effect of the tropical grass *Brachiaria brizantha* (Hochst. ex A. Rich.) Stap on microbial population and activity in petroleum contaminated soil. Microbiological Research 161: 80-91
- Marin Ş, Lacrimioara Ş, Ceclia R (2011) Evaluation of performance parameters for trace element analysis in perennial plants using icp-as a technique. Journal of Plant Development 18: 87-93
- Li Q, Cai S, Mo C, Chu B, Peng L, et al. (2010) Toxic effects of heavy metals and their accumulation in vegetables grown in a saline soil. Ecotoxicology & Environmental Safety 73: 84-88
- Ashraf MA, Maah MJ, Yusoff I (2011) Heavy metals accumulation in plants growing in ex tin mining catchment. International Journal of Environmental Science & Technology 8: 401- 416
- Dadrasnia A, Agamuthu P (2013a) Dynamics of diesel fuel degradation in contaminated soil using organic wastes. International Journal of Environmental Science & Technology 10: 769-778
- Dadrasnia A, Agamuthu P (2010). Enhanced Degradation of Diesel-Contaminated Soil using Organic Wastes. Malaysian Journal of Science 29: 225-230
- Dominguez-Rosado E, Pichtel J (2005) Transformation of fulvic substances in the rhizosphere during phytoremediation of used motor oil. Journal of Environmental Science and Health, 39: 2369 – 2381
- Vouillamoz J, Milke MW (2001) Effect of compost in phytoremediation of diesel contaminated soils. Water Science Technology 43: 291 – 295
- Kim D, Woo SM, Yim J, Kim T, Thao N, et al. (2010) The feasibility of phytoremediation combined with bioethanol feedstock production on diesel-contaminated soil. Paper read at 1 – 6 August, 19th World Congress of Soil Science, Soil Solutions for a Changing World, Brisbane, Australia
- Palmroth M, Koskinen P, Kaksonen A, Münster U, Pichtel J, et al. (2007) Metabolic and phylogenetic analysis of microbial communities during phytoremediation of soil contaminated with weathered hydrocarbons and heavy metals. Biodegradation Journal 18: 769-782

23. Tan LP, Lim TT, Jie H (2007) Mass screening of tropical plants to study their potential as heavy metal phyto remediators. 10th Int. Conference on Environmental Science & Technology, Kos island, Greece
24. Clemente R, Almela C, Bernal MP (2006) A remediation strategy based on active phyto remediation followed by natural attenuation in a soil contaminated by pyrite waste. Environmental Pollution 143: 397-406
25. Palmroth MRT, Pichtel J, Puhakka JA (2002) Phyto remediation of subarctic soil contaminated with diesel fuel. Bioresource Technology 84: 221-228
26. Dadrasnia A, Agamuthu P (2013b) Organic wastes to enhance phyto-treatment of diesel-contaminated soil. International Journal of waste management & research. 31: 1133-1139
27. Gerhardt KE, Xiao-Dong H, Glick BR, Greenberg BM (2009) Phyto remediation and rhizo remediation of organic soil contaminants: Potential and challenges. Plant Science. 176: 20-30
28. Muratova A, Turkovskaya OV, Hübner T, Kusch P (2003) Studies of the Efficacy of Alfalfa and Reed in the Phyto remediation of Hydrocarbon-Polluted Soil. Applied Biochemistry and Microbiology 39: 599-605
29. Adesodun JK, Atayese M, Agbaje T, Osadiaye B, Mafe O, et al. (2010) Phyto remediation Potentials of Sunflowers for Metals in Soils Contaminated with Zinc and Lead Nitrates. Water, Air, Soil Pollution 207: 195-201

Citation: Agamuthu P, Dadrasnia A (2013) Dynamics Phyto remediation of Zn and Diesel Fuel in Co-contaminated Soil using Biowastes. J Bioremed Biodeg S4: 006. doi:[10.4172/2155-6199.S4-006](https://doi.org/10.4172/2155-6199.S4-006)

This article was originally published in a special issue, **Biofuels & their Applications** handled by Editor(s). Dr. Kirill I Kostyanovskiy, Texas Agri Life Research & Extension Center, USA

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/biobiogroup>

