

Ecotoxicity Evaluation of Industrial Discharge Waters and Metallic Solutions using Two Organisms (*Lactuca sativa* and *Daphnia magna*)

Anne Priac, Amandine Poupene, Coline Druart and Grégorio Crini*

Chrono-Environment, UMR 6249 usc INRA, University of Franche-Comté/CNRS, 16 route de Gray, 25000 Besançon, France

Abstract

Surface treatment industrial discharge water is a complex anthropogenic source of pollutants, including organic pollutants (PAHs, VOCs...) and numerous metal ions. We attempted to identify the main toxicants comparing impact assessment of real polycontaminated effluents and reconstituted polymetallic solutions via ecotoxicological bioassays performed with *Daphnia magna* immobilization test (24 h) and *Lactuca sativa* germination test (168 h). We focused first on 2 (Ni and Zn) then on 5 metals (Ni, Zn, Co, Cr, Al). Our results showed differences between metal toxicity order: Zn>Al>Ni>Cr~Co, for daphnids and Ni>Zn>Al~Co>Cr for lettuce. However, discharge waters remained more toxic than synthetic solutions: those 5 metals were not entirely responsible for the discharge water ecotoxicity. We also found *D. magna* to be more sensitive than *L. sativa*. This last assessment should be interpreted with care, knowing that immobilization and germination tests are respectively acute and chronic toxicity bioassays. Thus, battery tests are appropriate to evaluate industrial discharge water samples, and should be increasingly used as eco toxicological standards.

Keywords: Bioassay; Heavy metals; Reconstituted solutions; Waste water

Introduction

Industrial discharge waters, especially those from the surface treatment (ST) industry, released into the aquatic ecosystems have their own set of various environmental and sanitary issues, due to the fact that various loads of hazardous substances including: metallic trace elements (MTE; mostly Zn, Ni, Cu, Cr, Sn and Al), organic matter (oils, solvents, etc.) and diverse organics such as polycyclic aromatic hydrocarbons (PAHs) and volatile organic compounds (VOCs) [1]. ST industries are, like other industrial sectors, subject to specific release regulations, notably for metals. Although the discharge waters usually respect the regulatory standards, the present metals could be assimilated by fauna and flora and thus lead to long term toxic effects on the environment [2,3]. Nowadays, while pollutant mixtures present in discharge water after treatment are relatively easy to characterize chemically, assessing their impact on the environment is usually difficult and has rarely been reported [4]. Finally, the toxicity of treated ST waste remains poorly defined.

To assess the biological and chemical quality of water, 4 main kinds of approach can be used: (1) Chemical analysis to characterize the water mass studied qualitatively and quantitatively, (2) Comparing the analytical data to ecotoxicological information available in the literature to reach an *a priori* assessment of the hazard of substances (as in Draft Assessment Reports for pesticides), (3) Laboratory bioassays to assess the toxicity of substances and (4) *in situ* studies using native organisms or via active bio indication to assess the risk of natural populations exposed to substances released in the environment. Laboratory bioassays for water quality assessment are numerous and offer a large choice of indicators [5-7]. Three different types of standardized bioassays are the most commonly used, notably for the regulatory framework for chemicals management. They represent 3 trophic levels: primary producers with algae, primary consumers with crustaceans and secondary consumers with fish. Among them, the short-term bioassay based on the immobilization of a freshwater crustacean, *Daphnia magna*, is a test also used in the ecotoxicological assessment of industrial discharge waters. Nevertheless, it was pointed out that toxicity strongly relies on the choice of bio indicators and

the endpoints used in the bioassays since sensitivity varies among taxonomic groups and species [8-10]. Consequently, it may be very useful to assess discharge water thanks to various bio-indicators in order to increase the ecological representativeness, to include a panel of sensitivity and to avoid a major risk of environmental effects and toxicity underestimation [11,12]. Recently, phytotoxicity tests using plants such as *Lactuca sativa* have been also proposed to assess the impact of industrial effluents by our group for the first time [3]. Our results demonstrated that these tests were simple, quick and reliable. Moreover, the use of these bioassays also presented the advantage of being inexpensive and not requiring major equipment as also reported in other works [8,10,13]. However, these tests were mainly used to assess the toxicity of single substances, such as metals (Table 1) [10,14-28] and there is a lack of studies concerning the impact of complex matrices such as discharge waters [29] or synthetic solutions of several metals.

The aim of this work was to assess the environmental impact of industrial discharge waters poly-contaminated with metals and to determine which metal(s) is (are) most responsible for the toxicity through the use of 2 bio-indicators *Daphnia magna* and *Lactuca sativa* via reconstituted solutions.

Materials and methods

Toxicity bioassays

Standardized germination tests [30] were performed following the method previously described in detail by Charles et al. [3]. The test assessed the germination of 30 plump lettuce seeds (*Lactuca sativa*

*Corresponding author: Grégorio Crini, Chrono-Environment, UMR 6249 usc INRA, University of Franche-Comté/CNRS, 16 route de Gray, 25000 Besançon, France, Tel: +33381665701; E-mail: gregorio.crimi@univ-fcomte.fr

Received September 04, 2014; Accepted September 22, 2014; Published October 01, 2014

Citation: Priac A, Poupene A, Druart C, Crini G (2014) Ecotoxicity Evaluation of Industrial Discharge Waters and Metallic Solutions using Two Organisms (*Lactuca sativa* and *Daphnia magna*). J Pollut Eff Cont 2: 117 doi: [10.4172/2375-4397.1000117](http://dx.doi.org/10.4172/2375-4397.1000117)

Copyright: © 2014 Crini G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Element	Bioassay indicator	Index	Endpoint	Concentration [mg L ⁻¹]	Reference
Zn	<i>L. sativa</i> (var <i>n.r.</i>)	96h EC ₅₀	Root elongation	1	[14]
	<i>P. subcapitata</i>	72h EC ₅₀	Growth rate	0.042	[15]
	<i>D. magna</i>	48h LC ₅₀	Death	0.970	[16]
	<i>G. pulex</i>	48h LC ₅₀	Death	4.920	[17]
Ni	<i>L. sativa</i> (var <i>Tro.</i>)	NOEC	Growth rate	1.8	[10]
	<i>P. subcapitata</i>	96h EC ₅₀	Population	0.233	[18]
	<i>D. magna</i>	48h LC ₅₀	<i>n.r.</i>	6.9	[19]
	<i>G. sp</i>	96h LC ₅₀	<i>n.r.</i>	13	[20]
Cr	<i>L. sativa</i> (var <i>Rav.</i>)	72h EC ₅₀	Growth rate	5.9	[21]
	<i>P. subcapitata</i>	72h EC ₅₀	Population	0.030	[22]
	<i>D. magna</i>	48h LC ₅₀	Immobilization	0.290	[23]
	<i>G. pulex</i>	48h LC ₅₀	Death	0.809	[24]
Co	<i>D. magna</i>	48h LC ₅₀	Death	4.4	[25]
Al	<i>D. magna</i>	48h LC ₅₀	Immobilization	3.9	[26]
Cu	<i>L. sativa</i> (var <i>n.r.</i>)	96h EC ₅₀	Root elongation	3	[14]
	<i>P. subcapitata</i>	72h EC ₅₀	Growth rate	0.020	[15]
	<i>D. magna</i>	48h LC ₅₀	Death	0.0111	[27]
	<i>G. pulex</i>	48h LC ₅₀	Death	0.047	[28]

Specific lettuce varieties: *n.r.*, non reported; *Tro.*, Trocadero; *Rav.*, Ravel
 End-points: LC₅₀, lethal concentration for 50% of the individuals tested; NOEC, no observed effect concentration.

Table 1: Toxicities of metallic trace elements (published data) on different organisms (*Lactuca sativa*, *Daphnia magna*, *Pseudoskinneriella subcapitata* and *Gammarus pulex* or *sp.*).

Sample	Concentrations [mg L ⁻¹]			EC ₅₀ [% of DW]	Germination (%)
	Ni	Zn	<i>D. magna</i>		
DW1	0.34	1.91	17	54	
DW2	0.28	1.51	5.2	58	
DW3	0.54	2.46	21	42	
DW4	0.35	1.84	11.3	48	
DW5	0.51	2.06	32	44	

Every control GR was higher than the required 90% of seed germination
 All concentrations were above the quantification limits

Table 2: Ecotoxicity (EC₅₀ and GR) of different discharge water samples (DW1 to DW5) on *Daphnia magna* and *Lactuca sativa* respectively in relation to Zn and Ni concentrations.

(*L.*) *var* Batavia Dorée de Printemps) watered with Reverse Osmosis Water ROW (controls; pH=6 ± 0.2), Discharge Water DW or Synthetic Solution SS, in triplicates, for 7 days, in the dark at 24 ± 1°C, on a filter paper substrate. As recommended by the standard, DW or SS pH must be between 5.5 and 9. To validate the test, germination rates (GR) of controls must be higher than or equal to 90%.

Bioassays using *Daphnia magna* were carried out by an accredited analysis laboratory (CARSO, Lyon, France). The test was performed according to the “Inhibition Protocol Mobility” described in the standardized biomonitoring test ISO [31].

Industrial discharge waters & synthetic solutions

Five DW (denoted DW1 to DW5) were firstly collected in a ST company in Franche-Comté over a one-year period. Effluents were average sample characteristic of that day’s activity. As the 2 main issues to be dealt with in DWs were Ni and Zn (specific company threshold emission values for these 2 MTE were 3.5 mg L⁻¹), DW5 Ni and Zn concentrations were mimicked in single and binary solutions S: S1 (Zn=2 mg L⁻¹), S2 (Ni=0.5 mg L⁻¹) and S3 (Zn=2 mg L⁻¹ and Ni=0.5 mg L⁻¹). Each solution ecotoxicity was evaluated with germination test. We also determined Ni and Zn EC₅₀ (concentrations range: 0 to 300 mg L⁻¹) for *L. sativa* and *D. magna*.

Four others DWs (DW6 to DW9) were then collected in the same

company. Ni, Zn, Al, Cr and Co concentrations of these DWs were mimicked in mixture SS denoted SS6 to SS9. Each solution ecotoxicity was evaluated with germination and immobilization test (Table 2). We also determined Al, Co and Cr EC₅₀ (concentrations range: 0 to 1000 mg L⁻¹) for *L. sativa* and *D. magna*.

For each of these 9 DWs, EC₅₀ (expressed in percentage of DW) was determined through lettuce germination and daphnids immobilization tests. DWs samples were diluted with ROW. Every metallic synthetic solutions were prepared in ROW from sulfate salts of Al, Co, Cr, Ni and Zn (purchased from Fisher Scientific, France).

Chemical analyses

For each DW sample and synthetic solution, pH was determined (pH meter, model 3110, WTW, Alès, France). Metal concentrations were measured by spectrophotometry (cuvette test and/or reagent tests; portable Spectroflex 6100, WTW, Alès, France) or by ICP-AES (ThermoFisher, iCAP 6500 radial model, Courtaboeuf, France) after acid digestion for DWs, following a previously reported method [1]. All results are expressed in mg L⁻¹.

Statistical analysis

Germination rates of control, DW5, S1, S2 and S3 were compared using the Kruskal-Wallis test, with a significance threshold of *p*<0.05. All statistical analyses were performed with R (2.15.1) (R Development Core Team, 2013). Dose-dependent curves and EC₅₀ values were calculated with Hill’s model using the macro Excel Regtox free version EV 7.0.6.

Results and discussion

The toxicity of the first 5 DWs was studied through 2 bio-indicators (Table 2). The results showed deleterious effects on both bio-indicators since EC₅₀ were low for daphnids (below 32%) and lettuce seed germination rates were significantly lower than those of controls (>90%). Due to activities of the industry focused on in our study, investigations of toxicity were firstly led on Ni and Zn. Concentrations

Bio-indicator	EC ₅₀ [mg L ⁻¹]				
	Al	Co	Cr	Ni	Zn
<i>D. magna</i>	8.45	11.69	10.36	9.8	6.35
<i>L. sativa</i>	237	247.7	265	58.3	154.3

Table 3: EC₅₀ values for daphnia and lettuce for 5 MTE detected in DW samples ICP-AES analysis (Al, Co, Cr, Ni and Zn).




Figure 1: Germination (%) of *L. sativa* seeds for the DW5 sample and synthetic solutions S1, S2 and S3. Different letters indicated significant differences (p-value <0.01).

Sample	Metal ion [mg L ⁻¹]					EC ₅₀ [% of DW or SS]	
	Al	Co	Cr	Ni	Zn	<i>D. magna</i>	<i>L. sativa</i>
DW6	5.09	2.75	0.15	0.62	2.67	6.1	45
SS6	5.06	2.72	0.15	0.63	2.69	56.8	86
DW7	5.70	4.05	0.35	0.74	1.97	5	68
SS7	5.58	4.06	0.24	0.70	1.96	52.1	84
DW8	2.66	1.69	0.24	0.31	1.45	ND	66
SS8	3.64	2.28	0.29	0.32	1.49	72.2	84
DW9	5.36	3.58	0.25	0.40	2.05	18.4	68
SS9	5.26	3.71	0.26	0.41	2.79	47.9	86

ND, not determined

All concentrations were above the quantification limits

Table 4: Concentrations of 5 metals (mg L⁻¹) in 4 discharge waters (DW6 to 9) and synthetic solutions (SS6 to 9) in relation to toxicity on *D. magna* and *L. sativa* (EC₅₀ in % of DW or SS).

of both these suspected toxicants are presented in Table 2 and showed daily variability (as previously reported by Charles et al. [3]). From an analytical point of view, the chemical composition in Ni and Zn can be ranked as follows: DW3 > DW5 > DW1 ~ DW 4 > DW2. For the 2 bioassays, the sample toxicity range (decreasing order) was:

- DW2 > DW4 > DW1 > DW3 > DW5 for *D. magna* and
- DW3 > DW5 > DW4 > DW1 > DW2 for *L. sativa*.

The more toxic DWs for *D. magna* were the less toxic for *L. sativa*. This was also confirmed by Castillo et al. [32] studying the impact of final tannery industrial effluent (daphnids EC₅₀ 24 h=77.9%) and lettuce (EC_{50-root growth} 120 h>90%). Despite its low coefficient of variation CV (14%), it appeared that lettuce DW toxicity could be linked to Ni and Zn concentrations (except for DW4 and DW1 which were inverted). For daphnids, no correlation was shown (CV=59%). This variability was explained by the production activity, as suggested by Hitchcock et al. [33] who calculated a CV reaching 133.7% for the mortality of nematodes exposed to industrial effluents (pulp and paper industries).

To verify the hypothesis that Ni and Zn concentrations in the DW can be linked to lettuce ecotoxicological response, we ran germination tests on synthetic solutions S1, S2 and S3 containing Ni and Zn, alone or in a mixture, in the same concentrations as those found in DW5. We also performed ecotoxicological tests on both *D. magna* and *L. sativa* (Table 3), to assess individual EC₅₀ of nickel and zinc. The results showed that GR of single (S1 and S2) and binary (S3) synthetic solutions were not significantly different from the control (Figure 1). This result was not surprising in regard to the values of EC₅₀ determined in *L. sativa* for Ni and Zn (Table 3) which were far above the concentrations found in DWs. However, the EC₅₀ results were not expected considering those found in the literature (Table 1). Indeed, toxicity values for other endpoints were much lower, of the order of 1 mg L⁻¹, both for Zn and Ni [10]. This major disparity could be explained by the variety of lettuce used for the assay, as criticized by Priac et al. (unpublished work) who demonstrated that among 4 varieties, Batavia (used in the present paper) was the least sensitive. Significant differences were found between the germination rates of lettuce exposed to synthetic solutions of Zn and Ni and those exposed to the DWs at same concentrations (Figure 1). Similar experiments and interpretations were reported by Yoo et al. [34] with Cu, Ag and cyanides, to reproduce an effluent from a lead frame manufacturing factory. Unlike our results, those of Yoo et al. [34] demonstrated that these 3 substances were responsible for the toxicity of the effluent on daphnids since they observed a similarity in the toxicity of the real and the synthetic effluents. In the present study, the DW toxicity observed on *L. sativa* was not explained only by the presence and the concentrations in Zn and Ni.

Investigations were conducted on a larger number of metals potentially responsible for the toxicity of DWs. Among 23 elements measured, 15 were present at quantifiable levels at least once, and 5 of them (Al, Co, Cr, Ni and Zn) were selected for the following experiments owing to their concentrations in DW6 to DW9 (higher than 1 mg L⁻¹; Table 4) and/or their known effects on the environment (Table 1). DWs 6, 7 and 9 appeared to be much more toxic (EC₅₀ 6.1, 5, 18.4% of the sample) than their respective SS (56.8, 52.1, 47.9% of the solution tested) on *D. magna*. Results showed the same tendency for the GR of *L. sativa*, but not as dramatic: for instance daphnid EC₅₀ values were 5 and 52.1% for DW7 and SS7, respectively, whereas lettuce EC₅₀ values were 68 and 84%. Like for Ni and Zn, the presence of Al, Co and Cr did not explain all the DW toxicity on both *D. magna* and *L. sativa*, even though daphnid EC₅₀ values showed these 5 metals to be toxic (Table 3).

To our knowledge, few studies have assessed the environmental impact of discharge water or synthetic solutions on more than one bio-indicator [7,12,14,35,36]. Bioassay batteries have already been shown to be a relevant way to evaluate toxicity, irrespective of the ecosystem studied [7,36,37]. Sensitivity differences observed between daphnids and lettuce (Tables 2-4) also occurred on comparison with data from the literature (Table 1). General differences can be explained by bioassay endpoint (acute or chronic toxicities) or protocol variability (bioindicator subspecies or cultivars, animal gender, lapse of exposure, number of individuals per Petri dish or tube, etc.; [38]). Yet, it appears that differences between bioindicator sensitivity remain in bibliographic data. For 3 metals for which we found comparative results (Zn, Ni, Cr), toxicity ranges were different for lettuce (Zn<Ni<<Cr) compared to algae, daphnids and gammarids (Cr<Zn<<Ni) as described in Table 1. Table 4 also shows single EC₅₀ differences between indicators: toxicity range for daphnids being (from less to more toxic) Cr, Ni and Zn while the lettuce toxicity range was Cr, Zn and Ni. Another difference

between these 2 bioindicators was related to the order of magnitude of the EC₅₀ (e.g. lettuce nickel EC₅₀: 58.3 mg L⁻¹ vs daphnids: 9.8 mg L⁻¹).

Conclusions

In this study, the 2 bioindicators *Lactuca sativa* and *Daphnia magna* were proved to be pertinent to assess the ecotoxicity of polycontaminated discharge water from the surface treatment industry. The results showed that metal-based synthetic single and mixed solutions were less toxic than the discharge water, meaning that the ecotoxicity of these effluents could not be explained only by the 5 metals chosen in this work. Consequently, it would be interesting to lead future investigations not only towards a more exhaustive determination of the chemical composition of discharge water but also possible interactions (e.g. additivity, antagonism, synergy) between metals and/or trace organics and/or other minerals. Results also demonstrated that lettuce was more resistant than daphnids to the discharge waters and synthetic solutions. Ecotoxicological assessments complete chemical analyses as they integrate all chemical interactions. As reported in this study the use of a battery of tests was a relevant tool to include the whole variability of toxicity.

Acknowledgment

The authors are grateful to *Ville de Besançon* which funded Anne Priac's PhD, to Sophie Gavaille and Céline Lagarrigue from the *Agence de l'Eau Rhône Méditerranée Corse*, the *Conseil Régional de Franche-Comté*, and the FEDER (*Fonds Européens de Développement Régional*) for financial support (NIRHOFEX Program 2013-2016).

Conflict of Interests

The authors declare that they have no conflict of interest.

References

- Morin-Crini N, Druart C, Gavaille S, Lagarrigue C, Crini G (2013) Analytical monitoring of the chemicals present in the discharge water generated by the surface treatment industry. J Environ Protect 4: 53-60.
- Komjarova I, Blust R (2009) Effect of Na, Ca and pH on simultaneous uptake of Cd, Cu, Ni, Pb, and Zn in the water flea *Daphnia magna* measured using stable isotopes. Aquatic Toxicol 94: 81-86.
- Charles J, Sancey B, Morin-Crini N, Badot PM, Degiorgi F, et al. (2011) Evaluation of the ecotoxicity of polycontaminated industrial effluents using the lettuce plant (*Lactuca sativa*) as a bioindicator. Ecotox Environ Saf 74: 2057-2064.
- Kadirova ZC, Hojamberdiev M, Katsumata KI, Isobe T, Matsushita N, et al. (2014) Photodegradation of gaseous acetaldehyde and methylene blue in aqueous solution with titanium dioxide-loaded activated carbon fiber polymer materials and aquatic plant ecotoxicity tests. Environ Sci Pollut R 21: 4309-4319.
- Cairns JJ, Pratt JR (1989) The scientific basis of bioassays. Hydrobiologia 188: 5-20.
- Farré M, Barceló D (2003) Toxicity testing of wastewater and sewage sludge by biosensors, bioassays and chemical analysis. Trac-Trend Anal Chem 22: 299-310.
- Pandard P, Devillers J, Charissou AM, Poulsen V, Jourdain MJ, et al. (2006) Selecting a battery of bioassays for ecotoxicological characterization of wastes. Sci Total Environ 363: 114-125.
- Wang W, Freemark K (1995) The use of plants for environmental monitoring and assessment. Ecotoxicol Environ Saf 30: 289-301.
- Wangberg SA, Bergström B, Blanck H, Svanberg O (1995) The relative sensitivity and sensitivity patterns of short-term toxicity tests applied to industrial waste water. Environ Toxic Water 10: 81-90.
- Di Salvatore M, Carafa AM, Carratu G (2008) Assessment of heavy metals phytotoxicity using seed germination and root elongation tests: A comparison of two growth substrates. Chemosphere 73: 1461-1464.
- Chapman PM, Fairbrother A, Brown D (1998) A critical evaluation of safety (uncertainty) factors for ecological risk assessment. Environ Toxicol Chem 17: 99-108.
- Hernando MD, Fernandez-Alba AR, Tauler R, Barcelo D (2005) Toxicity assays applied to wastewater treatment. Talanta 65: 358-366.
- Dutka BJ (1989) Methods for microbiological and toxicological analysis of waters, wastewaters and sediments. National Water Research Institute NWRI, Environment Canada.
- Fjällborg B, Li B, Nilsson E, Dave G (2006) Toxicity identification evaluation of five metals performed with two organisms (*Daphnia magna* and *Lactuca sativa*). Arch Environ Contam Toxicol 50: 196-204.
- Aruoja V, Dubourguier HC, Kasemets K, Kahru A (2009) Toxicity of nanoparticles of CuO, ZnO and TiO₂ to microalgae *Pseudokirchneriella subcapitata*. Sci Total Environ 407: 1461-1468.
- Gale NL, Wixson BG, Erten M (1992) An evaluation of the acute toxicity of lead, zinc and cadmium in Missouri Ozark groundwater. Trace Subst Environ Health 25: 169-183.
- McLoughlin N, Yin D, Maltby L, Wood RM, Yu H (2000) Evaluation of sensitivity and specificity of two crustacean biochemical biomarkers. Environ Toxicol Chem 19: 2085-2092.
- Chen CY, Lin KC, Yang DT (1997) Comparison of relative toxicity relationships based on batch and continuous algal toxicity tests. Chemosphere 35: 1959-1965.
- Cabejzek I, Stasiak M (1960) Investigation on the influence of some metals on the biocoenosis of water with the use of *Daphnia magna* as an indicator (Part I). Roczn Zabl Hig Warsaw 11: 303-312.
- Rehboldt R, Lasko L, Shaw C, Wirhowski E (1973) The acute toxicity of some heavy metals ions toward benthic organisms. Bull Environ Contam Toxicol 10: 291-294.
- Adema DMM, Henzen L (1989) A comparison of plant toxicities of some industrial chemicals in soil culture and soilless cultures. Ecotoxicol Environ Saf 18: 219-229.
- Nyholm N (1990) Expression of results from growth-inhibition toxicity tests with algae. Arch Environ Contam Toxicol 19: 518-522.
- Diamantino TC, Guilhermino L, Almeida MF, Soares AMVM (2000) Toxicity of sodium molybdate and sodium dichromate to *Daphnia magna* Straus evaluated in acute, chronic, and acetylcholinesterase inhibition tests. Ecotoxicol Environ Saf 45: 253-259.
- EPA (1993) Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms (EPA/600/4-90/02TF), US Environmental Protection Agency; Office of Toxic Substances: Washington, D.C.
- EPA (1975) Methods for acute toxicity test with fish, macroinvertebrates and amphibians (EPA/660/3-75-009), US Environmental Protection Agency; Office of Toxic Substances: Washington, D.C.
- Biesinger KE, Christensen GM (1972) Effects of various metals on the survival, growth, reproduction, and metabolism of *Daphnia magna*. J Fish Res Board Can 29: 1691-1700.
- Long KE, Van Genderen EJ, Klaine SJ (2004) The effects of low hardness and pH on copper toxicity to *Daphnia magna*. Environ Toxicol Chem 23: 72-75.
- Taylor EJ, Maund SJ, Pascoe D (1991) Toxicity of four common pollutants to the freshwater macroinvertebrates *Chironomus riparius* Meigen (Insecta: diptera) and *Gammarus pulex* (L.) (Crustacea: amphipoda). Arch Environ Contam Toxicol 21: 371-376.
- Backhaus T, Altenburger R, Arrhenius Å, Blanck H, Faust M, et al. (2003) The BEAM-project: prediction and assessment of mixture toxicities in the aquatic environment. Cont Shelf Res 23: 1757-1769.
- ISO (2005) Soil Quality – Determination of the effects of pollutants on soil flora – Screening test for emergence of lettuce seedlings (*Lactuca sativa* L.). International Organisation for Standardization ISO 17126. Geneva, Switzerland.
- ISO (1989) Water Quality – Determination of the inhibition of the mobility of *Daphnia magna* Straus (Cladocera, Crustacea). Acute Toxicity Assay. International Organisation for Standardization ISO 6341. Geneva, Switzerland.
- Hitchcock DR, Black HC, Williams PL (1997) Investigations into using the nematode *Caenorhabditis elegans* for municipal and industrial wastewater toxicity testing. Arch Environ Contam Toxicol 33: 252-260.
- Castillo GC, Vila IC, Neild E (2000) Ecotoxicity assessment of metals and wastewater using multitrophic assays. Environ Toxicol 15: 370-375.

34. Yoo J, Kim SB, Kim WK, Jung J (2014) Toxicity identification of effluent from a semiconductor lead frame manufacturing factory. J In Eng Chem 20: 494-498.
35. Bohorquez-Echeverry P, Campos-Pinilla C (2007) Assessment of *Lactuca sativa* and *Selenastrum capricornotum* like indicators of water toxicity. Universitas Scientiarum 12: 83-98.
36. Wilke BM, Riepert F, Koch C, Kühne T (2008) Ecotoxicological characterization of hazardous wastes. Ecotox Environ Saf 70: 283-293.
37. Ewell WS, Gorsuch JW, Kringle RO, Robillard KA, Spiegel RC (1986) Simultaneous evaluation of the acute effects of chemicals on seven aquatic species. Environ Toxicol Chem 5: 831-840.
38. Alexander PD, Alloway BJ, Dourado AM (2006) Genotypic variations in the accumulation of Cd, Cu, Pb and Zn exhibited by six commonly grown vegetables. Environ Pollut 144: 736-745.

Citation: Priac A, Poupenev A, Druart C, Crini G (2014) Ecotoxicity Evaluation of Industrial Discharge Waters and Metallic Solutions using Two Organisms (*Lactuca sativa* and *Daphnia magna*). J Pollut Eff Cont 2: 117 doi: [10.4172/2375-4397.1000117](https://doi.org/10.4172/2375-4397.1000117)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

