

Effect of Different Treatments on Excretion of Tetrodotoxin after Oral Administration in Rat

Hsiao-Chin Jen¹, Tzu-Chun Lin², Ya-Jung Wu^{1,3}, Yu-Chun Chen³, Cheng Hong Hsieh⁴ and Deng-Fwu Hwang^{3,4*}

¹Department of Food Technology and Marketing Management, Taipei College of Maritime Technology, Taiwan

²Department of Veterinary Medicine, National Pingtung University of Science and Technology, Pingtung, Taiwan

³Department of Food Science and Centre of Excellence for Marine Bioenvironment and Biotechnology, National Taiwan Ocean University, Taiwan

⁴Department of Health and Nutrition Biotechnology, Asia University, Taiwan

*Corresponding author: Deng-Fwu Hwang, Department of Food Science and Center of Excellence in Bioenvironment and Biotechnology, National Taiwan Ocean University, Taiwan, Tel: +886-2-24622192; E-mail: dfhwang@mail.ntou.edu.tw

Received date: January 23, 2015; Accepted date: April 14, 2015; Published date: April 25, 2015

Copyright: © 2015 Jen HC et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The levels of tetrodotoxin (TTX) in the serum and urine were investigated by using LC-MS method in healthy male wistar rats after oral administration with single dose of 60 nmole TTX mixture. After treatment, male wistar rats were randomly divided into three groups: (1) rats drink distilled water; (2) rats drink saline solution; (3) rats drink alkaline electrolyzed (AE) water. The serum and urine samples were collected at intervals between 2 hr to 60 hr after orally administration. It was found that TTX can be detected from serum and urine up to 60 hr the max that was tested when intoxicated poisoning occurred. The highest amount of TTX appeared after 4 hr and 6 hr in the serum and urine, respectively. When the rats took AE water, TTX was secreted earlier from the serum and urine. It indicates that AE water enhances the excretion of TTX in the rats. Moreover, LC-MS is useful for detecting TTX in serum and urine samples and may have diagnostic potential for TTX intoxication.

Keywords: Tetrodotoxin (TTX); Excretion; Rat; LC-MS

Introduction

Outbreaks of tetrodotoxin (TTX) food poisoning are still reported in various countries including Taiwan [1-6]. Until now the causative agent of these food poisonings was identified from the toxin in food. However, the suspected food was sometimes discarded and no leftover was retained. Hence, TTX diagnosis from patients is very important. TTX is a heterocyclic guanine compound, and acts by blocking the flux of sodium ions through voltage-gated sodium channels on excitable cell membranes [5].

In the past, the principle impediment to analyze TTX in the serum and urine was lack of an uncomplicated and reproducible method with sufficient sensitivity to detect the picogram to nanogram quantities thought to be present [7,8]. Hence, a new approach for examining TTX in the serum and blood was established by using C18 cartridge column, ultrafiltration and liquid chromatography- mass spectrometry (LC-MS) [9-15].

The purpose of this study was to establish a method for detecting trace concentration of TTX in the serum and urine of exposed rats in order to determine the time course of excretion and to determine the effectiveness of different supportive treatments. The results are suspected to be useful for enhancing potential method for TTX excretion in the patient.

Materials and Methods

Materials

The dried powder of mixed authentic tetrodotoxin (TTX), 4-epiTTX and anhydro-tetrodotoxin (anh-TTX) were obtained from Wako Pure Chemical Industries (Tokyo, Japan). The purity of TTX was 94%. The toxins were dissolved with 0.03 M acetic acid and kept at 20°C until use. Male Wistar rats (n=24) weighing between 210 g and 260 g (234 g ± 12 g) were purchased from National Laboratory Animal Centre, Taipei.

Experimental treatment of rats

Male Wistar (n=18) rats were fed individually in stainless steel wire bottom cages with a controlled environment (25°C, 50%-60% humidity, 12 hr light per day) for 2 weeks. The animals were fed a laboratory diet (PMI Feeds, USA). Experimental rats were each orally administered with a single dose of 60 nmole TTX (19.14 µg), which is toxic to lower than lethal dose to rats and is previously obtained by using rat bioassay to substitute mouse assay [14]. Then these rats were divided into three groups supplied with distilled water (as control), saline solution and alkaline electrolyzed (AE) water in free access, respectively. The saline solution was dissolved 0.9% NaCl in distilled water. The AE water (pH=9.0) was made by an electrolyzer (ROX-20TA, Hoshizaki Electric Co. Ltd., Toyooka, Aichi, Japan). Immediately after dosing, animals were transferred to metabolic cages, and the blood and urine samples were collected separately at 2hr, 4hr, 6hr, 8hr, 12hr, 24hr, 36hr, 48hr and 60 hr after dosing. The blood of rats were taken from the tail veins of each rat. The serum of blood samples were collected by centrifugation (2,000 x g for 15 min). The serum and urine samples were frozen at -20° until assay was carried

out. The other 6 rats were used to collect the serum and urine for establishing the standard curves and recoveries of TTX in the serum and urine by spiked with authentic TTX.

Sample preparation

The sample cleanup procedure used was based on the procedure developed by Andrinolo et al. [12]. Each (1.0 ml) of the serum and urine samples was thawed and mixed with 200 μ l of 0.5 M acetic acid to keep it acidic. The samples were immediately centrifuged at 10,000 \times g for 10 min. The supernatant was passed through a 3 ml prepared cartridge column (C18 Sep-Pak cartridges, particle size 37 μ m-55 μ m, sorbent weight 500 mg, Millipore, Waters, MA). The prepared cartridges were previously washed with 10 ml methanol and then 10 ml water before use. After applying sample into the cartridge, 10 ml 0.3% acetic acid was eluted. The eluant was freeze-dried, dissolved in 2 ml 0.3% acetic acid and filtered through a 3,000MW cut-off Ultrafree microcentrifuge filter (Micron YM-3, Millipore, Waters). The filtrate was freeze-dried and dissolved in a small amount (1 ml) of distilled water.

LC-MS analysis for TTXs

The chromatographic system was composed of a Waters Model 2690 quaternary LC pump coupled analysis. TTX was performed on a reverse-phased column (Puresil C-18 GP, 4.6 mm \times 250 mm, Waters) and the mobile phase was aqueous 30 mM heptafluorobutyric acid-1 mM ammonium acetate (pH 5.0) at flow rate of 1.0 ml/min for TTX. The injection volume was 10 μ l. The ZMD 2000 mass spectrometer equipped with an atmospheric pressure ion source and an electrospray ionization (ESI) interface (Finnigan MAT, CA, USA) was employed for detection. The electrospray interface was typically operated using the following setting fragmentor, 150 V; nebulizer, N₂ (60 psi); dry gas N₂ (10 l/min, 350 $^{\circ}$ C; V-cap, 3,500 V [9-11].

Standard curves, detection limit and recovery

The standard curve of TTX was prepared from 1 ng/ml, 5 ng/ml, 10 ng/ml, 50 ng/ml, 100 ng/ml of TTX. The determination was performed three times (n=3). Data from the curve were subjected to linear regression analysis and peak area (y) versus amount of TTX (x) was plotted. The detection limit that was evaluated according to the rates of sample peak area to noise area was more than three times. Spiked samples were prepared by dissolving approximate 1 μ g, 5 μ g, 10 μ g of TTX in normal serum and urine (1.0 ml). The sample were mixed with 500 μ l of 0.5 M acetic acid, centrifuged, passed through the cartridge column, and extracted with methanol as described above.

Statistical analysis

Statistical analysis for differences among rats in the experimental groups was performed by the one-way analysis of variance (ANOVA). When the ANOVA identified differences among groups, multiple comparisons among means were made using Duncan's new multiple range tests. A p<0.05 was considered statistically significant.

Results

When using LC-MS to determine TTX toxin, the standard curve was linear in the range of 1-100 ng/ml for serum and urine. The linear equations were $y=817x+2190$ ($r=0.996$) and $y=11.78x+1143.38$ ($r=0.992$) for serum and urine, respectively. The recovery of TTX in

the serum and urine were 91.2% and 94.3%, respectively. The detection limit was 0.5 ng/ml of TTX. Figure 1 contains LC-MS chromatograms in selected ions at m/z 302 and 320 for standard TTX, 4-epiTTX and anh-TTX, respectively.

Spiked samples were prepared by dissolving 18 nmole of TTX in the serum and urine (Figure 2). Figure 3 shows LC-MS chromatograms of the serum and urine samples of rat (4 hr). The rats were orally administered with a dose of 60 nmole of TTX, the serum and urine samples were collected during a 60 hr experimental course. The LC-MS could detect TTX amount in serum up to 36 hr and in the urine up to 60 hr. It was found that the highest amount of TTX appeared after 4 hr and 6 hr in the serum and urine, respectively. Hence, TTX can be detected from serum up to 36 hr and from urine up to 60 hr when intoxicated poisoning occurs. After 2 hr, TTX was secreted from the urine. The maximum peaks of TTX appeared at 6 hr in urine when the rats took each kind of water. By the end of the collection period (60 hr), almost TTX was excreted from the urine. Hence, TTX could be perfectly secreted after 60 hr of food poisoning.

The effect of other water (saline solution and AE water) on TTX metabolism in rats was also evaluated in this study. When the rats took saline solution and AE water, the time course of TTX excretion in rats is shown in Figure 4. When the rats took AE water, it was found that TTX could be secreted earlier from the urine at 4 hr. After 2 hr, TTX was secreted into the urine and clearly secreted after 60 hr. It appears that administration of AE water could stimulate a more rapid excretion of TTX through the urine (Figure 5).






Figure 1: LC-MS chromatograms in selected ion monitored (SIM) mode at m/z 302, 320 for authentic TTX, 4-epi TTX and anh-TTX.



Discussion

This is the first experimental effort to resemble TTX toxin incidents in rats. We have developed a simple technique for the determination of TTX in serum and urine of humans [9-11]. In this study, the unambiguous identification of TTX toxin in the rat was obtained using LC-MS method. The technique was applied to experimentally exposed animals and proved to be useful in confirming the toxins in even the smallest amount was demonstrated. LC-MS is quite rapid technique and has been considered as possible replacement for the mouse bioassay [13-17].



It is not known whether chemical conversion of toxin in the body happens when the mouse assay is used. Therefore, LC-MS method provided the reliable and simple alternative for trace TTX toxin in the serum and urine. In addition, there are few reports in the literature of TTX concentrations being determined in poisoned patients. Kawatsu et al. [18] indicated urine samples from 12 TTX poisoned patients, values in the range 6ng/ml-100 ng/ml were found. Oda et al. [19] mentioned urine excretion values of 281 μg/day, for the second day after ingestion, dropping to 16 μg/day on day 4 and undetectable on day 5. Urine is the major excretion route of TTX intoxication for human beings [15]. The maximum peaks of TTX appeared at 6 hr in urine when the rats took distilled water. By the end of the collection period (60 hr), almost TTX was excreted in the urine.

Although alkaline electrolyzed water is not used for medical treatment so far, it could promote the secretion of TTX in rats in this study. Hence, alkaline electrolyzed water may be considered as a useful solvent to treat the TTX intoxicated patient for promoting the recovery.



The time course of TTX in the rat is more consistent with each other, suggesting that TTX is similarly redistributed from the serum to its target sites at sodium channels. Alkaline electrolyzed water has high pH value (9.0). This character may elevate the pH value of intestine and change TTX (pKa 8.76) and anh-TTX (pKa 7.95) to molecule type (not ionic type). The molecule type of TTX exhibits more lipid soluble.

Therefore, TTX could be quickly absorbed and entered the metabolism system [1,20].

Our results demonstrate that LC-MS method is capable of detecting small quantities of TTX in rat serum and urine samples. TTX was detectable up to 60 hr. Although the toxicokinetics of TTX elimination will differ for rats and humans, this study indicates that this approach could be useful for diagnosing or confirming a diagnosis of TTX in humans.

Acknowledgement

This study was partially supported by National Science Council, Taiwan, R.O.C., and Center of Excellence on Marine Bioenvironment and Biotechnology, National Taiwan Ocean University.

References

1. Hwang DF, Noguchi T (2007) Tetrodotoxin poisoning. *Adv Food Nutr Res* 52: 141-236.
2. Hwang PA, Tsai YH, Lin SI, Hwang DF (2007) The gastropods possessing TTX and/or PSP. *Food Rev Int* 23: 321-340.
3. Noguchi T, Ebesu JSM (2001) Puffer poisoning: Epidemiology and treatment. *J Toxicol* 20: 1-10.
4. Noguchi T, Arakawa O (2008) Tetrodotoxin--distribution and accumulation in aquatic organisms, and cases of human intoxication. *Mar Drugs* 6: 220-242.
5. Yu FH, Catterall WA (2003) Overview of the voltage-gated sodium channel family. *Genome Biol* 4: 207.
6. Hwang PA, Tsai YH, Lu YH, Hwang DF (2003) Paralytic toxins in three new gastropod (Olividae) species implicated in food poisoning in southern Taiwan. *Toxicon* 41: 529-533.
7. Shui LM, Chen K, Wang JY, Mei HZ, Wang AZ, et al. (2003) Tetrodotoxin-associated snail poisoning in Zhoushan: a 25-year retrospective analysis. *J Food Prot* 66: 110-114.
8. O'Leary MA, Schneider JJ, Isbister GK (2004) Use of high performance liquid chromatography to measure tetrodotoxin in serum and urine of poisoned patients. *Toxicon* 44: 549-553.
9. Jen HC, Lin SJ, Tsai YH, Chen CH, Lin ZC, et al. (2008) Tetrodotoxin poisoning evidenced by solid-phase extraction combining with liquid chromatography-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 871: 95-100.
10. Tsai YH, Hwang DF, Cheng CA, Hwang CC, Deng JF (2006) Determination of tetrodotoxin in human urine and blood using C18 cartridge column, ultrafiltration and LC-MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 832: 75-80.
11. Hwang PA, Tsai YH, Deng JF, Cheng CA, Ho PH, et al. (2005) Identification of tetrodotoxin in a marine gastropod (*Nassarius glans*) responsible for human morbidity and mortality in Taiwan. *J Food Prot* 68: 1696-1701.
12. Andrinolo D, Michea LF, Lagos N (1999) Toxic effects, pharmacokinetics and clearance of saxitoxin, a component of paralytic shellfish poison (PSP), in cats. *Toxicon* 37: 447-464.
13. Lin CL, Hsieh CH, Hsieh YL, Tai YC, Hwang DF (2013) Application of LC-MS/MS in identification of toxin in the causative gastropod and victim samples. *J Mar Sci Tech* 21:52-57.
14. Wu YJ, Lin CL2, Chen CH2, Hsieh CH3, Jen HC, et al. (2014) Toxin and species identification of toxic octopus implicated into food poisoning in Taiwan. *Toxicon* 91: 96-102.
15. Jen HC, Nguyen TA, Wu YJ, Hoang T, Arakawa O, et al. (2015) Tetrodotoxin and paralytic shellfish poisons in gastropod species from Vietnam analyzed by high-performance liquid chromatography and liquid chromatography-tandem mass spectrometry. *J Food Drug Analysis* 22:178-188.
16. Hwang DF, Jeng SS (1991) Bioassay of tetrodotoxin by using ICR strain male mouse. *J Chin Biochem Soc* 20: 80-86.
17. Yasumoto T (1991) Tetrodotoxin, In: Environmental Health Bureau, Ministry of Health and Welfare (ed.), Standard Methods of Analysis in Food Safety Regulation-Chemistry. Japan Food Hygiene Association, Tokyo. pp. 296-300
18. Kawatsu K, Shibata T, Hamano Y (1999) Application of immunoaffinity chromatography for detection of tetrodotoxin from urine samples of poisoned patients. *Toxicon* 37: 325-333.
19. Oda K, Araki K, Totoki T, Shibasaki H (1989) Nerve conduction study of human tetrodotoxication. *Neurology* 39: 743-745.
20. Goto T, Kishi Y, Takahashi S, Hirata Y (1965) Tetrodotoxin. *Tetrahedron* 21: 2059-2088.