

Effect of Fruit Pectin on Growth of Lactic Acid Bacteria

Emon Chatterjee^{1*}, Suba GA Manuel² and Syed Shamimul Hassan³

¹Department of Biochemistry, Mount Carmel College, #58 Palace Road, Bengaluru, Karnataka, India

²Department of Life Science, Mount Carmel College, #58 Palace Road, Bengaluru, Karnataka, India

³School of Life Science, IGNOU, Maidan Garhi, New Delhi, India

*Corresponding author: Emon Chatterjee, Department of Biochemistry, Mount Carmel College, #58 Palace Road, Bengaluru, Karnataka, India, Tel: 9632082520; E-mail: emon.chatterjee@gmail.com

Received date: Feb 26, 2016; Accepted date: Apr 19, 2016; Published date: Apr 25, 2016

Copyright: © 2016 Chatterjee E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The concept of modulating gut health through diet is not new and dates back to at least the beginning of the 20th century. However, it is only recently that sound scientific rationales have been proposed and investigated. Three microflora modulation tools have emerged, the addition of exogenous living microorganisms to foods (i.e., probiotics), the selective stimulation of the growth and activity of beneficial microorganisms indigenous to the gut (i.e., prebiotics), and a combination of both approaches (i.e., synbiotics). Fruit wastes which are highly perishable and seasonal, is a problem to the processing industries and pollution monitoring agencies. A valuable byproduct that can be obtained from fruit wastes is pectin. An effort was taken to extract pectin from different fruit waste (*Musa* sp. and *Citrus limetta* and rind of *Citrullus lanatus* and putrefied fruits of *Solanum lycopersicum* and *Psidium guajava*). An attempt was made to observe the enhancement of growth of Lactic Acid Bacteria (LAB-*Lactobacillus casei*, *L. acidophilus*, *Bifidobacterium bifidum*) by introducing the pectin samples from the above fruit waste. It was observed that pectin was able to enhance the growth of the bacteria and the titrable acidity considerably. Hence it can be concluded that pectin that is extracted from fruit waste can be used to enhance the growth of LAB. The current study aims to prove pectin as a potential prebiotic.

Keywords: Pectin; *Musa acuminata*; *Citrus limetta*; *Citrullus lanatus*; *Solanum lycopersicum*; *Psidium guajava*; *Lactobacillus casei*; *L. acidophilus*; *Bifidobacterium bifidum*

Introduction

The food industry produces large volumes of wastes, both solids and liquids resulting from the production, preparation and consumption of food. These wastes pose increasing disposal and potentially severe pollution problems and represent a loss of valuable biomass and nutrients. Thus new methods and policies are needed for waste handling and treatment in order to prevent pollution of the environment [1]. By-product recovery from fruit wastes can improve the overall economics of processing units and can reduce the environmental pollution effectively. A valuable byproduct that can be obtained from fruit wastes is pectin. Pectin designates that water soluble pectinic acid (colloidal polygalacturonic acids) of varying methyl ester content and degree of neutralization, which are capable of forming gels with sugar, and acids, under suitable condition (GITCO). It is used in pharmaceutical preparation as filler, as an agglutinated in blood therapy and also to glaze candied fruits [2]. Pectins are mainly used as gelling agents but can also be used as thickener and water binder. The classical application is giving the jelly-like consistency to jams or marmalades which would otherwise be sweet juices. For household use, pectin is an ingredient in jelling sugar where it is diluted to the right concentration with sugar and some citric acid to adjust the pH. In some countries, pectin is also available as a solution or extract or as a blended powder for home jam making. Pectins can also be used to stabilize acidic protein drinks, such as drinking yogurt and as a fat substitute in baked foods. Typical levels of pectin used as a food

additive is about 0.5-1.0% this are about the same amount of pectin as in fresh fruit.

Pectin can also be a potential prebiotic. For a food ingredient to be classified as a prebiotic, it must neither be hydrolyzed nor absorbed in the upper part of the gastrointestinal tract; be a selective substrate for one or a limited number of potentially beneficial commensal bacteria in the colon, thus stimulating the bacteria to grow, become metabolically activated, or both; and be able to alter the colonic microflora toward a more healthier composition [3]. Although any food ingredient that enters the large intestine is a candidate prebiotic, it is the selectivity of the fermentation in the mixed culture environment that is critical. At present, most searches for prebiotics are directed toward the growth of lactic acid-producing microorganisms. This is due to their purported health-promoting properties. However, it may be that future developments in the study of prebiotics may include aspects of their effect on pathogenic flora components [4]. The term synbiotic is used when a product contains both probiotics and prebiotics. A synergy between probiotics and prebiotics is termed as a synbiotic system. The primary intention of using a synbiotic is to give a layer of protection for the bacteria during their travel through the gastrointestinal tract [3]. Hence the present study aims at extraction of pectin from peels of *Citrus limetta*, *Citrullus lanatus*, *Musa acuminata*, and the putrefied fractions of *Solanum lycopersicum* and *Psidium guajava* followed by the determination of the prebiotic potential of the different extracted pectin.

Materials and Methods

Extraction of pectin

Fruit extracts from rind of *Musa acuminata* c.v. banana, *Citrus limetta* c.v. sweet lime, *Citrullus lanatus* c.v. watermelon, *Malus domestica* c.v. apple and putrefied fruits of *Solanum lycopersicum* c.v. tomato and *Psidium guajava* c.v. guava were homogenized using deionized water (1:1.5 w/v). Lemon juice was added to the homogenate to adjust the pH to 2-2.5. The homogenate was then boiled at 80°C for 15 minutes. The boiled homogenate was filtered through a cloth filter followed by Whatman filter of under vacuum. The filtrate was then treated with isopropanol (1:1) to precipitate the pectin. The precipitated pectin was then recovered from the filtrate and dried for further use.

Probiotic culture maintenance

The Starter culture of *Lactobacillus acidophilus* was procured from Microbial Type Culture Collection and Gene Bank Institute of Microbial Technology, Chandigarh (MTCC number 10307) and the culture of *Bifidobacterium bifidum* and *Lactobacillus casei* was procured from the Department of Dairy Microbiology, Dairy Science College UAS, Bangalore. Probiotic strains were maintained individually in sterile MRS (Himedia) (De Man, Rogosa and Sharpe) broth at 37°C for 4 hours, and then refrigerated at 4°C. Subculturing was done in MRS media every 7 days.

Growth of LAB

Probiotic cultures were grown in MRS broth and were incubated at 37°C for 48 hrs. Growth was measured at 48 hrs and 60 hrs by optical density (660 nm) measurements. The effect of pectin on the growth of LAB was assessed by the addition of pectin (0.4%) from different fruit waste to the MRS broth maintained at different dilutions. The dilutions were plated on the MRS agar medium. All the experiments were done in triplicates.

Study of the titrable acidity

The titrable acidity of the samples was determined according to AOAC method. The milk samples (pasteurized skim milk) were curdled (6 hrs) using above LAB strains with pectin (0.4%). The curd was diluted and titrated against 0.1N sodium hydroxide, using phenolphthalein as indicator to a faint pink end point.

$$\% \text{ Titratable Acidity} = \frac{9 \times \text{titer value} \times \text{normality of NaOH} \times \text{Dilution Factor}}{\text{Weight of the sample taken in gms}}$$

Results and Discussion

Extraction of pectin

Pectins are structural polysaccharides present within all dicotyledonous plant cell walls. The primary structural feature of pectin is a linear, 4 α linked D-Galacturonic acid chain with varying degrees of methylation. The main raw materials used to produce commercial pectin are apple pomace and citrus peel [5]. The maximum amount of pectin (dry weight) was obtained from *Citrus limetta* (14.49%) and least quantity of pectin obtained from *Citrullus lanatus* (3.41%), whereas *Musa sp.* and *Psidium guajava* has similar

pectin concentration (5.57% and 5.19%) (Table 1). Simmaky and Jaanaki [6] reported the amount of pectin extracted from orange peels was 31.5 g which was equivalent to 15.25% of total orange peel weight, and the amount of pectin extracted from lemon peels was 41.5 g which was equivalent to 20.75% as total lemon peels weight which is similar to the current finding.

Fruit Wastes	Wet Weight (gms)	Dry weight (gms)	Percentage of pectin
<i>Solanum lycopersicum</i>	105	7.5	7.14
<i>Citrus limetta</i>	182.26	26.79	14.69
<i>Musa sp</i>	1409.33	78.6	5.57
<i>Citrullus lanatus</i>	454	15.5	3.41
<i>Psidium guajava</i>	1200	62.3	5.19

Table 1: Quantity of pectin obtained from fruit wastes.

Growth of LAB

The MRS broth culture was maintained at pH 6.0 (normal pH of the broth ranges between 5.7-6.3) and the colony forming units/ml was counted and was found to be almost similar for all the LAB samples (Table 2a).

Sample	Colony forming unit/mL (10 ⁻⁴) Mean ± SD	Colony forming unit/mL (10 ⁻⁵) Mean ± SD	Colony forming unit/mL (10 ⁻⁷) Mean ± SD
<i>L. acidophilus</i>	226 ± 8	34 ± 6.24	2 ± 1
<i>B. bifidum</i>	281.66 ± 8.08	20.33 ± 8.50	2 ± 1
<i>L. casei</i>	TNTC*	45 ± 7.00	2 ± 1

Table 2a: Growth of LAB at pH 6.0. *TNTC=Too numerous to count.

The optical density was measured at 660 nm, for *L. acidophilus* culture it was found to be 0.8 at 48 hrs of incubation and 0.77 by 60 hrs of incubation at 37°C. Whereas *B. bifidum* O.D. value was 0.6 at 48 hrs and 0.56 at 60 hrs of incubation. There was no significant change in the growth of LAB on incubation for 60 hrs, *L. casei* the OD was 1.004 at 48 hrs and 0.985 at 60 hrs. There was a reduction in the absorbance when the cultures were incubated more than 48 hrs (Table 2b).

Sample	OD at 660 nm at 48 hrs of incubation	OD at 660 nm at 60 hrs of incubation
<i>L. acidophilus</i>	0.8	0.77
<i>B. bifidum</i>	0.6	0.56
<i>L. casei</i>	1.004	0.985

Table 2b: Growth of LAB at 48 and 60 hrs of incubation.

The various growth responses of the bacteria to the pectin samples at various pH was also observed by Palframan et al. [7] stated that the pH at which the fermentation is carried out has a direct effect on the substrate metabolism due to the change in enzyme activity at different

pH. It was found that pectin samples increased the growth of Bifidobacteria. Similar results were also reported by Olano-Martin et al. [8], they investigated the effect of various pectin and pectin-oligosaccharides on Bifidobacteria which is accordance to the current study which indicates the enhancement of growth of LAB in the presence of the prebiotic pectin (Table 2c). It was observed that *L. casei* showed the maximum growth (2.4 OD at 660 nm) with pectin from *S. lycopersicum* as compared to *L. acidophilus* and *B. bifidum*. Yeo also indicated that prebiotics in the form of soy products are capable of significantly enhancing the growth of *L. acidophilus*.

Sample Pectin	Probiotic Culture	O.D. at 660 nm after 48 hours of incubation
Nil	Commercial curd	0.6
Musa sp	<i>L. acidophilus</i>	1.8
	<i>L. casei</i>	2.2
	<i>B. bifidum</i>	0.9
<i>C. lanatus</i>	<i>L. acidophilus</i>	1.9
	<i>L. casei</i>	1.8
	<i>B. bifidum</i>	0.7
<i>S. lycopersicum</i>	<i>L. acidophilus</i>	1.9
	<i>L. casei</i>	2.4
	<i>B. bifidum</i>	1.2
<i>Psidium sp</i>	<i>L. acidophilus</i>	1.7
	<i>L. casei</i>	2
	<i>B. bifidum</i>	0.9

Table 2c: Effect of pectin on the growth of LAB.

Titration acidity of synbiotic

Lactic acid is one of major products of lactose degradation in milk and milk products due to the bacterial fermentation. Depending on the microorganisms involved, fermentation of milk proceeds via glycolysis pathway will produce lactic acid while via pentose phosphate pathway with formation of lactic and acetic acids [9]. The titration acidity (as percentage lactic acid) of the synbiotic product ranged between 0.45 to 0.9% depending upon the type of culture (Table 3). High titration acidity (0.94–0.95%) was observed when all three stains of LAB was cultured together with the pectin samples. There was an increase in titration acidity of individual LAB culture when compared to the control (without pectin). The titration acidity of individual LAB culture varied between 0.52–0.57% with the pectin. The findings corresponded to Gomes and Malcata [10] study on acid tolerance of *L. acidophilus*, which varied from 0.3% to 1.9% titration acidity, with an optimum pH lying at 5.5 ± 6.0.

Prebiotic sample	Probiotic Culture	% Titration acidity
Nil	Commercial curd	0.1
Musa sp.	<i>L. acidophilus</i>	0.5
	<i>B. bifidum</i>	0.57

	<i>L. casei</i>	0.55
	<i>L. acidophilus</i> + <i>B. bifidum</i> + <i>L. casei</i>	0.95
<i>C. lanatus</i>	<i>L. acidophilus</i>	0.54
	<i>B. bifidum</i>	0.58
	<i>L. casei</i>	0.56
	<i>L. acidophilus</i> + <i>B. bifidum</i> + <i>L. casei</i>	0.94
<i>S. lycopersicum</i>	<i>L. acidophilus</i>	0.54
	<i>B. bifidum</i>	0.55
	<i>L. casei</i>	0.56
	<i>L. acidophilus</i> + <i>B. bifidum</i> + <i>L. casei</i>	0.94
<i>C. limetta</i>	<i>L. acidophilus</i>	0.5
	<i>B. bifidum</i>	0.56
	<i>L. casei</i>	0.57
	<i>L. acidophilus</i> + <i>B. bifidum</i> + <i>L. casei</i>	0.95
<i>P. guajava</i>	<i>L. acidophilus</i>	0.52
	<i>B. bifidum</i>	0.57
	<i>L. casei</i>	0.55
	<i>L. acidophilus</i> + <i>B. bifidum</i> + <i>L. casei</i>	0.95

Table 3: Titration acidity of the Synbiotic.

Conclusion

Pectin is already being used in microencapsulation of probiotics [11] in order to create more effective delivery systems for the probiotic bacteria. Further understanding of the protective mechanisms conferred by pectin will further enhance the numerous beneficial effects of probiotics. The results of the current study suggests that natural pectin from fruit sources can be used as an efficient prebiotic.

References

1. Laufenberg G, Kunz B, Nystroem M (2003) Transformation of vegetable waste into Value added (Product A) The Upgrading (Concept b) practical implementations. *Bioresource Technology* 87: 167-198.
2. Westerlund E, Aman P, Anderson R, Anderson RE, Rahman SMM (1991) Chemical characterization of water-soluble pectin in papaya fruit. *Carbohydrate Polymers* 15: 67-78.
3. Gibson GR, Collins MD (1999) Concept of balanced colonic microbiota, prebiotics and synbiotics. In: Hanson LA, Yolken RH (eds). *Probiotics other nutritional factors, and intestinal microflora* 42: 139-152.
4. Park SF, Kroll RG (1993) Expression of listeriolysin and phosphatidylinositol-specific phospholipase C is repressed by the plant-derived molecule cellobiose in *Listeria monocytogenes*. *Mol Microbiol* 8: 653-661.
5. Wang S, Chen F, Wu J, Wang Z, Liao X, et al. (2007) Optimization of pectin extraction assisted by microwave from apple pomace using response surface methodology. *J Food Eng* 78: 693-700.
6. Simmaky S, Jaanaki G (2014) Extraction and characterization of Pectin From Yellow Passion Fruit (*Passiflora edulis f. flavicarpa L.*) Endocarp Peel SAIMT Research Symposium on Engineering Advancements 2014 (SAITM-RSEA 2014). pp: 27-29.

-
7. Palframan RJ, Gibson GR, Rastall RA (2002) Effect of pH and dose on the growth of gut bacteria on prebiotic carbohydrates in vitro. *Anaerobe* 8: 287-292.
 8. Olano-Martin E, Rimbach GH, Gibson GR, Rastall RA (2003) Pectin and pectic-oligosaccharides induce apoptosis in in-vitro human colonic adenocarcinoma cells, *Anticancer Res* 23: 341-346.
 9. Urbiene S, Leskauskaitė D (2006) Formation of some organic acids during fermentation of milk. *Pol J Food Nutr Sci* 155: 277-281.
 10. Gomes AMP, Malcata FX (1999) Bifidobacterium spp. and Lactobacillus acidophilus: biological, biochemical, technological and therapeutical properties relevant for use as probiotics. *Trends in Food Science & Technology* 10: 139-157.
 11. Aquilah NS, Akhiar M (2010) Enhancement of probiotic survival by microencapsulation with alginate and prebiotics. *Basic Biotechnology* 6: 1-5.