

Effect of Heating and of Short Exposure to Sunlight on Carotenoids Content of Crude Palm Oil

Dongho Dongmo Fabrice Fabien¹, Ngono Ngane Annie¹, Demasse Mawamba Adélaïde¹, Schweigert Florian² and Gouado Inocent^{1*}

¹Department of Biochemistry, Faculty of Science, University of Douala, Douala, Cameroon

²Institute of Nutritional Science, University of Potsdam, Arthur-Scheuert-Allee 114-116, 14558 Bergholz-Rehbrücke, Germany

Abstract

This study was done to evaluate the effect of short exposure to sunlight and of heating on crude palm oil (CPO) quality particularly on its carotenoids content. Firstly, CPO was exposed to sunlight for 14 hrs. Then, samples were collected and kept at 4°C for analysis of carotenoids content, free fatty acid (FFA) and peroxide value (PV). Secondly, CPO was heated at 50°C, 120°C, 200°C or 400°C for 30 min, 60 min or 120 min and samples were cooled down and kept at 4°C until analysis of carotenoids content, FFA and PV. Finally, we studied the effect of heating of CPO in the food matrix (maize cake). Thus, maize cake was steamed on gas stove (100 ± 5°C) during exactly 1, 2, 3 or 4h and kept at 4°C until analysis of moisture and carotenoids contents. The results showed that short exposure to sunlight did not significantly affect carotenoids content, FFA and PV of CPO. However, heating accelerated the formation of peroxides and degradation of carotenoids. Destruction of carotenoids increased with both temperature and duration of exposure to heat. FFA did not significantly change during heating. Likewise, during heating of CPO in the food matrix, carotenoids content decreased significantly with cooking time. These results suggest that short exposure to sunlight does not have a significant effect on carotenoids content of CPO. But, its heating (directly or in the food matrix) results in significant degradation of carotenoids.

Keywords: Crude palm oil; Carotenoids; Heating; Sunlight; Photo-oxidation; Thermo-oxidation

Introduction

Vitamin A Deficiency (VAD) remains a major nutritional problem worldwide, despite substantial progress in recent decades [1-5]. It is estimated that more than 500,000 preschool children become blind each year because of the VAD among which 250,000 die because of resulting immune dysfunction [6]. The distribution of vitamin A (VA) capsules permitted good coverage of child populations at relatively low cost. However, it is widely recognized that food approaches should be part of means to fight against VAD in sustainable way. They must cover all approaches improving the nutrition focused on food systems or circuits taking into account nutrition education, production, marketing, processing and consumption [7]. In developing countries, these dietary approaches are very encouraging in the case of VA. Indeed, although provitamin A rich fruits and vegetables are not always available and accessible in these countries, crude palm oil (CPO) is generally available and cheap for everybody [6,8]. This oil extracted from the pulp of palm fruits (*Elaeis guineensis*) is the main source of provitamin A, principally β -carotene [8,9]. CPO has a special place among vegetable sources of VA, because of the absence of plant matrix and presence of lipid environment favorable to the absorption of carotenoids. Moreover, it provides to the organism energy, essential fatty acids, phytosterols, phosphatides and vitamin E [9].

The presence of unsaturated fatty acids exposes palm oil (PO) to lipid oxidation reactions. In the oxidation process, oxygen reacts with the double bonds of fatty acids to form peroxides and/or free radicals. These reactions result in the qualitative and nutritional alterations (rancidity, loss of essential fatty acids and vitamins) and even the toxicity caused by oxidative products (peroxide, aldehydes). They are influenced by extrinsic factors (temperature, light) and intrinsic factors (fatty acid composition, presence of pro- or antioxidants). Natural antioxidants found in many foods which can act as $1O_2$ quenchers help to reduce the oxidation reaction. So, a food will be more resistant to oxidation as its natural antioxidants content is high [10]. It is the case

of CPO which is very rich in carotenoids and vitamin E. However, the development of these reactions may be enhanced by cooking, refining or during storage [11]. In fact, some authors showed the negative effects of refining [12], heating [13-15] and light [14,16] on nutritional, sensory and/or physicochemical qualities of vegetable oils including carotenoids degradation.

In 2010, Cameroon was the 11th CPO producer in the world with 190 MT per year [17]. This oil is therefore the most accessible and available for consumption in this country. The survey on the CPO selling conditions and use in households in Douala town and its surroundings showed that sellers exposed oil to open air during the distribution. The maximum average exposure time to liquidate a sample of around 20l was 6 hrs. Moreover, in order to liquefy CPO (which coagulates at ambient temperature), sellers most often exposed it to sunlight and/or heated it slightly.

The survey also showed that housewives frequently used CPO to cook many foods. They usually used it heated either before cooking (bleaching) or during cooking. To bleach oil, they heated it either for long time at low temperature or for short time at high temperature. Moreover, those using the frying at commercial scale (doughnuts, chickens, fish etc.) declared that they always heated oil at high temperature for a long time and often recycled it. Depending on the food, CPO is added either at the beginning, during or at the end of the cooking. So, it results in dishes with oil distributed on the surface of

***Corresponding author:** Gouado Inocent, Department of Biochemistry, Faculty of Science, University of Douala, P.O. Box 24157 Douala-Cameroon, Tel: +237 99 83 30 65; E mail: gouadoi@yahoo.fr

Received February 13, 2014; **Accepted** March 24, 2014; **Published** April 03, 2014

Citation: Dongho Dongmo FF, Ngono Ngane A, Demasse Mawamba A, Schweigert F, Gouado I (2014) Effect of Heating and of Short Exposure to Sunlight on Carotenoids Content of Crude Palm Oil. J Food Process Technol 5: 314. doi:10.4172/2157-7110.1000314

Copyright: © 2014 Dongho Dongmo FF, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

food and those with oil uniformly (mixed with all) integrated to the matrix of food.

The objective of this study was to evaluate the effect of heating and of short exposure to sunlight on carotenoids from CPO. More specifically, we firstly assessed the effect of exposure to sunlight and of bleaching on carotenoids content and chemical quality (free fatty acids, peroxide value) of CPO. Secondly, we evaluated the retention of carotenoids from CPO during cooking of maize (*Zea mais*) cake, food commonly cooked in many households and eaten by many people in Cameroon.

Materials and Methods

Sampling

CPO used in this study was obtained in the area of *Moungo*, Littoral Region Cameroon. It was collected in the plastic cans of 20l and brought to Biochemistry Laboratory (University of Douala). Maize, wraps (banana leaves) and others ingredients were obtained in *Bonamoussadi* market, Douala Cameroon. We used white variety of maize because it was not only the most used by the housewives, but also the most available as compared to the yellow variety. Moreover, contrary to white variety, yellow variety contains carotenoids (around 0.4 mg/100 g flesh mater for dry seed). Thereby, these carotenoids could influence the study by interfering with carotenoids from CPO. Maize flour was obtained from maize seeds by crushing with grinder, then packed in plastic bags and brought in Laboratory. We firstly determined the composition of this flour and of Banana leaves used as wraps. We found that the flour contained around 12.4% of water; 70.7% of carbohydrates; 9.4% of proteins; 1.4% of fibers; 1.5% of ashes, 4.2% of lipids and 11.4µg of carotenoids per 100g of fresh mater. As for banana leaves, they contained around 87% of water; 12.6% of organic matter and 1.4% of ashes.

Methods

Three independent samples were done for each study of this work.

Study of the effect of exposure to sunlight: Three samples of 10l each of CPO contained in green plastic basins were exposed to sunlight during 14 hrs (7 hrs of exposure per day during 2 days) at 79359 ± 41628 Lux (1200-144000 Lux). Basins used had an interior diameter of 38 cm and a wall with 0.3 mm of thickness. Surface of oil directly exposed to sunlight was therefore 1134 cm² and expose volume to surface ratio was 8.82 ml/cm². Between the two exposition days, the containers were sealed and kept hideaway of light. In Parallel, a control was done without light (<2 Lux). Samples of 50ml were collected after 1, 2, 3, 5, 7, 9, 12 and 14 hrs. These samples were kept at 4°C until analysis of carotenoids content, free fatty acid (FFA) and peroxide value (PV).

Study of the effect of heating: For the study of direct heating effect (bleaching), 03 samples of 1l each of CPO contained in aluminum pot were heated on hotplate (CPC-250600D) at 50°C, 120°C, 200°C or 400°C during 30, 60 and 120 min for each temperature. This pot had an interior of 20 cm and a wall of 0.15 mm. The volume of oil heated to surface ratio was 3.19 ml/cm². The heated oil were allowed to cool down at room temperature and kept at 4°C until analysis of carotenoids content, FFA and PV.

As for the effect of heating in the food matrix (case of maize cake), given that the results from the effect of blanching could inform on the comportment of carotenoids during the cooking of meal in which CPO is distributed in surface of food, we chose to work with those where

CPO is mixed with the matrix of food. Furthermore, we chose to work with dishes cooked at long time and in which CPO is added at the beginning of cooking because the risk of carotenoids losses would be high there.

This is why we choose to work with maize cake. Maize cake was prepared according to the data collected close to housewives. Briefly, maize flour is mixed with water, salt and oil to form a paste. This paste is then packed and steamed with firewood or gas stove ($105 \pm 5^\circ\text{C}$). These data showed that the average time of cooking is around 2 hrs independently of the size of package. Nevertheless, some housewives affirmed that they preferred to cook for a short time while others preferred to cook for a long time. This is why we looked the effect of cooking time. Thus, around 4l of water was added to 5 kg of dry maize flour and mixed until obtaining a homogenous paste. This paste was watered with approximately 500 ml of CPO. Then, the paste was salted and mixed until the oil was absorbed completely. The portions of roughly 300 g of this paste were packed in the banana leaves. The packages obtained were steamed in an aluminum pot on gas stove. We carried out many tests by cooking the cake during exactly 1, 2, 3 and 4 hrs. After cooking, the samples were cooled down for the analysis of moisture and carotenoids contents. 500 ml of CPO and maize cake without oil were done under the same conditions as controls.

Analysis methods: Moisture content was analyzed by gravimetric method according to AOAC protocol [18]. Briefly, samples were dried until constant weight in an electric oven (Binder FDL 115) at 105°C. Moisture content was then calculated as percent of water loss.

Carotenoids content was evaluated by photometry (icheckTM Carotene; BioAnalyt GmbH, Teltow, Germany) at 446 nm. The procedure of analysis in CPO and in maize cake was different. Concerning CPO, a sample of 20 µl of oil was diluted with 2 ml of hexane. After vigorous shakeup, the mixture was read with photometer which gives the carotenoids content of solution in mg/l. The carotenoids content of CPO was calculated from this value by taking into account the dilution factor. As for maize cake, previous extraction of carotenoids was necessary. Thus, approximately 2 g of crushed food was added to a tube containing the mixture of ethanol/hexane (1:1). After centrifugation at 3000 revs/min for 5 min, the hexane phase was transferred to another tube. The procedure was repeated until the total discoloration of the residue. The volume of the hexane phases obtained was noted and 2 ml was used to read carotenoids content in photometer. The carotenoids content of maize cake in mg/100 g fresh mater was calculated from this value according to the mass of food used and the total volume of hexane phase obtained. As for carotenoids contents in mg/100 g dry mater, it was evaluated taking into account the moisture content of cake. Carotenoids losses were deducted from carotenoids contents in mg/100 g dry mater of food before and after cooking.

FFA content of oil was evaluated by titrimetric method according to AOAC protocol [18]. Briefly, 25 ml of ethanol was added to around 2 g of oil sample. The mixture was brought to boil in a water bath and then cooled down. After adding 2 drops of phenolphthalein as indicator, 0.1N NaOH was used to titrate the mixture with constant shaking for proper mixing until end-point (appearance of violet color). The free fatty acid content was calculated as followed, $\text{FFA} = V \times N \times M / 10 \times W$, where V=volume of NaOH, N=normality of NaOH (0.1N), M=molecular weight of palmitic acid (256 g/mol) and W=weight of the sample.

As for PV, it was analyzed by titrimetric method according to AOAC protocol [18]. Briefly, mixture of CH₃COOH/CHCl₃ (3:2) was added to 2 g of oil sample. Then, added 0.5 ml of saturated KI; agitated

during 1min and added 15 ml of distilled water and 0.5 ml of starch. Thereafter, this solution was titrated with 0.1N $\text{Na}_2\text{S}_2\text{O}_3$ until total disappearance of blue color. Meanwhile, we performed a blank test without oil. Peroxide value was calculated from the equation, $\text{PV (meq/Kg)} = 1000 \times (V_2 - V_1) \times N / M$, where M=mass of oil taken, V_2 =volume of $\text{Na}_2\text{S}_2\text{O}_3$ for essay, V_1 =volume of $\text{Na}_2\text{S}_2\text{O}_3$ for blank and N=normality of $\text{Na}_2\text{S}_2\text{O}_3$ (0.1N).

Data analysis: Data were processed using format designed in Microsoft Excel version 2010. Statistical analyzes were performed with Graph Pad Prism package version 5.00 (San Diego California USA). T-Student and one-way ANOVA (analysis of variance) tests were used to determine the influence of one factor. Two-way ANOVA was used to determine simultaneous influence of two factors and of their interaction. P-values were used as measure of significance and $P < 0.05$ was considered significant.

Results

Effect of sunlight on CPO

Sunlight did not have a significant effect on carotenoids content of CPO after 14 hrs of exposure (Table 1). Indeed, carotenoids content did not significantly vary during exposure and there was no significant difference between test and control (from 629.80 ± 39.00 mg/l to 629.63 ± 38.95 mg/l for test and from 629.80 ± 39.00 mg/l to 629.77 ± 38.95 mg/l for control).

Likewise, FFA did not vary significantly during exposure as well for test as for control (Table 2). In fact, FFA passed from $3.500 \pm 0.200\%$ to $3.540 \pm 0.200\%$ for test and from $3.500 \pm 0.200\%$ to $3.533 \pm 0.215\%$ for control.

Similarly, PV did not change significantly during exposure. But, contrary to carotenoids content and FFA, it varied more quickly in the test compared to the control. Indeed, the results showed a slight increase of PV in the test at the 2nd hour of exposure whereas it is only at the 5th hour that we noted this increase in the control. Moreover, after 14h of exposure, the PV increased of around 0.07 meq/kg in the test against only 0.02 meq/kg for the control. Although this result was not significant, it confirms that sunlight carries a pro-oxidizing activity on CPO (Table 3).

Effect of heating on CPO

For the study of the effect of direct heating, the results presented in Table 4 showed that bleaching significantly affected the carotenoids content of CPO. Statistical analysis showed that the degradation increased as well with temperature ($P < 0.0001$) as well as with time ($P < 0.0001$) of heating. There was a significant interaction ($P = 0.0224$) between the effect of the temperature and time of heating: at the higher temperature, the shorter time needed for carotenoids degradation.

Likewise, PV was significantly affected during bleaching (Table 5). Statistic showed that the formation of peroxides increased as well with temperature ($P < 0.0001$) as with time ($P < 0.0001$) of heating. There was a significant interaction ($P < 0.0001$) between the effect of the temperature and time of heating: at the higher temperature, the shorter time needed for peroxides formation. Nevertheless, let us note that the results also showed that at high temperatures (from 120°C), PV increased at the beginning of heating and decreased after certain heating time (from 60 min depending of temperature).

Contrary to carotenoids content and PV, FFA did not significantly vary during heating. Indeed, the result presented in Table 6 showed

that it neither significantly changed with heating time nor with heating temperature.

As for the study of the effect of heating in the food matrix, the results (Table 7) showed that carotenoids content of maize cake with CPO and of controls (CPO and maize cake without oil) significantly

		Carotenoids content (mg/l)		P ¹
		Test	Control	
Exposure time (h)	0	629.80 ± 39.00	629.80 ± 39.00	ns
	1	629.80 ± 39.00	629.80 ± 39.00	ns
	2	629.80 ± 39.00	629.80 ± 39.00	ns
	3	629.80 ± 39.00	629.80 ± 39.00	ns
	5	629.77 ± 38.95	629.80 ± 39.00	ns
	7	629.77 ± 38.95	629.80 ± 39.00	ns
	9	629.73 ± 39.00	629.80 ± 39.00	ns
	12	629.63 ± 38.95	629.80 ± 39.00	ns
	14	629.63 ± 38.95	629.77 ± 38.95	ns
P ²		ns	ns	

The values are given as mean ± standard deviation

¹Comparison of carotenoids content between assay and control at the same exposure time

²Comparison of carotenoids content between different exposure time

ns: No significant difference at $P < 0.05$

Table 1: Variation of carotenoids content of CPO (essay and control) according to exposure time.

		FFA (%)		P ¹
		Test	Control	
Exposure time (h)	0	3.500 ± 0.200	3.500 ± 0.200	ns
	1	3.500 ± 0.200	3.500 ± 0.200	ns
	2	3.500 ± 0.200	3.500 ± 0.200	ns
	3	3.503 ± 0.205	3.500 ± 0.200	ns
	5	3.503 ± 0.205	3.507 ± 0.200	ns
	7	3.510 ± 0.200	3.517 ± 0.220	ns
	9	3.513 ± 0.200	3.520 ± 0.220	ns
	12	3.520 ± 0.201	3.523 ± 0.225	ns
	14	3.540 ± 0.200	3.533 ± 0.215	ns
P ²		ns	ns	

The values are given as mean ± standard deviation

¹Comparison of FFA between essay and control at the same exposure time

²Comparison of FFA between different exposure time

ns: No significant difference at $P < 0.05$

Table 2: Variation of FFA of CPO (assay and control) according to exposure time.

		PV (meq/kg)		P ¹
		Test	Control	
Exposure time (h)	0	1.167 ± 0.252	1.167 ± 0.252	ns
	1	1.167 ± 0.252	1.167 ± 0.252	ns
	2	1.170 ± 0.252	1.167 ± 0.252	ns
	3	1.173 ± 0.257	1.167 ± 0.252	ns
	5	1.180 ± 0.256	1.170 ± 0.252	ns
	7	1.183 ± 0.261	1.173 ± 0.252	ns
	9	1.200 ± 0.271	1.173 ± 0.257	ns
	12	1.207 ± 0.276	1.177 ± 0.257	ns
	14	1.237 ± 0.235	1.183 ± 0.278	ns
P ²		ns	ns	

The values are given as mean ± standard deviation

¹Comparison of PV between essay and control at the same exposure time

²Comparison of PV between different exposure time

ns: No significant difference at $P < 0.05$

Table 3: Variation of PV of CPO (essay and control) according to exposure time.

Carotenoids content (mg/l)		Heating temperature (°C)				P ¹
		50	120	200	400	
Heating time (h)	0	626.60 ± 2.60 ^{a,‡}	626.60 ± 2.60 ^{a,‡}	626.60 ± 2.60 ^{a,‡}	626.60 ± 2.60 ^{a,‡}	ns
	0.5	621.40 ± 2.60 ^{a,‡}	613.60 ± 5.20 ^{a,‡}	478.40 ± 15.60 ^{b,‡}	140.40 ± 23.40 ^{c,‡}	<0.0001
	1	616.20 ± 2.60 ^{a,‡}	542.10 ± 3.90 ^{b,‡}	457.60 ± 15.60 ^{c,†}	87.75 ± 21.45 ^{d,†}	<0.0001
	2	613.60 ± 5.20 ^{a,‡}	368.94 ± 13.26 ^{b,†}	297.96 ± 28.60 ^{c,‡}	78.52 ± 20.28 ^{d,‡}	<0.0001
P ²		0.0074	<0.0001	<0.0001	<0.0001	

The values are given as mean ± standard deviation

The values of the same line superscripted with the same letter are not significantly different at P<0.05

The values of the same column superscripted with the sign #, † or ‡ are not significantly different at P<0.05

¹Comparison of carotenoids content between different heating temperatures at the same heating time

²Comparison of carotenoids content between different heating times at the same heating temperature

ns: No significant difference at P<0.05

Table 4: Variation of carotenoids content of heated CPO according to the time and the temperature of heating.

PV (meq/kg)		Heating temperature (°C)				P ¹
		50	120	200	400	
Heating time (h)	0	0.983 ± 0.104 ^{a,‡}	0.983 ± 0.104 ^{a,‡}	0.983 ± 0.104 ^{a,‡}	0.983 ± 0.104 ^{a,‡}	ns
	0.5	1.013 ± 0.105 ^{a,‡}	1.260 ± 0.078 ^{a,‡}	2.577 ± 0.585 ^{b,‡}	4.387 ± 0.361 ^{c,‡}	<0.0001
	1	1.050 ± 0.118 ^{a,‡}	1.390 ± 0.017 ^{a,†}	2.743 ± 0.630 ^{b,‡}	4.240 ± 0.246 ^{c,‡}	<0.0001
	2	1.110 ± 0.115 ^{a,‡}	1.363 ± 0.015 ^{a,†}	1.997 ± 0.105 ^{d,†}	4.033 ± 0.153 ^{c,‡}	<0.0001
P ²		ns	0.0002	0.0045	<0.0001	

The values are given as mean ± standard deviation

The values of the same line superscripted with the same letter are not significantly different at P<0.05

The values of the same column superscripted with the sign #, † or ‡ are not significantly different at P<0.05

¹Comparison of PV between different heating temperatures at the same heating time

²Comparison of PV between different heating time at the same heating temperature

ns: No significant difference at P<0.05

Table 5: Variation of PV of heated CPO according to the time and the temperature of heating.

FFA (%)		Heating temperature (°C)				P ¹
		50	120	200	400	
Heating time (h)	0	3.567 ± 0.379	3.567 ± 0.379	3.567 ± 0.379	3.567 ± 0.379	ns
	0.5	3.587 ± 0.405	3.583 ± 0.390	3.597 ± 0.385	3.590 ± 0.403	ns
	1	3.597 ± 0.422	3.610 ± 0.419	3.627 ± 0.410	3.623 ± 0.414	ns
	2	3.660 ± 0.478	3.620 ± 0.419	3.650 ± 0.407	3.637 ± 0.411	ns
P ²		ns	ns	ns	ns	

The values are given as mean ± standard deviation

¹Comparison of FFA between different heating temperatures at the same heating time

²Comparison of FFA between different heating time at the same heating temperature

ns: No significant difference at P<0.05

Table 6: Variation of FFA of heated CPO according to the time and the temperature of heating.

	Before cooking	After cooking				P
		1h	2h	3h	4h	
CONTROLS						
Crude palm oil						
mg/l	823 ± 74 ^a	738 ± 77 ^b	681 ± 54 ^c	566 ± 55 ^d	502 ± 81 ^e	<0.0001
Maize cake without oil						
µg/100 g of fresh mater	6.27 ± 0.75 ^a	5.50 ± 0.70 ^b	3.73 ± 0.55 ^c	3.50 ± 0.53 ^d	3.13 ± 0.51 ^d	<0.0001
µg/100 g of dry mater	18.53 ± 2.80 ^a	14.43 ± 2.32 ^b	8.67 ± 0.97 ^c	8.17 ± 0.68 ^c	6.70 ± 1.18 ^c	<0.0001
ASSAY (Maize cake with oil)						
mg/100 g of fresh mater	2.00 ± 0.57 ^a	1.90 ± 0.52 ^{a,b}	1.80 ± 0.52 ^{b,c}	1.40 ± 0.36 ^{c,d}	1.20 ± 0.26 ^d	<0.0001
mg/100 g of dry mater	5.80 ± 1.5 ^a	4.90 ± 1.40 ^{a,b}	4.20 ± 1.30 ^{b,c}	3.50 ± 0.93 ^{c,d}	2.80 ± 0.67 ^d	<0.0001


The values are given as mean ± standard deviation

The values of the same line superscripted with the same letter are not significantly different at P<0.05

Table 7: Variation of carotenoids content of maize cake with CPO and of controls according to the cooking time.

decreased (P<0.0001) during cooking. Figure 1 gives the variations of carotenoids losses in the maize cake with CPO and in the controls. It shows that carotenoids losses significantly increased (P<0.0001) with cooking time as well for maize cake with CPO as for controls. In fact, these losses passed from 15.33 ± 3.75% after 1 hrs of cooking to 51.00 ±

3.00% after 4 hrs in maize cake with CPO; from 22.33 ± 3.06% to 65.67 ± 4.73% in maize cake without CPO and from 10.33 ± 1.51% to 39.33 ± 4.93% in CPO. Moreover, it was found that whatever the cooking time, the carotenoids losses were significantly lower in CPO compared to maize cake. Likewise, they were significantly lower in maize cake



Vertical bars represent standard deviation; for the same cooking time, values with different letters are significantly different at $p < 0.05$

Figure 1: Variation of carotenoids losses in maize cake with CPO and in controls (CPO and maize cake without oil) according to the cooking time.

with CPO compared to maize cake without oil. By example, after 1h of cooking, we noted $10.33 \pm 1.51\%$ of losses in CPO, $15.33 \pm 3.75\%$ in maize cake with CPO and $22.33 \pm 3.06\%$ in maize cake without oil. After 4 hrs of cooking, we noted $39.33 \pm 4.93\%$; $51.00 \pm 3.00\%$ and $65.67 \pm 4.73\%$ of losses respectively in CPO, cake with CPO and cake without oil.

Discussion

The objective of this work was to evaluate the effect of sunlight and of heating on the CPO quality particularly on its carotenoids concentration. We noted that short exposure to sunlight did not significantly affect carotenoids content, FFA and PV of CPO. This result can be partially explained by the exposure time which was very short. In fact, some study showed that it is after five day of exposure to sunlight that one could observe a significant variation of physico-chemical parameters of vegetable oil [16]. Besides, another authors noted that FFA and PV significantly varied in CPO only after two weeks of storage [19]. This result could be explained by the distance of incident radiations on exposed oil and indirectly by the radiations intensity. Indeed, formal studies showed that the exposure of CPO on red, green or blue light (that incidence distance is short compared to sunlight) led to high increase of FFA compared to exposure to sunlight [20]. Moreover, other authors remarked that exposure of olive oil on UV light and on white light resulted to high reduction of β -carotene and high increase of FFA and PV compared to exposure to sunlight [14]. One could explain this result by high content and diversity of natural antioxidants of CPO (carotenoids, tocopherol and tocotrienol). This hypothesis is supported by the results of the study of Zeb et al. [14] who noted, contrary to our study, that exposure of

olive oil to sunlight during 14 hrs resulted to a significant decrease of β -carotene and high increase of PV. This contradictory result could be explained by the fact that antioxidants content of olive oil is lower compared to CPO. Moreover, Schroeder et al. [21] showed the synergic mechanism of antioxidants during frying with CPO.

Study of the heating effect on CPO was done in two parts. The first part on the effect of direct heating (bleaching) and the second part on the effect of heating in the matrix of food (case of maize cake). We noted that direct heating resulted to significant decrease of carotenoids content and significant increase of PV. The reduction of carotenoids could be attributed directly to the effect of heating or indirectly to the effect of reactive oxygen species (singlet oxygen, superoxide radical, hydroxyl radical) issued from lipids oxidation. In fact, these compounds react with polyene chain of carotenoids to generate carotenoids-adducts radicals [22,23]. This work confirms the synergic effect of time and temperature of heating on the lipids oxidation. These results corroborate with those of Ahmed et al. [13] who had the same observations during heating of red palm olein. Likewise, Zeb et al. [14] showed a significant decrease of β -carotene and a significant increase of PV in the olive oil samples heated at 72°C compared to those stored at 4°C or at ambient temperature (25°C). As Ahmed et al. [13], we observed that at high temperature, PV increased at the beginning of heating and decreased at high heating time. In fact, the primary products of lipid oxidation are hydroperoxides, therefore the result of PV give a clear indication of oxidation. The reduction of PV at elevated temperature could be attributed to the rapid decomposition of hydroperoxides to secondary oxidation product [13]. Opposing to carotenoids content and PV, we noted that heating did not significantly affect FFA. This result is in accordance with those of Ahmed et al. [13]

and Okiy and Oke [24] who had similar observations. Some authors noted a significant increasing of FFA of oils with time and temperature during frying [25]. The difference between the two studies is due to the fact that in the present study there was no frying process. Frying food results in increasing of the water content of the oil and subsequently increases the hydrolysis and FFA content. The bleaching of CPO carried on a significant oxidizing activity (proportional to heating time and temperature) resulting not only to the reduction of carotenoids content, but also to the formation of dangerous compounds (hydroperoxides) which may react with cellular components such as proteins, nucleic acids, lipids, carbohydrates and poly-unsaturated acids [26,27].

The results of heating of CPO in the matrix of foods on the carotenoids content are in accordance with the observations done during the direct heating. In fact, during cooking one noted a significant diminution of carotenoids content of maize cake with CPO and of controls (maize cake without oil and CPO) according to cooking time. It was remarkably noted that independently of cooking time, carotenoids losses in maize cake with CPO were significantly lower compared to the control maize without oil and higher compared to the control CPO. This result agrees with the study of Ahmed et al. [28] who showed that during storage, cakes made with palm oil resisted better to oxidation reactions compared to those made with ordinary margarine. As previously, this could be explained by high content and diversity of natural antioxidants of CPO (carotenoids, tocopherol and tocotrienol).

Conclusions and Recommendations

We can conclude that although sunlight does not have a significant effect on carotenoids content of CPO at the short term, it carries on a small pro-oxidant activity. Contrary, heating of CPO carried on a significant oxidizing activity (proportional to heating time and temperature) resulting not only to the reduction of carotenoids content, but also to the formation of hydroperoxides which are dangerous compounds. Likewise, the heating of CPO in the matrix of food results to significant reduction (proportional to heating time) of carotenoids content. Thus, we recommend to traders to avoid exposing CPO in open air during distribution. To housewives, we recommend to avoid bleaching of CPO, to cook foods requiring uncolored oils, they could use less colored oils (cotton oil, olive oil, soya oil, groundnut oil or refined palm oil) or blend them with CPO. Also they could develop the strategies to reduce maximally carotenoids losses during cooking of foods in which CPO is mixed with the matrix of food by varying for example some cooking parameters as cooking time, quantity of oil used or kind of package.

Acknowledgements

The authors acknowledge BioAnalyt GmbH (Teltow, Germany) for gracefully provided us the photometer (ichekTM carotene) for carotenoids analysis.

References

1. Kollo B, De Bernadi R, Sibetcheu D, Nankap M, Tata Ngoh J, et al. (2001) Report of National survey on vitamin A deficiency and anemia in Cameroon. Edit. MSP, UNICEF, WHO, HK World Wide, Sigh and Life.
2. Strobel M, Tinz J, Biesalski HK (2007) The importance of β -carotene as a source of vitamin A with special regard to pregnant and breast feeding women. *Eur J Nutr* 46: 1-20.
3. Gouado I, Demasse MA, Meyimgo OSR, Touridomon SI, Tchouanguép MF (2008) Provitamin A carotenoid content of dried fermented cassava flour: The effect of palm oil addition during processing. *Int J Food Eng* 4: 4.
4. WHO (2009) Global prevalence of vitamin A deficiency in populations at risk 1995–2005. WHO Global Database on Vitamin A deficiency. Geneva, World Health Organization.
5. Ministry of public health in Cameroon, Helen Keller International Cameroon, UNICEF (2011) National survey of micronutrient status and consumption of fortifiable foods. Yaoundé (Cameroon): Helen Keller International Cameroon.
6. Edem DO (2009) Vitamin A: A review. *Asian J Clin Nutr* 1: 65-82.
7. Delisle H, Zagré N, Bakari S, Codjia P, Zandong R (2003) A food system approach to Vitamin A deficiency. *Food Nutr Agric*. 32: 40-50.
8. Zeb A, Mehmood S (2004) Carotenoids contents from various sources and their potential health applications. *Pakistan J Nutr*. 3: 199-204.
9. Mukherjee S, Mitra A (2009) Health effects of palm oils. *J Hum Ecol* 26: 197-203.
10. Cillard J, Cillard P (2006) Mécanismes de la peroxydation lipidique et des antioxydants. *Oléagineux, Corps Gras, Lipides* 13: 24-29.
11. Gary RL, Tong W, Vijai KSS (2005) Storage, handling, and transport of oils and fats. *Bailey's Industrial Oil and Fat Products*, (6thedn) Fereidoon Shahidi, John Wiley & Sons, Inc., USA.
12. Verhé R, Verleyen T, Hoed VV, Greyt WD (2006) Influence of refining of vegetable oils on minor components. *J Oil Palm Res Special issue-April 2006*:168-179.
13. Ahmed AS, Abdulah A, Aini IN (2006a) Changes of β -carotene content during heating of red palm olein. *J Oil Palm Res. Special issue-April 2006*: 99-102.
14. Zeb A, Khan S, Khan I, Imbran M (2008) Effect of temperature, UV, sun and white lights on the stability of olive oil. *J Chem Soc Pak*. 30: 790-794.
15. Combe N, Rossignol-Castera A (2010) Vegetable oils and frying. *Cahier Nutr Diet* 45: 44-51.
16. Fekarurhobo GK, Obomanu FG, Maduelosi (2009) Effects of short-term exposure to sunlight on the quality of some edible vegetable oils. *Res J App Sc*. 4: 152-156.
17. Oil World (2011) Palm Oil: Statistics 2011. Oil World report 2011.
18. AOAC (1980) Official methods of analysis (11thedn) WILLIAM HORWITZ edv, Washington DC, USA.
19. Ngando EGF, Mpondo MEA, Dikotto EEL, Koona P (2011) Assessment of the quality of crude palm oil from smallholders in Cameroon. *J Stored Products Postharvest Res*. 2: 52-58.
20. Oyem HH (2011) Monitoring the free fatty acid level of crude palm oil stored under light of different wavelengths. *Am J Food Tech*. 6: 701-704.
21. Schroeder MT, Becker EM, Skibsted LH (2006) Molecular mechanism of antioxidant synergism of tocotrienols and carotenoids in palm oil. *J Agric Food Chem*. 54: 3445–3453.
22. Mezgebe AG (2011) The relation between the in vitro bioaccessibility of β -carotene and structural characteristics of carrot pieces during thermal processing (sterilization and pasteurization) in a retort. *Afr J Food Sci*. 5: 845-850.
23. Bonnie TP, Choo YM (1999) Oxidation and thermal degradation of carotenoids. *J Oil Palm Res*. 2: 62-78.
24. Okiy DA, OKE OL (1981) Chemical changes in palm oil at the early stages of heating. *J Nigerian Inst Oil Palm Res* 6: 101-113.
25. Warner K, MOUNTS TL (1993) Frying stability of soybean and canola oils with modified fatty acids composition. *J Amer Oil Chem Soc* 70: 983-988.
26. Isong EU, Ebong PE, Ifon ET, Umoh IB, Eka OU (1997) Thermoxidized Palm Oil Induces reproductive Toxicity in Healthy and Malnourished Rats. *Plant Foods Hum Nutr*. 51: 159-166.
27. Yaeko I, Shunichi Y, Mitsuru U (1984) Effect of ingestion of thermally oxidized frying oil on peroxidative criteria in rats. *Lipids* 19:324-331.
28. Ahmed AS, Abdulah A, Aini IN, Gapor ABMD (2006) Evaluation of storage stability of yellow cake made with red palm fat. *J Oil Palm Res. Special issue-April 2006*: 81-86.

Citation: Dongho Dongmo FF, Annie NN, Adélaïde DM, Florian S, Inocent G (2014) Effect of Heating and of Short Exposure to Sunlight on Carotenoids Content of Crude Palm Oil. *J Food Process Technol* 5: 314. doi:[10.4172/2157-7110.1000314](https://doi.org/10.4172/2157-7110.1000314)