

Effect of Organic and Inorganic Fertilizers on Natural Food Composition and Performance of African Catfish (*Clarias gariepinus*) Fry Produced Under Artificial Propagation

Sebastian S Mosha^{1*}, Jeremiah Kang'ombe², Wilson Jere² and Nazael Madalla³

¹Training Department, Ministry of Agriculture Training Institute (MATI), P.O Box 121, Mtwara, Tanzania

²Department of Aquaculture and Fisheries Science, Lilongwe University of Agriculture and Natural Resource (LUANAR), P.O Box 219, Lilongwe, Malawi

³Department of Animal Science and Production, Sokoine University of Agriculture (SUA), P.O Box 3004, Morogoro, Tanzania

Abstract

Two experiments were conducted to assess the effect of organic and inorganic fertilizers on natural food composition and performance of African catfish (*Clarias gariepinus*) fry produced under artificial propagation. In both experiments chicken manure, di-ammonium phosphate and no fertilizer were used as treatments. The first experiment was conducted to determine the abundance and diversity of natural food produced in tanks fertilized with chicken manure and di-ammonium phosphate fertilizer. The second experiment was conducted to assess growth performance and survival of catfish fry produced in chicken manure and di-ammonium phosphate at 5fry/m² and 10fry/m² stocking densities. Nine concrete tanks and eighteen concrete tanks were used in experiment 1 and 2, respectively. The results indicated that abundance of natural food (*phytoplankton*) were significantly higher ($P < 0.05$) in di-ammonium phosphate (DAP) fertilizer applied tanks compared to other treatments. *Zooplankton* diversity was higher in tanks applied with chicken manure, followed by tanks applied with DAP fertilizer and least in tanks with no fertilizer. Fry growth performance was higher in fertilized tanks compared to unfertilized tanks. At low stocking density (5fry/m²) fry had better growth performance compared to high stocking density (10fry/m²) across all fertilizer types. Survival rates were not significantly different ($P > 0.05$) between chicken manure and DAP fertilized treatments across stocking densities though significantly differed ($P < 0.05$) from the control. Water quality parameters were found to be within the optimum range for cat fish in both experiments. In conclusion, the study indicated that higher *phytoplankton* abundance are attained with DAP fertilized tanks and *zooplankton* diversity were higher in chicken manure applied tanks. Therefore, it is recommended that for better growth and survival in aquaculture practices, catfish fry should be raised in DAP or chicken manure fertilized tanks at low stocking density as suggested by this study.

Keywords: DAP fertilizer; Chicken manure; Catfish fry; Stocking density; Fry performance

Introduction

In recent times, the African catfish, *Clarias gariepinus*, has gained popularity in aquaculture sector of Tanzania. However, the rearing of *C. gariepinus* larvae to juveniles has proved to be challenging due to their small size and lack of functional digestive system [1,2]. Great losses are in the hatchery, as fry weans over from yolk absorption to exogenous feeding [3]. This is due to inability to accept large sized feeds and assimilate protein from dry formulated diets [1,4]. According to Agadjihouede et al. [5], *Artemia* constitutes an excellent starting food in larviculture of *C. gariepinus*. However, its cost is high and not available in developing countries especially in the rural fish farming [6]. Due to this fact, it has been found important to provide the larvae with live feeds such as *zooplankton* or algae first before they are sequentially acclimatized to accepting formulated diets [2,7].

Primary production in fish pond/tank is limited by phosphate, in particular, and nitrogen in general [8]. Fertilizer application stimulates the growth of decomposers such as bacteria and fungi which breakdown toxic waste products that can accumulate with the use of prepared feeds [9]. It is an inexpensive method of feeding and plankton is necessary for smaller fish which are small to eat supplemental feeds [10]. Ponds/tanks with fertilized water will turn a rich green or reddish color when the plankton becomes abundant and food will be available for the fish [11].

Survival and growth of catfish larvae and fry are influenced by several factors such as stocking density [12,13] and water quality [14,15]. Growth of juvenile African catfish is directly density dependent and is normally highly aggressive when confined in small numbers in a large volume of water [16,17]. Aggressiveness, territorial defense and

development of hierarchies and individual dominance are often reduced at high stocking density and fish may start to be stressed once certain threshold densities are attained [18,19]. Aggression can result in stock losses, reduced food conversion efficiency and slower growth. So, there must be a density from which both growth and production decreased with increasing stocking density [17]. Water quality, mainly dissolved oxygen and pH levels are considered as the limiting factors in intensive fish culture. According to Brazil and Wolters [14]; Pangni et al. [15], growth increase in channel catfish larvae reared in tanks was due to high levels of dissolved oxygen. The decreasing trend of dissolved oxygen in tanks with high stocking densities would be attributed to the gradual increase in biomass, resulting in higher oxygen consumption at varied stocking densities [15].

Among fertilizers used to increase natural food in ponds and tanks, chicken manure and di-ammonium phosphate (DAP) fertilizers are very cheap and locally available to fish farmers [20]. They contain a good combination of nitrogen and phosphorus in different proportions which increase the quantity of primary producers [21,22]. According to Lin et

*Corresponding author: Sebastian S Mosha, Training Department, Ministry of Agriculture Training Institute (MATI), P.O Box 121, Mtwara, Tanzania, Tel: 255-22-2780077; E-mail: sebylm2012@yahoo.com

Received July 17, 2016; Accepted August 26, 2016; Published August 28, 2016

Citation: Mosha SS, Kang'ombe J, Jere W, Madalla N (2016) Effect of Organic and Inorganic Fertilizers on Natural Food Composition and Performance of African Catfish (*Clarias gariepinus*) Fry Produced Under Artificial Propagation. J Aquac Res Development 7: 441. doi: [10.4172/2155-9546.1000441](https://doi.org/10.4172/2155-9546.1000441)

Copyright: © 2016 Mosha SS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

al. [23], fertilization in fish ponds/tanks is known worldwide to improve pond productivity by promoting the growth of *phytoplankton* thereby increasing natural food available to fish. A lot of progress has been made so far in aquaculture, however, larvae rearing remains the bottleneck in *C. gariepinus* production. The use of natural live food at earlier stage of fry development seems to be one of the solutions to improve growth performance and survival. Therefore, the study was designed to assess the growth performance and survival rate of the fry reared in chicken manure and DAP applied tanks at different stocking density.

Materials and Methods

Experimental design

The study was conducted from the last week of March 2015 to June 2015 at aquaculture research facility in Magadu farm at Sokoine University of Agriculture (SUA) Morogoro, Tanzania. Two experiments were conducted for 1 week and 2 months, respectively. Experiment 1, involved determination of abundance and diversity of natural food in tanks applied with different type of fertilizer. While Experiment 2, involved evaluation of growth performance and survival of African catfish fry stocked under two stocking densities (5fry/m² and 10fry/m²), in tanks applied with different type of fertilizer [24].

Experiment 1: Determination of abundance and diversity of natural food

(a) Fertilizer application: Fresh chicken manure from layers kept in the cages were collected from poultry unit belonging to Department of Animal Science and Production, SUA [25]. Nitrogen (2.55%) and Phosphorus (0.95%) in chicken manure was determined using proximate analysis at Animal Science laboratory according to Association of Official Analytical Chemists [26]. Di-ammonium phosphate fertilizer was purchased from a local agro-input shop in Morogoro containing 18% N and 46% P as indicated on the package [27]. Several trials were conducted to find the appropriate rate of fertilizers application which produced algae bloom in the tanks before starting the experiment. Nine concrete tanks having a area of 7 m² each were laid out with 10 cm of soil and then filled with water to depth of 0.8 m. Three treatments: 2 kg of chicken manure, 42 g of di-ammonium phosphate fertilizer and no fertilizer (control) were randomly assigned to the tanks and replicated three times.

(b) Water sampling for plankton analysis: One week after fertilization water samples were collected with 10 L bucket from 4 locations within each concrete tank and this was done for all 9 tanks at 08:00 am. Water sample was placed into plankton net (KC Denmark A/S Denmark) with a 20 µm mesh size and left for 20 min. The concentrated sample from plankton net was transferred to 200 ml plastic bottles. Five drops of 70% alcohol were added to each sample to fix the organisms and taken to Faculty of Veterinary Medicine Laboratory (SUA) for analysis.

(c) Plankton identification and quantification: Plankton identification was done on a light microscope (BRESENER, Germany) 100X magnification using identification keys according to UNESCO [28]. 1 ml of water sample was taken from the collecting bottles (200 ml bottles) using micropipette and transferred to a Sedgewick Rafter cell (Wild Supply Company, England) then covered with slide at the top and placed under microscope. From 10 randomly selected squares of cell, planktonic organisms were enumerated and numerical abundance was calculated. *Phytoplankton* and *Zooplankton* abundance were calculated using the following formulas as described by Greenberg et al. [29] and Wetzel and Likens [30], respectively:

$$\text{Phytoplankton Abundance} = \frac{C}{F \times V}$$

Where C is the number of organisms counted, F is the number of fields counted and V is the volume of sample settled.

$$\text{Zooplankton} / L = \frac{C \times V_a}{V_b \times V_c}$$

Where C is the number of *zooplankton* counted, V_a is the volume of the concentrated sample (L), V_b is the volume of counted sample (L) and V_c is the volume of water filtered (L).

In case of plankton diversity Shannon-Wiener diversity index (H') and evenness (J') were used. Diversity index of plankton was calculated by using the formula as described by Krebs [31]:

$$H' = \frac{n \ln(n) - \sum_{i=1}^k f_i \ln(f_i)}{n}$$

Where k is the number of categories, f_i is the number of observations in category i, n is the sample size.

Species evenness or homogeneity or relative diversity (J') was calculated from the observed species diversity and from the equation of H_{max} as described by Sundar et al. [32]. Index of species evenness was measured by using the following formula:

$$J' = \frac{H'}{H'_{\max}}$$

Where, H'_{max} = ln(k)

After plankton quantitative analysis, *phytoplankton* and *zooplankton* were further identified up to genus level following the guide of Bellinger [33].

Experiment 2: Determination of growth and survival of African catfish (*C. gariepinus*) fry

Fertilization of tanks in this experiment was done one week before fry stocking. Total 18 concrete tanks (7 m² each) were used. Six tanks were fertilized with chicken manure and the other 6 tanks fertilized with diammonium phosphate at a rate of 286 g/m² and 6 g/m², respectively. While the remaining 6 tanks were not fertilized (control). The experiment was laid out in a Factorial Design where catfish fry assigned randomly in 18 concrete tanks. *Clarias gariepinus* fry (1 month old) an average weight of 0.5 g ± 0.01 g from hatchery tank were randomly assigned to treatment tanks. Two stocking densities (5fry/m² and 10fry/m²) were used. Each stocking density was assigned at once and replicated three times. All 18 concrete tanks were covered with plastic nets to prevent predators. Fertilization was repeated every week for a period of 2 months. During this period, all treatments were supplemented with a formulated diet containing 55% crude protein and fed at 25% body weight per day.

(a) Data collection: Once every week in the rearing period of 2 months, the catfish fry from each concrete tank were harvested by using seine net. Then placed in 10L buckets and taken to hatchery building where they were weighed by using analytical balance (BOECO, Germany) and recorded. The following formula was used to determine fry performance and survival rate as outlined by Kang'ombe et al. [34].

Weight gain (g) = Final mean weight (g) - Initial mean weight (g)

$$\text{Increase mean weight}(\%) = \frac{\text{Final mean weight}(\text{g}) - \text{Initial mean weight}(\text{g})}{\text{Initial mean weight}(\text{g})} \times 100$$

$$\text{Specific growth rate (\%)} = \frac{\ln(\text{Final mean weight (g)} - \text{Initial mean weight (g)})}{\text{Time (days)}} \times 100$$

$$\text{Survival rate (\%)} = \frac{\text{Initial number of fry} - \text{Number of dead fry}}{\text{Initial number of fry}} \times 100$$

(b) Physico-chemical parameters of water: In both experiments, water quality parameters were monitored. Temperature (°C) and dissolved oxygen (mg L⁻¹) were measured on daily basis (08:00-16:00hrs) using a multi-probe water checker (OAKTON, Singapore). The pH was measured every day at 8:00 hrs using pH meter (EXSTICK, USA). Ammonia and nitrite (ppm) were calculated once a week by using Salicylate Method and Diazotization Method, respectively.

(c) Statistical data analysis: One way and two way analysis of variance (ANOVA) (Experiments 1, 2 respectively) were used to compare differences between treatment means at 5% level of significance. Post-hoc analysis was done where significant differences existed between treatments means using Tukey's Test. Analyses were performed using SPSS software version 20 [35].

Results

Abundance and diversity of different species of live food produced in chicken manure and di-ammonia phosphate (DAP) fertilizers

Generally, fertilized tanks had higher abundance and diversity of *phytoplankton* and *zooplankton* compared to unfertilized tanks (Tables 1 and 2). *Phytoplankton* abundance was represented by four diverse groups, namely *chlorophyta*, *cynophyta*, *euglenophyta* and *diatomeae* [36]. Among the *phytoplankton*, *chlorophyta*, *cryophyta* and *diatomeae* were highly significant (P<0.05) in tanks fertilized with chicken manure and DAP fertilizer compared to control. *Euglenophyta* showed to be highly significant (P<0.05) when DAP fertilizer was used in comparison to other treatments (Table 1). *Phytoplankton* diversity was higher in tanks applied with DAP fertilizer, followed by tanks applied with chicken manure and least in tanks with no fertilizer. The diversity was indicated with the high values of Shannon-Wiener indices (H') and evenness (J'). *Zooplankton* was represented by major three groups, namely rotifers, copepods and cladocerans [37]. All *zooplankton* groups had significantly (P<0.05) higher abundance in tanks fertilized with DAP fertilizer as compared to chicken manure and control. Chicken manure had high values of Shannon-Wiener indices (H') and evenness (J') compared to other treatments. This indicated that *zooplankton* diversity was higher in tanks applied with chicken manure compared to other treatments (Table 2).

Effect of fertilizers type and stocking density on growth performance of (*C. gariepinus*) fry

Fry performance was higher in fertilized tanks compared to unfertilized tanks. At low stocking density (5fry/m²) fry had better growth performance compared to high stocking density (10fry/m²) across all fertilizer types (Figure 1). Individual final mean weights (g), weight gain (g) and specific growth rate (%) were significantly different (P<0.05) between stocking densities. While were not significantly different (P>0.05) between chicken manure and di-ammonium phosphate (DAP) fertilizers but significant higher than the control. Initial mean weight (g) and increase mean weight (g) were not significantly different (P>0.05) across stocking densities and fertilizers type (Table 3).

Effect of fertilizers type and stocking density on survival of (*C. gariepinus*) fry

Survival rate was significantly (P<0.05) higher under stocking density and the interaction but were not significantly different among fertilizer types. Higher survival rate observed in tanks fertilized with DAP fertilizer and control tanks under low stocking density (Table 3).

Effect of fertilizers type and stocking density on water quality parameters

There were no significant differences (P>0.05) in morning and afternoon temperature, dissolved oxygen (morning), and ammonia between stocking density and fertilizer type. Afternoon dissolved oxygen and nitrite were significantly (P<0.05) higher among stocking densities and fertilizer types. The range in pH values were similar to all stocking densities under chicken manure and DAP fertilizer while under control different ranges were observed (Table 4).

Discussion

The results indicated that abundance of plankton was higher in tanks fertilized with DAP fertilizer than in chicken manure and the control. Similar results have been reported by Gamal [38], Padmavathi [39] and Kumar et al. [22]. The significantly higher abundance and diversity of *phytoplankton* is likely due to easily available of nutrients to the water. Boyd and Massaut [40] reported that inorganic fertilizers have much higher concentrations of nutrients such as nitrogenous compounds than manures.

Rotifer, copepods and cladocerans showed a higher abundance as *phytoplankton* trend. This might be due to feeding effects of *zooplankton* on *phytoplankton* [41,42]. Similarly, Ghosh et al. [43]; Kumar et al. [22]

Treatments	Phytoplankton				Zooplankton		
	Chlorophyta	Cynophyta	Euglenophyta	Diatomeae	Rotifers	Copepods	Cladocerans
Chicken manure	52750 ± 10528 ^a	18833 ± 3346 ^a	9917 ± 1109 ^a	10375 ± 2376 ^a	157 ± 23 ^a	257 ± 8 ^a	167 ± 14 ^a
DAP fertilizer	71833 ± 11383 ^a	25458 ± 4616 ^a	14917 ± 1500 ^b	19833 ± 3840 ^b	1207 ± 388 ^b	667 ± 133 ^b	640 ± 69 ^b
No fertilizer	12542 ± 2648 ^b	3333 ± 417 ^b	2500 ± 224 ^c	1542 ± 384 ^b	33 ± 4 ^a	40 ± 10 ^a	17 ± 7 ^a
P-Values	0.001	0.001	0.0001	0.001	0.004	0.0001	0.0001

Means with different superscript within columns indicate significant differences (Tukey's multiple range test at P<0.05).

Table 1: Phytoplankton and zooplankton abundance (individuals/L) observed in tanks fertilized with different type of fertilizers (Mean ± SE).

Treatments	Phytoplankton/L			Zooplankton/L		
	H'	H' _{max}	J	H'	H' _{max}	J
Chicken manure	1.13	1.3863	0.8151	1.0728	1.0986	0.9765
DAP fertilizer	1.1797	1.3863	0.8509	1.0526	1.0986	0.9581
No fertilizer	1.0489	1.3863	0.7566	1.0431	1.0986	0.9494

Table 2: Shannon-Wiener indices of phytoplankton and zooplankton diversity (individuals/L) in tanks applied with chicken manure DAP fertilizer and no manure application.

Parameters	Initial mean weight (g)	Individual final mean weight (g)	Weight gain (g)	Increase mean weight (g)	Specific growth rate (%)	Survival rate (%)
Stocking densities						
5fry/m ²	0.5 ± 0.01	24.1 ± 2.1 ^a	6.3 ± 0.4 ^a	90.7 ± 10.9	21.6 ± 1.7 ^a	97.8 ± 0.9 ^a
10fry/m ²	0.5 ± 0.01	18.3 ± 2.1 ^b	4.6 ± 0.4 ^b	80.6 ± 10.9	16.5 ± 1.7 ^b	87.2 ± 0.9 ^b
P-values	0.17	0.047	0.008	0.62	0.034	0.0001
Fertilizers						
Chicken manure	0.5 ± 0.01	25.6 ± 2.5 ^a	6.5 ± 0.5 ^a	90.8 ± 13.4	22.5 ± 2.1 ^a	92.1 ± 1.2
DAP Fertilizer	0.5 ± 0.01	20.5 ± 2.5 ^a	5.2 ± 0.5 ^a	84.3 ± 13.4	18.7 ± 2.1 ^a	93.9 ± 1.2
No fertilizer	0.5 ± 0.01	17.5 ± 2.5 ^b	4.5 ± 0.5 ^b	81.7 ± 13.4	15.8 ± 2.1 ^b	91.5 ± 1.2
P-values	1.0	0.047	0.034	0.886	0.042	0.32
Stocking density*Fertilizer						
5fry/m ² *chicken manure	0.5 ± 0.01	29.2 ± 3.6	7.6 ± 0.8	92.6 ± 19.0	25.6 ± 2.9	93.3 ± 1.7 ^a
5fry/m ² *DAP Fertilizer	0.5 ± 0.01	24.3 ± 3.6	5.9 ± 0.8	90.9 ± 19.0	20.6 ± 2.9	100.0 ± 1.7 ^b
5fry/m ² *no fertilizer	0.5 ± 0.01	18.8 ± 3.6	5.2 ± 0.8	88.5 ± 19.0	15.6 ± 2.9	100.0 ± 1.7 ^b
10fry/m ² *chicken manure	0.5 ± 0.01	21.9 ± 3.6	5.4 ± 0.8	89.0 ± 19.0	19.6 ± 2.9	90.8 ± 1.7 ^a
10fry/m ² *DAP Fertilizer	0.5 ± 0.01	16.7 ± 3.55	4.4 ± 0.8	77.7 ± 19.0	16.7 ± 2.9	87.9 ± 1.7 ^a
10fry/m ² *no fertilizer	0.5 ± 0.01	16.3 ± 3.6	3.9 ± 0.8	74.9 ± 19.0	13.1 ± 2.9	82.9 ± 1.7 ^c
P-values	1.0	0.73	0.87	0.95	0.94	0.0001

Means with different letters in the same columns are significantly different (Tukey's multiple range test at P<0.05)

Table 3: Growth parameters and survival observed in chicken manure, di-ammonium fertilizer and no fertilizer at different stocking densities (Mean ± SE).

Parameters	Temp (°C) at am	Temp (°C) at pm	DO (mgL ⁻¹) at am	DO (mgL ⁻¹) at pm	NO ₂ (ppm)	NH ₃ (ppm)	pH ranges
Stocking densities							
5fry/m ²	26.8 ± 0.1	28.6 ± 0.1	4.2 ± 0.1	6.2 ± 0.1	0.2 ± 0.07	0.5 ± 0.2	5.0-8.0
10fry/m ²	26.8 ± 0.1	28.5 ± 0.1	4.3 ± 0.1	6.3 ± 0.1	0.3 ± 0.07	0.5 ± 0.2	5.0-8.0
P-values	0.61	0.71	0.29	0.43	0.21	0.86	
Fertilizers							
Chicken manure	26.7 ± 0.1	28.6 ± 0.1	4.3 ± 0.8	6.1 ± 0.1 ^a	0.2 ± 0.09 ^{ab}	0.3 ± 0.2	6.0-8.0
DAP Fertilizer	26.8 ± 0.1	28.5 ± 0.1	4.3 ± 0.8	6.5 ± 0.1 ^b	0.4 ± 0.09 ^a	0.8 ± 0.2	5.0-7.0
No fertilizer	26.8 ± 0.1	28.6 ± 0.1	4.3 ± 0.8	6.0 ± 0.1 ^a	0.04 ± 0.09 ^b	0.3 ± 0.2	7.0-8.0
P-values	0.78	0.85	0.96	0.016	0.031	0.113	
Stocking density*Fertilizer							
5fry/m ² *chicken manure	26.7 ± 0.2	28.7 ± 0.2	4.2 ± 0.1	6.1 ± 0.2 ^a	0.1 ± 0.1 ^a	0.4 ± 0.3	6.0-8.0
10fry/m ² *chicken manure	26.8 ± 0.2	28.5 ± 0.2	4.3 ± 0.1	6.1 ± 0.2 ^a	0.3 ± 0.1 ^a	0.3 ± 0.3	6.0-8.0
5fry/m ² *DAP fertilizer	26.8 ± 0.2	28.5 ± 0.2	4.2 ± 0.1	6.3 ± 0.2 ^b	0.3 ± 0.1 ^b	0.9 ± 0.3	5.0-7.0
10fry/m ² *DAP fertilizer	26.8 ± 0.2	28.5 ± 0.2	4.3 ± 0.1	6.7 ± 0.2 ^b	0.5 ± 0.1 ^b	0.9 ± 0.3	5.0-7.0
5fry/m ² *no fertilizer	26.8 ± 0.2	28.7 ± 0.2	4.2 ± 0.1	6.0 ± 0.2 ^a	0.02 ± 0.1 ^c	0.3 ± 0.3	6.0-8.0
10fry/m ² *no fertilizer	26.9 ± 0.2	28.6 ± 0.2	4.3 ± 0.1	6.0 ± 0.2 ^a	0.07 ± 0.1 ^c	0.3 ± 0.3	7.0-8.0
P-values	0.95	0.88	0.92	0.03	0.018	0.99	

Means with different letters in the same columns are significantly different (Tukey's multiple range test at P<0.05)

Table 4: Water quality parameters observed in fertilizer type and different stocking densities (Mean ± SE).

reported rotifers to be the common and dominant *zooplankton* group in fertilized ponds. In addition, Mischke and Zimba [44] reported that copepod nauplii and cladocerans were significantly higher in ponds fertilized with inorganic fertilizer (Urea + Super phosphate) than in control ponds and organically fertilized ponds. Higher *zooplankton* diversities were recorded in chicken manure compared to other treatments. This might be due to *phytoplankton* particle size and cell abundance which influences *zooplankton* communities [45]. The larger size of *phytoplankton* cells were not consumed by smaller *zooplankton* hence high abundance of *phytoplankton* communities in DAP fertilized tanks than *zooplankton*. Similarly, Soderberg [46] reported that organic fertilizers require bacteria and other microbes for decomposition, and thus offer a wider diversity of fish ponds, particularly *zooplankton*. Also, chicken manure have been reported to provide a substrate for *zooplankton* production [47,48] which enhance high diversity.

Among water quality parameters, temperature and dissolved

oxygen were within the optimal range. In the present study, pH ranges were similar to those reported by Kumar et al. [22]. Concentration of NO₂ and NH₃ were high in the chicken manure and DAP fertilizer treatments as those reported by Boyd [49]. The reason might be due to temperature and pH dependence of ammonia and ammonium ion from the fertilizer. Ammonium fertilizer such as DAP increases total ammonia nitrogen concentrations in water while manures have low nitrogen content and tend to decompose slowly and incompletely [49].

African catfish fry had higher performance in fertilized tanks than unfertilized. The same results have been reported by Muendo et al. [50]. Individual mean weight and weight gain showed to be high at low stocking density (5fry/m²) compared to high stocking density (10fry/m²) across all fertilizers type (Table 3). This indicated that the performance of catfish fry was influenced by stocking density and fertilizers type. Similarly, Jamabo and Keremah [51] reported significantly higher growth rate at a stocking density of 5fry compared




Figure 1: Average weight of catfish fry reared in tanks fertilized with different types of fertilizer at different stocking densities for 8 weeks.

to 10fry and 15fry per 55m³ tank. Specific growth rate of African catfish (*Clarias gariepinus*) fry were observed to be improved in tanks fertilized with chicken manure and DAP fertilizers compared to unfertilized tanks under both stocking densities. Hossain et al. [52], Jamabo and Keremah [51] reported that growth of catfish fry was strongly influenced by stocking density. This might be due to high natural food organisms and good water quality found in fertilized tanks at low stocking densities compared to high stocking densities [53]. Therefore, these results indicated that the addition of fertilizers to the fish ponds/tanks increase the level of nitrogenous compounds and minerals which considered as a good source for *phytoplankton* growth. Lin et al. [54], Dang and Dalsgaard [55]; Saad et al. [56] reported that addition of manures and fertilizers to the fish ponds improve the level of *phytoplankton* that is responsible for growth fish. This was supported by Silva et al. [57] who reported that *phytoplankton* and *zooplankton* are rich source of protein often containing 40-60% protein on a dry matter basis and is sufficient to support excellent fish growth. Saad et al. [56] found that, nitrogenous and phosphorus compounds fertilizers improve the growth *phytoplankton* and *zooplankton* that in turn improve growth performance of the fish.

Survival rate can be influenced by stocking density and fertilizers type. Similarly, Haylor [58] found that survival rates in African catfish fry were directly related to stocking density. The study recorded high survival rate in DAP fertilized tanks and unfertilized tanks compared to chicken manure under low stocking density. While the lowest survival rate recorded in tanks with no fertilizer at high stocking density. This might be due to low dissolved oxygen recorded. Similar results with different fish species were reported by Sophin and Preston [59] that poorer survival rates of tilapia were observed in the ponds fertilized with fresh manure as compared to the inorganic fertilizers (urea and DAP) and unfertilized ponds. In addition, aggressive behavior of catfish in taking in food and cannibalism in control tanks might influence low survival rate. Similar results have been reported by Gamal [38] who found that high stocking density (5fisherlings/m²) using chemical fertilizer (urea + monophosphate) and chicken manure had the highest survival rate (96.48%) of catfish and indicated that predation behavior of catfish increased in the higher stocking densities. In addition, Baskerville-Bridges and Kling [60] reported greater risk of mortality at high stocking densities as a result of deteriorating water quality.

Water quality parameters in this study were within recommended

range for growth and survival of catfish fry. There were no significant differences ($P>0.05$) in temperature, dissolved oxygen during the morning and ammonia within stocking densities and fertilizer types. Temperature in all fertilizer types and stocking density was within optimal growth and survival range of catfish fry which is 26°C to 29°C [61,62]. Dissolved oxygen during the afternoon was higher in DAP fertilized tanks compared to chicken and control which was above recommended value of 6mg/l [63]. This might be due to high abundance of planktons [38]. The concentration of NO₂ was high at 10fry/m² under DAP fertilizer compared to other treatments. This might be due to ammonia utilization by *phytoplankton* [64] or oxidation of ammonia nitrite especially in high dissolved oxygen level conditions [65]. In addition pH ranges were optimal as recorded by Tucker 1991. However, were similarly higher under both stocking densities in chicken manure and DAP fertilized tanks compared to control. Boyd [66] reported that the application of ammonium and urea-based fertilizers can cause acidification of pond water because of nitrification, which produces two hydrogen ions from each ammonium ion. Similarly, Saad et al. [56] reported high pH values in treatments that received inorganic and organic fertilizers compared to group given feed only. Therefore, the results indicated that addition of fertilizers to the fish tank increased some of the physicochemical properties of the water.

Conclusion

In conclusion, the study indicated that, DAP fertilizer had highest plankton abundance and diversity compared to chicken manure and no fertilizer application. Hence it is thus recommended to use DAP fertilizer for production of live food in fish tanks. Higher growth performance was recorded at low stocking density (5fry/m²) across all fertilizer types. Survival rate of catfish fry did not depend on fertilizer type but it was depended on stocking density. Water quality parameters were no significant differences in chicken manure and di-ammonium phosphate at different stocking densities. Hence it is recommended to raise catfish fry in DAP or chicken manure fertilized tanks at low stocking density (5fry/m²) for better growth and survival in aquaculture practices.

Acknowledgements

We thank the Department of Aquaculture and Fisheries Science (LUANAR), and Department of Animal Science Production (SUA). We also thank the Innovative Agricultural Research Initiative (iAGRI) under Feed the Future Program and

RUFORUM for providing financial support to enable first author pursue MSc studies in Malawi.

References

- Govoni JJ, Boehlert GW, Watanabe Y (1986) The physiology of digestion in fish larvae. *Environmental Biology of Fish* 16: 59-77.
- Olurin KB, Iwuchukwu PO, Oladapo O (2012) Larval rearing of African catfish, *Clarias gariepinus* fed decapsulated *Artemia*, wild copepods or commercial starter diet. *African Journal of Food Science and Technology* 3: 182-185.
- Adewumi AA (2015) Growth performance and survival of *Clarias gariepinus* hatchlings fed different starter diets. *European Journal of Experimental Biology* 3: 1-5.
- Cahu CL, Zambonino-Infante JL (2001) Substitution of live food by formulated diets in marine fish larvae. *Aquaculture* 200: 161-180.
- Agadjihouede H, Chikou A, Bonou CA, Laleye PA (2012) Survival and Growth of *Clarias gariepinus* and *Heterobranchus longifilis* Larvae fed with fresh water zooplankton. *Journal of Agricultural Science and Technology* 2: 192-197.
- Schroeder GL (1980) Fish farming in manure loaded ponds. In integrated agriculture-aquaculture farming systems. In International Centre for Living Aquatic Resources Management, Manila, Philippines.
- Olurin KB, Oluwo AB (2010) Growth and survival of African catfish (*Clarias gariepinus*) larvae fed decapsulated *Artemia*, live daphnia, or commercial starter diet. *Journal of Aquaculture* 62: 50-55.
- De Graaf G, Janssen H (1996) Artificial reproduction and pond rearing of the African catfish *Clarias gariepinus* in sub-Saharan Africa. *Fisheries Technical Paper*. No. 362. Rome.
- Wurts WA (2004) Organic fertilization in culture ponds. *Aquaculture* 35: 64-66.
- Chenyambuga SW, Madalla N, Mnembuka B (2011) Management and value chain of Nile tilapia cultured in ponds of small-scale farmers in Morogoro region, Tanzania. IFFET paper, Sokoine University of Agriculture, Morogoro, Tanzania.
- Murnyak D (2010) Basics of raising tilapia and implementing aquaculture projects. In: *Fish Farming, ECHO Technical Note*, North Fort Myers.
- Schram E, Van der Heul JW, Kamstra A, Verdegem MCJ (2006) Stocking density dependent growth of dover (*Solea solea*). *Aquaculture* 252: 239-247.
- Nwipie GN, Erondu ES, Zabbey N (2015) Influence of stocking density on growth and survival of post fry of the African Mud Catfish, *Clarias gariepinus*. *Fish and Aquaculture* 6: 2-4.
- Brazil BL, Wolters WR (2002) Hatching success and fingerling growth of channel catfish cultured in ozonated hatchery water. *North American Journal of Aquaculture* 64: 144-149.
- Pangni K, Atse BC, Kouassi NJ (2008) Effect of stocking density on growth and survival of the African catfish *Chrysichthys nigrodigitatus*, *Claroteidae* (Lacépède 1803) larvae in circular tanks. *Livestock Research for Rural Development* 20: 1-9.
- Hengsawut K, Ward FJ, Jaruratjamorn P (1997) The effect of stocking density on yield, growth and mortality of African catfish (*Clarias gariepinus*, Burchell 1822) cultured in cages. *Aquaculture* 152: 67-76.
- Toko I, Fioebe ED, Koukpodé B, Kestemont P (2007) Rearing of African catfish (*Clarias gariepinus*) and vundu catfish (*Heterobranchus longifilis*) in traditional fish ponds (whedos): Effect of stocking density on growth, production and body composition. *Aquaculture* 262: 65-72.
- Hecht T, Uys W (1997) Effect of density on the feeding and aggressive behaviour in juvenile African catfish, *Clarias gariepinus*. *South African Journal of Science* 93: 537-541.
- Obirikorang KA, Madkour HA, Adjei-Boateng D (2014) A study of intra-cohort cannibalism in juveniles of the African Catfish, (*Clarias gariepinus*) under Controlled Conditions. *International Journal of Science and Technology* 3: 23-26.
- Diana JS (2012) Some principles of pond fertilization for Nile tilapia using organic and inorganic inputs. USA.
- Akand MA (1986) Effect of organic fertilizers on water quality and fish production in carp ponds. *Bangladesh Journal of Fisheries Research* 9: 47-53.
- Kumar SD, Kumar SP, Kumar VU, Anbuganapathi G (2014) Efficacy of bio-fertilizer enriched flower waste vermin-compost on production and growth of primary producers and freshwater aquarium fishes *Global Veterinaria* 13: 215-220.
- Lin CK, Teichert-Coddington DR, Green BW, Veverica KL (1997) Fertilization regimes. In: *Dynamics of Pond Aquaculture*, CRC Press, Boca Raton.
- Amisah S, Adjei-Boateng D, Afianu DD (2008) Effects of bamboo substrate and supplementary feed on growth and production of the African catfish, (*Clarias gariepinus*). *Journal for Applied Science and Environmental Management* 12: 25-28.
- Thingstad TF, Krom MD, Mantoura RFC, Flaten GAF, Groom S, et al. (2005) Nature of P-limitation in the ultra-oligotrophic Eastern Mediterranean. *Science* 309: 1068-1071.
- Association of Official Analytical Chemists (AOAC) (2002) *Official Methods of Analysis*. (15th edn). JRC-European Commission, Washington, DC, USA.
- Law CS, Abraham ER, Woodward EMS, Liddicoat MI, Fileman TW, et al (2005) The fate of phosphate in an in situ Lagrangian addition experiment in the Eastern Mediterranean. *Deep-Sea Research II* 52: 2911-2927.
- United Nations Educational Scientific and Cultural Organization (UNESCO) (2006) *Manual on aquatic cyanobacteria*. A photo guide and a synopsis of their toxicology, Institute of Ecology/Limnology, Lund University, Sweden.
- Greenberg AE, Clesceri LS, Eaton AD (1992) *Standard methods for the examination of water and wastewater*, (18th edn). American Public Health Association. Port City Press, Baltimore, Maryland.
- Wetzel RG, Likens GE (1991) *Limnological Analyses*, (2nd edn). Springer-Verlag, New York, USA.
- Krebs CJ (1989) *Ecological methods for field and laboratory investigations*. Harper Collins Publishers, New York.
- Sundar S, Raina HS, Mohon M, Singh B (1995) Ecology and fisheries potentials of the Guala River with special reference to proposed impoundment (Jamrani dam) on the system. *Journal of Inland Fisheries Society of India* 27: 33-45.
- Bellinger ED (1992) *A key to common algae*. The Institution of Water and Environmental Management, London, UK.
- Kang'ombe J, Brown JA, Halfyard LC (2006) Effect of using different types of organic animal manure on plankton abundance, and on growth and survival of *Tilapia rendalli* (Boulenger) in ponds. *Aquaculture Research* 37: 1360-1371.
- SPSS (1999) *SPSS*. SPSS, Chicago, IL, USA.
- Kumar A, Sharma LL, Aery NC (2008) Physicochemical characteristics and diatom as indicators of Trophic status of Kishore Sagar, Rajasthan In: *Proceedings of Taal 2007: The 12th World Lake Conference*, India.
- Lubzen E, Zmora O, Barr Y (2001) Biotechnology and aquaculture of rotifers. *Rotifera IX Developments in Hydrology* 153: 337-353.
- Gamal OE, Ibrahim NA, Abou ZMY (2008) Influence of fertilizers' types and stocking density on water quality and growth performance of Nile tilapia-African catfish in polyculture system. 8th International Symposium on Tilapia in Aquaculture, Sharkia, Egypt.
- Padmavathi P, Veeraiah K (2009) Studies on the influence of *Microcystis aeruginosa* on the ecology and fish production of carp culture ponds. *African Journal of Biotechnology* 8: 1911-1918.
- Boyd CE, Massaut L (1999) Risks associated with the use of chemicals in pond aquaculture. *Aquacultural Engineering* 20: 113-132.
- Guangjun LV (2013) Structure and diversity of zooplankton communities in four reservoirs with varying nutrient compositions in the Yangtze River Basin, China. Southwest University, Rongchong, China.
- Mischke CC, Wise DJ, Byars TS (2009) Evaluation of Zooplankton in Hatchery Diets for Channel Catfish Fry. *North American Journal of Aquaculture* 71: 312-314.
- Ghosh AL, Rao H, Banerjee SC (1974) Studies on hydrobiological conditions of sewage fed pond with a note of their role in fish culture. *Journal of Inland Fisheries Society of India* 6: 51-61.
- Mischke CC, Zimba PV (2004) Plankton community responses in earthen channel catfish nursery ponds under various fertilization regimes. *Aquaculture* 233: 219-235.
- Bell T (2002) The ecological consequences of unpalatable prey: Phytoplankton response to nutrient and predator additions. *Oikos* 99: 59-68.

46. Soderberg RW (2012) Organic and inorganic fertilization. In: Aquaculture pond fertilization: impacts of Nutrient Input on Production, Wiley-Blackwell, USA.
47. Food and Agriculture Organization (FAO) (2003) Seed production. An overview, FAO Fisheries and Aquaculture Department, Rome, Italy.
48. Hossain MY, Begum M, Ahmed ZF, Hoque MA, Karim MA, et al. (2006) A study on the effects of Iso-Phosphorus Fertilizers on plankton production in fish ponds. South Pacific Studies 26: 101-110.
49. Boyd CE (1997) Practical aspects of chemistry in pond aquaculture. Chapman & Hall, New York, USA.
50. Muendo PN, Milstein AA, Dam N, Gamal EI, Stoorvogel JJ, et al. (2006) Exploring the trophic structure in organically fertilized and feed driven tilapia culture environments using multivariate analyses. Aquaculture Research 37: 151-163.
51. Jamabo NA, Keremah RI (2009) Effects of stocking density on growth and survival of fingerlings *Clarias gariepinus* (Burchell, 1822). Journal of Fisheries International 4: 55-57.
52. Hossain MAR, Beveridge MCM (1998) The effects of density, light and shelter on the growth and survival of African catfish (*Clarias gariepinus* Burchell, 1822) fingerlings. Aquaculture 160: 251-258.
53. Bahnasawy MH (2009) Effect of dietary protein levels on growth performance and body composition of monosex Nile tilapia, *Oreochromis niloticus* L. reared in fertilized tanks. Pakistan Journal of Nutrition 8: 674-678.
54. Lin CK, Yang Y, Shivappa RB, Kabir CMA (1998) Optimization of nitrogen fertilization rate in freshwater tilapia production ponds. In: PD/A CRSP Sixteenth Annual Technical Report, Oregon State University, USA.
55. Dang ST, Dalsgaard A (2012) *Escherichia coli* contamination of fish raised in integrated pig-fish aquaculture systems in Vietnam. Journal of Food Protection 75: 1317-1319.
56. Saad TT, El-Hawary WN, Abel-lateef MA (2014) Growth and economic performance of *Oreochromis niloticus* in earthen ponds. Journal of Middle East Applied Science and Technology (JMEAST) 11: 463-475.
57. Silva SS, Anderson TA (1995) Fish nutrition in aquaculture, Chapman and Hall, New Delhi, India.
58. Haylor GS (1992) Controlled hatchery production of *Clarias gariepinus* (Burchell, 1822): growth and survival of fry at higher stocking density. Aquaculture of Fisheries Management 22: 405-422.
59. Sophin P, Preston TR (2001) Effect of processing pig manure in a biodigester as fertilizer input for ponds growing fish in polyculture. Livestock Research for Rural Development 13: 1-15.
60. Baskerville-Bridges B, Kling LJ (2000) Larval culture of Atlantic cod (*Gadus morhua*) at high stocking densities. Aquaculture 181: 61-69.
61. Tucker CS (1991) Water quantity and quality requirements for channel catfish hatcheries. Southern Regional Aquaculture Center, Mississippi State University, USA.
62. Ahmed GU, Islam MF, Khan MNA, Haque MM, Kibria MG (1997) Culture feasibility of African catfish (*Clarias gariepinus* Lin.) fry in glass tank and synthetic hapa system using supplemental diets. Bangladesh journal of Fish Research 2: 59-66.
63. Food and Agriculture Organization (FAO) (1996) Artificial reproduction and pond rearing of the African Catfish *Clarias gariepinus* in Sub-Saharan Africa-A Handbook, FAO Fisheries and Aquaculture Department, Rome, Italy.
64. Boyd CE, Tucker CS (1998) Pond aquaculture water quality management. Kluwer Academic Publishers. Boston.
65. Boyd CE (2000) Water quality. Kluwer Academic publishers, Massachusetts, USA.
66. Boyd CE (1990) Water quality in ponds for aqua culture. Alabama Agricultural Experiment Station, Auburn University, AL, USA.

Citation: Mosha SS, Kang'ombe J, Jere W, Madalla N (2016) Effect of Organic and Inorganic Fertilizers on Natural Food Composition and Performance of African Catfish (*Clarias gariepinus*) Fry Produced Under Artificial Propagation. J Aquac Res Development 7: 441. doi: [10.4172/2155-9546.1000441](https://doi.org/10.4172/2155-9546.1000441)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>