

Effect of Pesticide on Nitrate Reductase Activity in *Trigonella Foenum* i.e., Fenugreek

Pankaj SM* and Rohit P Gore

Department of Biotechnology, Yashwantrao Chavan Institute of Science, Satara. Maharashtra, India

*Corresponding author: Department of Biotechnology, Yashwantrao Chavan Institute of Science, Satara. Maharashtra, India, Tel: +919552991234; E-mail: mundada.pankaj77@gmail.com

Received date: Sep 05, 2015; Accepted date: Nov 23, 2015; Published date: Nov 26, 2015

Copyright: © 2015 Pankaj SM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.


Abstract

Nitrate reductase (NR) is the key enzyme of nitrate assimilation subjected to inhibition by excessive use of pesticides. Now a day's pesticides are used commonly at a large scale on field. In the present study, the effect of pesticides on nitrate reductase activity has been studied by in vivo assay in *Trigonella foenum-graecum*. The four pesticides Confidor, Omite, Karathane, Chlorpyrifos+Cypermethrin were used for study at various concentrations such as 25, 50, 100 and 200 µl/100 ml. The application of pesticides showed decrease in activity of nitrate reductase enzyme. Out of these pesticides confidor affected the nitrate reductase activity maximum. The nitrate reductase activity in control plant was found to be $0.460U \pm 0.005$ which was reduced up to $0.192 \pm 0.008U$ in presence of confidor. Even in the tissue specific assay it was seen that the leaves and stem of the plant, the NR activity was affected maximum than other parts. The use of pesticides lowers the NR activity of the *Trigonella* plants thereby affecting its nitrogen metabolism.

Keywords: Nitrate reductase; Confidor; Omite; Karathane; Chlorpyrifos+cypermethrin; In vivo NR-Assay

Introduction

Nitrate reductase (NR; EC 1.6.6.1.3) is one of the most important enzymes in the assimilation of exogenous nitrate—the predominant form of nitrogen available to green plants growing in soil. It is monomer composed of 100 KD polypeptide and one each of FAD, Heme ion and molybdenum



Nitrate reduction, the most important stage of nitrogen turnover in nature, has several functions:

1. Utilization of NO_3^- as a source of nitrogen (nitrate assimilation);
2. Production of metabolic energy during NO_3^- utilization as terminal acceptor of electrons (nitrate respiration), and
3. Dissipation of excess of reducing energy to maintain oxidation–reduction balance (nitrate dissimilation).

It is an enzyme that is being used to help clean up the environment, by conversion of nitrate to nitrite and has great potential to be part of the solution to the global problem of excess nitrate and related nitrogen nutrients in water sources [1]. Nitrate is currently one of the most hazardous pollutants [2-4]. Activity of this enzyme in plants gives a good estimate of the nitrogen status of the plant and is very often correlated with growth and yield. Nitrate reduction processes are perhaps most significant in maintaining water quality. NR enzyme is responsible not only for nitrate reduction, but also for nitrate uptake in *Chlamydomonas*, *Chlorella sorokiniana* [5,6]. Assimilatory Nitrate reduction, performed by a variety of bacterial and eukaryotic organisms [7,8] includes the reduction of nitrate and nitrite compounds, often mediated by nitrate or nitrite reductase enzymes, for

synthesis of amino acids. The ability of nitrate to induce nitrate reductase for nitrate metabolism in higher plants has been widely accepted [9]. The nitrate reductase activity is inhibited by heavy metals like Cr^{3+} and Cu^{2+} irreversibly [10]. It has also been reported that insecticides (kelthane and fenvalerate) caused some inhibition to enzyme activities [6]. In some experiments 15 different pesticides were evaluated to study their effect on growth and nitrogen assimilation of the *Azolla Mexicana* [11]. It was seen that phenolic herbicides were detrimental causing 75% reduction in nitrogen fixation and nitrate reduction. Choramber (3 amino, 2-5 dichlorobenzoic acid) and benomyl (methyl 1-butyl carbamoyl 2-bezimidazolyl carbamate) caused 84-99% reduction in nitrogen fixation [12]. The toxic insecticide Trichlorfon induced inhibition of nitrogen uptake in Cyanobacteria [13]. In the present study the effect of pesticide on nitrate reductase enzyme activity is studied in vivo in Fenugreek i.e., *Trigonella foenum*. The plant material was selected since it is easy to grow and have short life cycle.

Materials and Methods

Plant material

The present study involves the use of *Trigonella Foenum-graecum* i.e., Fenugreek. The seeds of the fenugreek were purchased from the market and grown in different pots in the laboratory. The plantlets were identified as the *Trigonella foenum-graecum* with the help of Department of Botany, Yashwantrao Chavan Institute of Science, Satara.

Pesticides

From a common survey of around 100 farmers using pesticides around Satara City, Confidor, Omite, Karathane, Chlorpyrifos

+Cypermethrin were selected as the most commonly used pesticides on fields for the present study.

In vivo Nr Assay

Nitrate reductase (NR) activity can be directly assayed using cut plant tissues. In a solution containing nitrate, phosphate buffer and propanol, nitrate can readily enter plant material. In this case, the plant cell supplies NR and the reductant NADH. The propanol in the assay medium increases cell permeability and makes the leaf segments essentially anaerobic. Under anaerobic conditions, the reduction of nitrite to ammonia is inhibited, and the conversion of nitrate to nitrite is enhanced. Because of the sensitivity of the reaction to molecular oxygen all operations should be carried out in dim room light to reduce photosynthetic activity. With this approach the NR activity associated with all four inducing treatments can be assayed by one group [14].





Weigh out four 0.5 g leaf samples for each of the inducing treatments, cut the tissue into 1.0-1.5 cm squares segments. Prepare duplicate screw cap test tubes for each inducing condition. You will need one set for time=0 (t₀) and another set for time=30 min. (t₃₀).

Place the leaf samples in the appropriate tubes. Add 10 ml of In vivo assay solution to each tube and cap the tube. (100 mM phosphate buffer, pH 7.5, 30 mM KNO₃, 5% (v/v) propanol) Place the two tubes in a boiling water bath for 5 min. and then allow to cool to room temperature. Place all tubes, t₀ and t₃₀, in a shaking water bath at 30°C and incubate for 30 min. At the end of the 30-min. incubation period, place the t₃₀ tubes in a boiling water bath for 5 min. and then cool to room temperature. To detect nitrite in the assay tubes, add 1 ml 1% sulfanilamide in 3 N HCl, and 1ml of 0.02% N-(1-naphthyl)-ethylenediaminehydrochloride. Mix thoroughly and place tubes in the dark at room temperature for 15 min. Determine OD of each standard tube at 540 nm and calculate the enzyme activity.

(Note: All the experiments were repeated three times and the data presented is the mean of the readings.)

Results

The results are shown in graphical representation (Graphs 1-4).



Discussion

In the present investigation from the result clearly indicates the inhibition of nitrate reductase activity after using the pesticides Confidor, Omite, Karathane, Chlorpyrifos+Cypermethrin, on *Trigonella Foenum-graecum*. Nitrate reductase activity of control pot of whole plant was found to be 0.460 units however it is severely

affected after using confidore and is reduced to 0.192 units. After studying different parts of the plant it is observed that leaves are the most affected by the use of pesticides than root and stem.

Different pesticides used in the current study showed the inhibitory effect on the nitrate reductase in the following order Confidor>Chlorpyrifos 50%+Cypermethrin 5% >Karathane>Omite.

All the pesticide under investigation showed the inhibitory effect on the nitrate reductase, thereby affecting the ability of the plant to utilize the nitrate.

Smiller studies on insecticides such as kelthane and fenvalrate caused inhibition of nitrate reductase in fungi from root nodules and faba bean (*Vicia faba* L.).

Chromium and copper metals irreversibly inhibited nitrogen reductase in *Aspergillus niger*.

Several pesticides have been proved to affect growth and nitrogen assimilation of *Azolla anbaena* symbiosis (especially *Azolla mexicana*). Toxic insecticide trichlorfon inhibits the nitrogen and Ammonium uptake in cyanobacteria. Cadmium, acetochlor and bensulfuron methyl have toxic effect on nitrogen metabolism and plant growth in rice, seedlings resulting in marked decrease in the fresh weight and the activities of nitrate reductase (NR).

Number of researchers proved in different plants use of pesticides leading to inhibition of nitrate reductase. In the present investigation the commonly used pesticides showed similar inhibitory effect on *Trigonella Foenum-graecum* and compelling us to use nitrate reductase friendly methodology in place of above pesticides.

References

1. Campbell ER, Campbell WH (1998) Determination of nitrate in aqueous matrices using nitrate reductase. In: Current protocols in field analytical chemistry, supplement 1, Chapter 5 "Water Quality Parameters-Anions", John Wiley and Sons, Inc.
2. Awasthi M, Rai LC (2005) Toxicity of Nickel, Zinc and Cadmium to nitrate uptake in free and immobilized cells of *Scenedesmus quadricauda*. *Ecotox Environ Safe* 61: 268-272.
3. Bier J (2002) Nitrate in groundwater: Sources, Impacts and Solutions. 6th Symposium on Groundwater contaminants, Fresno, CA.
4. Johnson CJ, Kross BC (1990) Continuing importance of nitrate contamination of groundwater and wells in rural areas. *Am J Int Med* 18: 449-456.
5. Knobloch O, Tischner R (1989) Characterization of nitrate reductase deficient mutants of *Chlorella sorokiniana*. *Plant Physiol* 89: 786-791.
6. Nichols GL, Shehata SAM, Syrett PJ (1978) Nitrate reductase deficient mutants of *Chlamydomonas reinhardtii*: Biochemical characteristics. *J Gen Microbiol* 108: 79-88.
7. Solomonson LP, Barber MJ (1990) Assimilatory nitrate reductase: functional properties and regulation. *Plant Mol Biol* 41: 225-253.
8. Zvyagil'skaya RA, Vartapetyan BB, Vov NPL (1996) Nitrate dissimilation in eukaryotes. *Appl Biochem Microbiol* 32: 165-169.
9. Beevers L, Schrader LE, Fleasher D, Hageman RH (1965) The role of light and nitrate and nitrite in the induction of nitrate reductase in radish cotyledons and maize seedlings. *Plant Physiology* 40: 691-698.
10. Abigail M Aiken, Brent M Peyton, William A Apel, James N Petersen (2003) Heavy metal-induced inhibition of *Aspergillus niger* nitrate reductase: applications for rapid contaminant detection in aqueous samples. *Analytica Chimica Acta* 480: 131-142.
11. Holst RW, Yopp JH, Kapusta G (1982) Effect of several pesticides on the growth and nitrogen assimilation of *Azolla-Anabaena* symbiosis. *Weed sci* 30: 54-58.
12. Omar SA, Abd-Alla MH (2000) Physiological aspects of fungi isolated from root nodules of faba bean (*Vicia faba* L.). *Microbiol Res Mar*: 154: 339-47.
13. Marco E, Orus MI (1992) Trichlorfon-Induced Inhibition of Nitrate and Ammonium Uptake in Cyanobacteria. *Journal of Experimental Botany* 44: 501-508.
14. Jaworski EG (1971) Nitrate reductase assay in intact plant tissues. *Biochem Biophys Res Commun* 43: 1274-1279.