

Effects of Fish Meal Substitution with Poultry By-product Meal on Growth Performance, Nutrients Utilization and Blood Contents of Juvenile Nile Tilapia (*Oreochromis niloticus*)

Yones AMM* and Metwalli AA

National Institute of Oceanography and Fisheries, Shakshouk Fish Research Station, El-Fayoum, Egypt

Abstract

Four isonitrogenous and isocaloric diets ($30.22 \pm 0.02\%$ CP and 19.007 ± 0.015 MJ kg^{-1} diet) were formulated to represent four dietary treatments. The first treatment (control) without poultry by-product (PM0), while the second, third and fourth diets formulated with substitution of fish meal by poultry by-product meal as 50, 75 and 100%, respectively. Each diet was fed to 100 juvenile tilapia (1.5 ± 0.05 g), in triplicate cement ponds of 2 m³. Fish were fed diets at a rate of 3% of its biomass daily divided into two equal portions. The highest ($P < 0.05$) growth performance parameters (final weight, weight gain, daily gain and specific growth rate) and best nutrient utilization (feed conversion ratio, protein efficiency ratio and net protein utilization) were recorded with PBM0 and PBM100% groups. The applied treatments showed insignificant effects on nutrients digestibility coefficient among dietary groups for Dry matter, Energy, CP, Fat and nitrogen free extract. No significant effects on dry matter, crude protein, fat and ash contents were recorded between dietary treatments. Juvenile tilapia fed the experimental diets showed insignificant different in blood contents ($P < 0.05$) between groups. The present study recommended substitution of 100% fish meal with poultry by-product meal in juvenile Nile tilapia diets.

Keywords: *Oreochromis niloticus*; Poultry by-product meal; Digestibility coefficient; Growth performance; Blood contents

Introduction

The global production of tilapia drastically increased from 124 thousand metric tons (Mt) in 1997 to 2.5 million Mt in 2010 [1]. This trend suggests that there will be even greater increases in the future. Among the cichlid species, it is the Nile tilapia (*Oreochromis niloticus*) that has dominated in different culture system. The tilapia market has expanded from a subsistence level to meet the protein needs of the middle class because of the year-round supply, delicious flavour and reasonable price of that fish [2]. Global tilapia production was recorded 3,500,000 metric tons in 2011, 3,800,000 metric ton in 2012, 4,850,000 metric ton in 2014 and by 2015, world production tilapia is forecast to reach 4.6-5.0 million metric ton [3,4].

Traditionally, fish meal has provided a major part of protein sources in formulated feeds because of its suitable protein quality. Science the recent scarcity and uncertain consistency of supply encourage its replacement by alternative protein sources that are of high quality, but less expensive has been investigated in many fish species. The limitations on the world's food supply provide additional motivation [5,6]. Therefore, numerous studies have undertaken to examine the effects of replacing fish meal by another source of protein such as animal by-product or plant-based protein in diets that can be fed to tilapia [7,8].

Animal by-products such as meat meal, bone meal and poultry by-product meal have considerable potential as feed ingredients in fish production system [9-12] and comparatively less expensive than fish meal [13]. These animal protein ingredients are good sources of amino acids with high protein content, total digestible dry matter and energy similar to fish meal [9]. Therefore, poultry by-product meal is considered a probable replacement for fish meal [14-21]. Many studies have also shown that animal protein ingredients can be useful for fish feed formulation and they are comparatively much less expensive than fish meal [13,22,23].

Some studies have shown that poultry by-product meal cannot replace more than 50% of fish meal in fish diets [24], but other studies have shown that with the recent improvement of the quality of poultry by-product meal it could replace 75% or 100% of fish meal without significant decrease in fish growth [25-28].

The present study aims to evaluate the effects of use poultry by-product meal as a alternative protein source to replace fish meal on growth performance, nutrient utilization, digestibility coefficient and some blood constituents of Nile tilapia (*Oreochromis niloticus*) reared in cement tanks.

Materials and Methods

Fish culture and experimental diets

The present study was conducted using the research facilities of the experimental station at Shakshouk, Fayoum Governorate, National Institute of Oceanography and Fisheries (NIOF). The system contained two water pumps and upstream sandy filter units at a point between the water source and tanks. Each pump was drowning the water from the lake Qaroun to collection cement pond and forced it through storage units and then to the rearing tanks in open system. Physicochemical characteristics of water tanks were examined every week, (Table 1) according to APHA [29].

*Corresponding author: Yones AMM, National Institute of Oceanography and Fisheries, Shakshouk Fish Research Station, El -Fayoum, Egypt, Tel: (+202) 7921342; E-mail: yones_552000@yahoo.com

Received September 10, 2015; Accepted October 19, 2015; Published January 15, 2016

Citation: Yones AMM, Metwalli AA (2015) Effects of Fish Meal Substitution with Poultry By-product Meal on Growth Performance, Nutrients Utilization and Blood Contents of Juvenile Nile Tilapia (*Oreochromis niloticus*). J Aquac Res Development 6: 389. doi:10.4172/2155-9546.1000389

Copyright: © 2015 Yones AMM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Diets				
Parameters	PBM0	PBM50	PBM75	PBM100
Temperature °C	27.0 ± 0.12	27.1 ± 0.12	27.4 ± 0.1	27.1 ± 0.15
pH	7.6 ± 0.1	7.8 ± 0.12	7.8 ± 0.11	7.6 ± 0.1
Dissolved oxygen (mg l ⁻¹)	6.3 ± 0.11	6.2 ± 0.12	6.2 ± 0.11	6.1 ± 0.12
Salinity (g l ⁻¹)	2.22 ± 0.1	2.23 ± 0.1	2.22 ± 0.1	2.22 ± 0.1
Unionized ammonia (mg l ⁻¹)	0.03 ± 0.01	0.028 ± 0.01	0.026 ± 0.002	0.028 ± 0.001

Table 1: Averages of water physicochemical characteristics parameters during experimental period.

Ingredientes	DM	CP	EE	NFE	CF	ASH
Fish meal	916	700	128	-	-	172
Poultry by-product meal	921	560	134	128	24	154
Soybean meal	915	480	44	368	42	66
Gluten meal	906	350	48	532	22	48
Wheat bran	904	144	34	666	92	68
Yellow maize	896	90	24	760	72	54

DM: Dry Matter, CP: Crude Protein, EE: Ether Extract, NFE: Nitrogen Free Extract, CF: Crude Fiber.

Table 2: Proximate composition of feed ingredients (g kg⁻¹ d.m) n=3.

The fry of Nile tilapia (*Oreochromis niloticus*) used in the present study were obtained after brood stock hatching in the station. The fry were acclimatized for two weeks in rearing tanks and fed on prepared powder diet contain 30% crude protein, formulated from the same ingredients use in the growth trial. Juvenile tilapia with an initial average weight of 1.5 ± 0.05 g were randomly distributed and stocked at 100 juvenile per tank in 12 cement tanks, each with a water volume of (2 m³) and the treatments were performed in triplicates. The diets were given at 3% of live body weight (BW) and offered in two equal portions at 10.00 a.m and 16.00 p.m. The experiment lasted 120 days after start.

Four isonitrogenous diets were formulated to contain an average of 30.11 ± 0.07% crude protein for meeting the recommended nutritional requirements of tilapia [30]. The ingredients were obtained from Zoocontrol fish Co at 6 October city, Egypt. Ingredients, diets formulation and chemical composition analysis are presented in Tables 2 and 3. The first diet was formulated without poultry by-product meal and considered as a control diet (PM0), Diets 2 (PM50), 3 (PM75) and 4 (PM100) were formulated to be comprised with partial and total inclusion levels of 50, 75 and 100% poultry by-product meal, respectively. All diets were processed into dry sinking pellet form, using California pelleting machine with 1 mm diameter.

Apparent digestibility coefficient

The experimental test diets with addition of 0.5% chromic oxide (Cr₂O₃) were fed to fishes after the growth trial lasted for a period of two week in order to study the apparent digestibility coefficient (ADC %) of nutrients. Any uneaten and fecal residues were siphoned out from the tank bottom after two hours of first feeding (10.00 a.m.) and discarded. Fish fecal samples were collected every afternoon before the second feeding, new fecal materials were carefully siphoned and collected using the filtration system developed by Choubert et al. [31]. After freeze-drying of 20 g fecal samples in each replicate tank, the feces were analyzed. Dry matter was calculated by gravimetric analysis at 105°C for 24 hrs. Chromic oxide levels were determined spectrometry (Spectra, AA220FSNarin) based on the method described by Bolin et al. [32].

The apparent digestibility coefficients ADC for test diets were calculated according to the equation described by Cho [33].

$$ADC(n) = 100 - \{100 (\%Cr_2O_3d) / \%Cr_2O_3f \times (\%Nf / \%Nd)\}$$

Where ADC (n)=apparent digestibility coefficients of a nutrient in the test diets; Cr₂O₃d=% chromic oxide of the diet; Cr₂O₃f=% chromic oxide of the feces; Nd=nutrient in the test diet; Nf=nutrients in feces.

Chemical analysis

The chemical composition of the experimental diets, feces and whole fish samples was performed via proximate composition analysis according to standard methods [34]. Briefly dry matter was determined gravimetrically in an oven dried samples at 105°C for 24h, protein (N×6.25) content was determined using Kjeldhal method and crude fat by chloroform-methanol extraction (2:1) using Soxhlet system. Ash was determined by incinerating samples in a Germany muffle furnace at 550°C for 18h. Nitrogen free extract (NFE) was calculated by the difference.

Diets samples were hydrolyzed in 6 N (HCL) at 106°C over 24 h in nitrogen-flushed glass vials before amino acid analysis. Total amino acids were analysed by high pressure liquid chromatography (HPLC) in a Pico-Tag amino acid analysis system (Water, Bedford, MA, (USA), using norleucine as internal standard and according to the procedure described by Cohen et al. [35].

Gross energy (MJ Kg⁻¹ diet) was calculated according to Schulz et al. [36] using the following calorific values: 23.9, 39.8 and 17.6 MJ g⁻¹ diet for protein, ether extract and nitrogen free extract, respectively. The metabolizable energy contents of the experimental diets were calculated as 18.9, 35.7 and 14.7 MJ g⁻¹ diet for protein, lipid and nitrogen free extract, respectively according to Jobling [37].

Diets				
Ingredients	PBM0	PBM50	PBM75	PBM100
Fish meal	200	100	50	-
Poultry by-product meal	-	150	200	260
Soybean meal	150	130	150	150
Corn gluten meal	110	100	100	100
Wheat brain	200	200	200	2100
Yellow corn maize	260	240	220	200
Fish oil	30	30	30	30
Sunflower oil	30	30	30	30
Vitamin and mineral mix ¹	15	15	15	15
Chromic oxide	5	5	5	5
Proximate analysis (n=3)				
Dry matter	928	922	924	926
Crude protein	301.9	301	301.8	300
Ether extract	110.4	115.9	116.6	118.8
Nitrogen free extract	453.9	451.3	449.2	447.8
Crude fiber	45.8	46.8	50.4	49.4
Crude ash	88	85	82	84
Gross energy (MJ kg ⁻¹ diet) ²	19.58	19.74	19.75	19.77
ME (MJ kg ⁻¹ diet) ³	16.31	16.44	16.46	16.49

¹Vitamin-mineral premix, mg Kg⁻¹ dry diets: vitamin A (as acetate), 7500 IU kg⁻¹ dry diet, Vitamin D3 (as cholecalciferol); 6000IU kg⁻¹ dry diet, vitamin E (as DL-L-tocopheryl-acetate); 150 IU kg⁻¹ dry diet, vitamin K (as menadione Na-bisulphate); 0.06 ascorbic acid (as ascorbyl polyphosphate), 150 D-biotin, 42 choline (as chloride) 3000; folic acid, 3 niacin (as nicotinic acid), 30 pantothenic acid, 60 pyridoxine, 15; ribflavine, 0.06; manganese sulphate, 0.18; potassium iodide, 0.02 zinc sulphate.

²Schulz et al. [36].

³Jobling [37].

Table 3: Formulation and approximate composition of experimental diets (g kg⁻¹).

Growth performance and feed utilization

Standard formulae were used to assess growth-feed utilization and other relevant parameters during the growth trial and these included, initial average weight, final average weight, total feed consumed, weight gain (g), average daily gain (g fish day⁻¹), specific growth rate (SGR% day⁻¹), feed conversion ratio (FCR), protein efficiency ratio (PER) and net protein utilization (NPU%).

Blood assays

Blood samples were collected using heparinized syringes from the caudal vein of the experimental fish at the end of the growth trial. Blood samples were centrifuged at 3000 rpm ×15 min at 4°C to allow separation of plasma which use to determine the blood parameters. Total plasma protein were carried out using Colorimetric method, (Roch Diagnostics, GmbH, Monnheim, Germany) as recorded by Ruane et al. [38]. Creatinine was determined according to Pincus [39]. The activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were carried out using Colorimetric method, Roch Diagnostics, GmbH, Monnheim, Germany kits according to Reitman and Frankel [40].

Statistical analysis

One way Analysis of Variance (ANOVA) was applied to test the effects of partially and totally replacement of poultry-by product meal on various growth parameters, chemical composition, blood constituents and apparent digestibility coefficients according to Snedecore and Cochran [41]. Duncan Multiple Range test was used to detect the significant differences between the means of treatments [42]. All analysis were performed using SAS (version 9.1 2004 SAS Institute, Cary, NC, USA) [43]. The level of significance was chosen at $p \leq 0.05$ and the results are presented as a group means (n=3 per tanks in each treatments ±S.E.M.).

Results

Physicochemical characteristics

Water physicochemical characteristics (Table 1) revealed that temperature, pH, dissolved oxygen, salinity and unionized ammonia are within the optimum ranges for rearing Nile tilapia according to Wangead et al., El-Shafai et al. and Ferreira et al. [44-46]. Similar physicochemical conditions were found in all tanks.

Chemical composition of diets

As can be seen in Table 3, the four experimental diets were almost similar in protein content (30.0-30.19%) and gross energy (19.58-19.77 MJ kg⁻¹ diets). However, they differed in their amino acids contents (Table 4). The all essential amino acids met the requirements of this species as recommended by NRC.

Growth performance

As presented in Table 5 averages of initial weights ranged between 1.5 to 1.6 g/fish with insignificant differences among the dietary groups indicating the random distribution of the experimental fish among treatment groups. Concerning final weights the fish fed the tested treatments recorded an insignificant values ($P < 0.05$) between each other. The same trend was observed with total gain in weight, the daily gain and specific growth rate of fish fed the different inclusion levels (50, 75 and 100% PM) of poultry by-product meal.

As shown in Table 5 average amounts of feed intake were found

Diets					
Amino acids	RE*	PBM0	PBM50	PBM75	PBM100
Essential amino acids					
Arginine	11.8	16.8	17.4	18.2	18.3
Histidine	8.4	9.2	8.8	8.9	9.1
Lysine	14.3	14.6	14.8	14.5	14.8
Leucine	9.5	20.4	21.6	22	22.4
Isoleucine	8.7	10.8	11.3	11.7	12
Valine	7.8	13.8	14.8	15.3	15.8
Methionine +Cystine	9	9.8	9.6	10.4	12.1
Threonine	10.5	10.3	11.1	11.6	11.9
Phenyl alanine+Tyrosine	15.5	20.7	20.4	24.9	24.5
Tryptophan	2.8	2.9	3.1	3	3.1
Non essential amino acids					
Glutamic acid	-	43	42.9	43.8	43.9
Aspartic acid	-	24.3	24.2	24.9	24.5
Glycine	-	18.8	19.2	18.9	17.9
Serine	-	11.7	15	16.4	17.7
Proline	-	18.4	20.4	21.1	21.8

*Requirements according to, NRC (2011).

Table 4: Amino acids contents of the experimental diets (g kg⁻¹) n=3.

Diets				
Parameters	PBM0	PBM50	PBM75	PBM100
Initial average weight (g fish ⁻¹)	1.5 ± 0.05	1.6 ± 0.06	1.6 ± 0.05	1.5 ± 0.06
Final average weight (g fish ⁻¹)	54.6 ± 2.5	53.8 ± 1.8	54.1 ± 2.0	54.3 ± 1.5
Gain in weight (g fish ⁻¹)	53.1 ^a ± 1.1	52.2 ^a ± 1.4	52.5 ^a ± 1.5	52.8 ^a ± 1.6
Average daily gain (g fish day ⁻¹)	0.44 ± 0.15	0.43 ± 0.2	0.43 ± 0.3	0.44 ± 0.1
Specific growth rate (% day ⁻¹) ¹	3.0 ± 0.5	2.92 ± 0.4	2.93 ± 0.2	2.99 ± 0.3
Feed consumed (g fish ⁻¹)	70.0 ± 3.0	71.0 ± 2.0	71.0 ± 4.0	71.0 ± 3.0
Feed conversion ratio ²	1.31 ± 0.15	1.36 ± 0.2	1.35 ± 0.2	1.34 ± 0.1
Protein efficiency ratio ³	2.51 ± 0.2	2.44 ± 0.15	2.45 ± 0.18	2.46 ± 0.16
Net Protein Utilization (NPU%) ⁴	39.75 ± 4.0	37.10 ± 3.0	37.98 ± 3.0	38.51 ± 4.0

Values are the mean ± S.E. of triplicate groups of each treatment.

¹Specific growth rate = $100 \times (\ln \text{ final weight} - \ln \text{ initial weight}) / 120 \text{ day}$.

²Feed conversion = (feed given per fish) / (weight gain per fish).

³Protein efficiency ratio = (weight gain per fish) / (protein intake per fish).

⁴Net protein utilization (%) = (final body protein - initial body protein) / (protein intake) × 100.

Table 5: Growth performance of Nile tilapia fed the experimental diets.

to be 70.0, 71.0, 71.0 and 71.0 g for the PBM0, PBM50, PBM75 and PBM100% groups, respectively. On the other hand, the best FCR (lowest) values were obtained by the PBM0 group without significant differences ($P < 0.05$) were registered among treatments. As presented in the same table the PER and NPU values showed an insignificant values between the different groups.

Apparent digestibility coefficient

Data on apparent digestibility coefficients in present study for dry matter (DM), energy (E), crude protein (CP), fat and nitrogen free extract (NFE) are presented in Table 6. Results revealed that apparent digestibility coefficients DM, E, CP, fat and NFE were not significantly affected with the inclusion levels of poultry by-product meal.

Blood characteristics

Blood characteristics revealed that insignificant differences

Diets				
Nutrients	PBM0	PBM50	PBM75	PBM100
Dry matter	85.8 ± 3.0	85.5 ± 2.0	85.6 ± 1.0	85.4 ± 2.0
Energy	90.4 ± 4.0	89.8 ± 2.0	89.6 ± 3.0	90.2 ± 1.0
Protein	94.4 ± 3.0	94.5 ± 4.0	94.3 ± 2.0	94.6 ± 3.0
Fat	95.5 ± 2.0	95.3 ± 2.0	95.4 ± 3.0	95.6 ± 3.0
Nitrogen free extract	71.8 ± 1.0	71.9 ± 3.0	71.6 ± 2.0	71.5 ± 4.0

Table 6: Apparent digestibility coefficients (%) for the experimental diets (Mean ± S.E n=3).

Diets				
Parameters	PBM0	PBM50	PBM75	PBM100
Total protein g/dl	6.4 ± 1.11	6.4 ± 0.8	6.42 ± 0.9	6.3 ± 0.6
Albumin g/dl	3.22 ± 0.33	3.24 ± 0.22	3.3 ± 0.32	3.26 ± 0.2
Urea mg/dl	7.2 ± 0.14	7.13 ± 0.12	7.3 ± 0.16	7.5 ± 0.15
Creatinine	1.12 ± 0.3	1.14 ± 0.34	1.16 ± 0.2	1.18 ± 0.26
Ast ¹	116.0 ± 0.65	117.0 ± 0.48	124.0 ± 0.56	126.0 ± 0.52
Alt ²	44.0 ± 0.41	44.0 ± 0.32	45.0 ± 0.22	44.5 ± 0.38

Values are the mean ± S.E. of triplicate groups of 20 fishes

1-Aspartate aminotransferase 2-Alanine aminotransferase

Table 7: An average of blood characteristics parameters of tilapia fed the experimental diets.

Diets					
Items	Initial	PBM0	PBM50	PBM75	PBM100
Dry matter	242 ± 1.6	271 ± 3.18	270 ± 3.6	276 ± 3.14	274 ± 1.16
Crude protein	152 ± 1.8	158 ± 2.16	152 ± 2.2	156 ± 2.2	154 ± 2.11
Crude lipid	32 ± 1.5	55 ± 1.8	62 ± 1.6	61 ± 1.4	58 ± 1.16
Crude ash	58 ± 1.2	58 ± 1.4	56 ± 1.6	59 ± 1.2	54 ± 1.6

Table 8: Carcass analysis of Nile tilapia fed the experimental diets (g kg⁻¹ wet basis) mean ± S.E. n=3.

(P<0.05) were detected between the experimental groups (Table 7) for total protein, albumin, creatinine, Aspartate aminotransferase (Ast) and Alanine aminotransferase (Alt) of tilapia, which indicate that substitution of fish meal by poultry by-product had no hazardous effects on blood parameters tested.

Carcass analysis

Results of whole fish body composition in terms of wet weight, were not significantly affected (P<0.05) by the increment of PBM levels. The chemical analysis showed that the applied dietary treatments had no significant effects (P<0.05) on whole body dry matter, crude protein, fat and ash content between groups (Table 8).

Discussion

The replacement of dietary FM in aqua feeds with readily available and more economical alternatives sources, such as poultry by-product meal is an important aim for each aquaculture industry and feed-manufacture company. This by-product of poultry processing industry is high in protein, low price and contains a favorable profile of indispensable amino acids for fish production.

The results of the present study indicated that PBM is a suitable replacement of fish meal in practical formulation diets for juvenile tilapia.

The growth performance (final weight gain, daily gain and specific growth rate) and nutrient utilization (feed conversion ratio, protein

efficiency ratio and net protein utilization) of tilapia has shown enhancement for dietary PBM without significant reduction when the replacement level of fish meal up to 100%. These findings are in agreement with other studies in tilapia [47]. However, PBM can be used to replace 75% of the fish meal in diets without amino acid supplementation for gilthead sea bream [26] up to 100%, red sea bream [27], sunshine bass [48]; grouper [49] and gible carp [50]. In contrast, fish meal could only be replaced with PBM at a level which did not exceed 50% for some marine fish species [51-53]. In the present study the growth performance and feed utilization recorded comparable results with fish meal diets. In contrast, Rawles et al. [16], recorded that PBM had lower growth than fish fed the control diet with FM in sunshine bass. They attributed that imbalance of some limiting amino acid content may have caused reduced growth performance in sunshine bass. They also speculated that reduced growth observed in this species may be due to reduced palatability for PBM. However, in the current study the balance in limiting amino acid, all diets were consumed similarly and different species may be enhance the growth performance in juvenile tilapia.

As can be seen from Table 5, the FCR, PER and NPU were not significantly differed (P<0.05) between groups and the best value was recorded by PBM0. Similar and comparable results of FCR, PER and NPU were recorded with tilapia [47,54,55].

Amongst the experimental diets in the present trial, fish receiving the PM0 and PM100 diets showed high growth performance and net protein utilization. This result is in agreement with the other several warm water finfish species obtained by Hernandez et al. and Pine et al. [47,56]. They recorded that poultry by-product has been shown to enhanced growth performance of Nile tilapia and sunshine bass.

The major reasons for different results may be due to the different fish species and the varying quality of tested PBM, which are significantly influenced by their processing methods [49].

The experimental diets in the present study showed a good digestibility coefficient for the tested diets. These results are in agreement with the results of Hernandez et al. [47]. Similar and comparable ADC values of feed dry matter, protein, lipid and energy were also observed by several authors in digestibility studies with tilapia fed the conventional commercial ingredient [55,57-60]. Similar results were recorded in rainbow trout by Burea et al. [9,23].

The carcass proximate composition of tilapia indicated that the dry matter, protein, lipid and ash were not affected by incorporation of 100% poultry by-product meal. Similar results have been reported for tilapia by Hernandez et al. [47] and sunshine bass [56], grouper [53] and gibel carp [50].

Measures of blood parameters not significantly different between treatments, where the total protein, albumin, urea, creatinine Ast and Alt are comparable with the control diet. These results are in agreement with the previous study in tilapia recorded by Metwalli, Yue et al. [61,62] and other species such as sturgeon [63] and sunshine bass [48].

Results of the present study suggest that potential replacing 100% of fish meal with poultry by-product meal in the feed of tilapia *Oreochromis niloticus*, without compromising growth performance, nutrients utilization and some blood contents. It is important to establish that alternative dietary sources to fish meal are not only supplied in the correct quantities and balance for optimal growth and feed efficiency, but can maintain optimal whole body composition and blood contents.

References

1. FAO (2010) FAO Fisheries Department Fishery Information Data and Statistics Units DISHSTAT Plus: Universal software for fishery statistical time series, Version, 23 2000 Data sets: Aquaculture production: quantities and values 1950-2009, capture production 1950-2009.
2. Vechklang K, Boonanuntanasam S, Ponchunchoovong S, Pirarat N, Wanapu C (2011) The potential for rice wine residual as alternative protein source in a practical diet for Nile tilapia (*Oreochromis niloticus*) at the juvenile stage Aquaculture Nutrition 17: 685-694.
3. Burden D (2012) Tilapia profile International and special projects Iowa State university.
4. Fitzsimmons (2014) World tilapia production 4 international trade and technical conference and exposition on tilapia, Kuala Lumpur Malaysia.
5. Naylor RL, Goldburg RB, Primavera JH, Kautsky N, Beveridge MCM, et al. (2000) Effect of aquaculture on world fish supplies Nature 405: 1017-1024.
6. New MB, Wijkstrom UN (2002) Use of fish meal and fish oil in aqua feed: further thoughts on the fish meal trap FAO fish circular No 975, FAO, Rome Italy.
7. Cavalheiro JMO, Souza EO, Bora PS (2007) Utilization of shrimp industry waste in the formulation of tilapia (*Oreochromis niloticus Linnaeus*) feed. Bioresour Technol 98: 602-606.
8. Nguyen TN, Davis DA (2009) Evaluation of alternative protein sources to replace fish meal in practical diets for Juvenile Tilapia, *Oreochromis* spp. J World Aquaculture Soc 40: 113-121.
9. Bureau DP, Harris AM, Bevan DJ, Simmons LA, Azevedo PA, et al. (2000) Feather meals and meat bone meals from different origins as protein sources in rainbow trout (*Oncorhynchus mykiss*) diets. Aquaculture 181: 281-291.
10. Millamena OM (2002) Replacement of fish meal by animal by-product meals in a practical diets for grow-out culture of grouper (*Epinephelus coioides*). Aquaculture 204: 75-84.
11. Wei Z, Kangsen M, Baigang Z, Fuzhen W, Yu U (2004) A study on the meat and bone meal and poultry by-product meal as protein substitutes of fish meal in practical diets for (*Litopenaeus vannamei*) juveniles. J Ocean Univ China (Ocean, Coast Sea Res) 3: 157-160.
12. Fasakin EA, Serwata RD, Davies SJ (2005) Comparative utilization of rendered animal derived products with or without composite mixture of soybean meal in hybrid tilapia (*Oreochromis niloticus* × *mossambicus*) diets. Aquaculture 249: 324-338.
13. Abdel-Warith A, Davies SJ, Russell P (2001) Inclusion of commercial poultry by-product meal as a protein replacement of fish meal in practical diets for the African cat fish (*Claris gariepinus*). Aquacult Res 32: 296-306.
14. Gaylord TG, Rawles SD (2005) The modification of poultry by-product meal for use in hybrid striped bass (*Morone chrysops* × *Morone saxatilis*) diets. J World Aquac Soc 36: 363-374.
15. Muzinic LA, Thompson KR, Metts LS, Dasupta S, Webster CW (2006) Use of turkey meal as partial and total replacement of fish meal in practical diets for sunshine bass (*Morone chrysops* × *Morone saxatilis*) grown in tanks Aquaculture 12: 71-81.
16. Rawles SD, Richie M, Gaylord TG, Webb J, Freeman DW, et al. (2006) Evaluation of poultry by-product meal in commercial diets for hybrid striped bass (*Morone chrysops* × *Morone saxatilis*) in recirculated tank production Aquaculture 259: 377-389.
17. Thompson KR, Metts LS, Muzinic LA, Dasupta S, Webster CD, et al. (2007) Use of turkey meal as a replacement for menhaden fish meal in practical diets for sunshine bass grown in cages N Am J Aquacult 69: 351-359.
18. Soltan MA (2009) Effect of dietary fish meal replacement by poultry by-product meal with different grain source and Enzyme supplementation on performance, feces recovery, body composition and nutrient balance of Nile tilapia. Pakistan Journal of Nutrition 8: 395-407.
19. El-Sayed AF (1998) Total replacement of fish meal with animal protein sources in Nile tilapia (*Oreochromis niloticus* L) feeds. Aquaculture 29: 275-280.
20. Yang Y, Xie SQ, Cui YB, Zhu XM, Lei W, et al. (2006) Partial and total replacement of fish meal with poultry by-product meal in diets for gibel carp, *Carassius auratus gibelio* Bloch. Aquaculture Research 37: 40-48.
21. Soaud IP, Rodgers LJ, Davis DA, Rouse DB (2008) Replacement of fish meal with poultry by-product meal in practical diets for red claw crayfish (*Cherax quadrinatus*). Aquacult Nutr 14: 139-142.
22. Rodriguez-Serna M, Olvera-Novoa MA, Carmona-Osaldá C (1996) Nutritional value of animal by-product meal in practical diets for Nile tilapia (*Oreochromis niloticus* L) fry. Aquacult Res 27: 67-73.
23. Bureau DP, Harris AM, Cho CY (1999) Apparent digestibility of rendered animal protein ingredients for rainbow trout (*Oncorhynchus mykiss*). Aquaculture 180: 345-358.
24. Fowler LG (1991) Poultry by-product meal as a dietary protein source in fall chinook salmon diets. Aquaculture 99: 309-321.
25. Alexis MN, Paparaskeva-Papoutsoglou E, Theochri V (1985) Formulation of practical diets for rainbow trout (*Salmo gairdneri*) made by partial or complete substitutes for fish meal by poultry by-product and certain plant by-products. Aquaculture 50: 61-73.
26. Nengas L, Alexis MN, Davies SJ (1999) High inclusion levels of poultry meal and related byproducts in diets for gilthead sea bream (*Sparus aurata* L). Aquaculture 179: 13-23.
27. Takag S, Hosokawa H, Shimeno S, Kawa MU (2000) Utilization of poultry by-product meal in a diet for red sea bream *Pagrus major*. Nippon Suisan Gakkaishi 66: 428-438.
28. Thompson KR, Rawles SD, Metts LS, Gannam AL, Brady YJ, et al. (2008) Digestibility of dry matter, protein, lipid and organic matter of two fish meals, two poultry by-product meals, soybean meal and distillers dried grains with solubles in practical diets for sunshine bass (*Morone chrysops* × *M saxatilis*). J World Aquac Soc 309: 352-363.
29. APHA (1992) Standard methods for the examination of water and waste water American Public Health Association, Washington, DC.
30. Jobling M (2011) National Research Council(NRC): Nutrient Requirements of fish and shrimp National Academy Press, Washington, DC, USA 20: 601-602.
31. Choubert G, De la Noue J, Luquet P (1982) Digestibility in fish: improved device for the automatic collection of faeces. Aquaculture 29: 185-189.
32. Bolin DW, King RP, Klosterman EW (1952) A simplified method for the determination of chromic oxide (Cr₂O₃) when used as an index substance. Science 116: 634-635.
33. Cho CY (1993) Digestibility of feedstuffs as a major factor in aquaculture waste management. Fish nutrition in practice. INRA, France.
34. AOAC (2006) Official methods of analysis of AOAC International (18 EDN) AOAC International Maryland, USA.
35. Cohen SA, Mamd M, Tavin TL (1989) The pico-Tag Method-A method of advanced techniques for amino acid analysis Bedford, MA, USA.
36. Schulz C, Wickert M, Kijora C, Ogunji J, Rennert B (2007) Evaluation of pea protein isolate as alternative protein source in diets for juvenile tilapia (*Oreochromis niloticus*). Aquaculture Research 38: 537-545.
37. Jobling M (1994) Fish bioenergetics, Fish and fisheries Series, Chapman & Hall-2-6 Boundary, London.
38. Ruane NM, Huisman EA, Comen J (2001) Plasma cortisol and metabolite level profiles in two isogenic strains of common carp. J Fish Biol 59: 1-12.
39. Pincus MR (1996) Interpreting laboratory results reference values and decision making, Clinical Diagnosis and Management by laboratory Methods Saunders, Philadelphia, USA.
40. Reitman A, Franks S (1957) Determination of aspartate glutamic amino transferase and alanin aminotransferase. American Journal of Clinical Pathology.
41. Snedecore, Gand W, Cochran WC (1987) Statistical Methods Iowa state Univ, USA.
42. Duncan DB (1955) Multiple ranges and multiple F tests. Biometric 11: 1-42.
43. SAS (1986) SAS User's Guide Version 6 Edition SAS Institute, Cary, NCUSA.
44. Wangead C, Greater A, Tansakul R (1988) Effects of acid water on survival and growth rate of Nile tilapia (*Oreochromis niloticus*) In: RSV Pulin, T Bhukaswan, K Tonguthai and J Maclean, Proceedings of the Second International Symposium on Tilapia in Aquaculture ICLARM Conference Proceedings No15, Department of Fisheries, Bangkok, Thailand ICLARM, Manila, Philippines.

45. El-Shafai SA, El-Gohary FA, Nasr FA, Der Steen NP, Gijzen HJ (2004) Chronic ammonia toxicity to duckweed-fed tilapia (*Oreochromis niloticus*). *Aquaculture* 232: 117-127.
46. Ferreira MW, Araujo FG, Costa DV, Rosa PV, Figueiredo HCP, et al. (2011) Influence of dietary oil sources on muscle composition and plasma lipoprotein concentration in Nile tilapia (*Oreochromis niloticus*). *Journal of World Aquaculture Society* 42: 24-33.
47. Hernandez C, Olvera-Novoa AM, Hardy RW, Hemosillo A, Reyes C, et al. (2010) Complete replacement of fish meal by poultry by-product meals in practical diets for fingerlings Nile tilapia (*Oreochromis niloticus*) digestibility and growth performance. *Aquaculture Nutrition* 16: 44-53.
48. Rawles SD, Thompson KR, Brady YJ, Metts LS Aksoy MY, et al. (2011) Effects of replacing fish meal with poultry by-product meal and soybean meal and reduced protein level on the performance and immune status of pond-grown sunshine bass (*Morone chrysops* × *Morone saxatilis*). *Aquacult Nutr* 17: e708-e721.
49. Shapawi R, Ng WK, Mostafa S (2007) Replacement of fish meal with poultry by-product meal in diets formulated for the humpback grouper (*Cromileptes altivelis*). *Aquaculture* 273: 118-389.
50. Hu M, Wang Y, Wang Q, Zhao M, Xiong B, et al. (2008) Replacement of fish meal by rendered animal protein ingredients with lysine and methionine supplementation to practical diets for gibel carp (*Carassius auratus gibelio*). *Aquaculture* 275: 260-265.
51. El-Sayed AF (1994) Evaluation of soybean meal, spirulina meal and chicken offal meal as protein sources for silver sea bream (*Rhabdosargus sarba*) fingerlings. *Aquaculture* 127: 169-175.
52. Rawles SD, Gaylord TG, McEntire ME, Freeman DW (2009) Evaluation of poultry by-product meal in commercial diets for hybrid striped bass (*Morone chrysops* × *Morone saxatilis*) in pond production. *J World Aquac Soc* 40: 141-156.
53. Wang Y, Li K, Han H, Zheng Z, Bureu DP (2008) Potential using a blend of rendered animal protein ingredients to replace fish meal in practical diets for Malabar grouper (*Epinephelus malabaricus*). *Aquaculture* 281: 133-117.
54. Gao W, Liu YGJ, Tian LX, Mai KS, Liang GY, et al. (2011) Protein sparing capability of dietary lipid in herbivorous and omnivorous freshwater finfish: a comparative case study on grass carp (*Ctenopharyngodon idella*) and tilapia (*Oreochromis niloticus* × *Oaureus*). *Aquaculture Nutrition* 17: 2-12.
55. Yones A, Abdel-Hakim NF (2010) Study on growth performance and apparent digestibility coefficients on some common plant protein ingredients used in formulated diets of Nile tilapia (*Oreochromis niloticus*). *Egyptian Journal Nutrition and Feeds* 13: 589-606.
56. Pine HJ, Daniels WH, Davis DA, Jiang M, Webster CD (2008) Replacement of fish meal with poultry by-product meal as a protein source in pond-raised sunshine bass (*Morone chrysops* × *Morone saxatilis*). *Journal World Aquaculture Society* 39: 586-597.
57. Shiao SY, Chuang JL, Sun CL (1987) Inclusion of soybean meal in tilapia (*Oreochromis niloticus* × *Oreochromis aureus*) diets at two protein levels. *Aquaculture* 65: 251-261.
58. Koprucu K, Ozdemir Y (2005) Apparent digestibility of selected feed ingredients for Nile tilapia (*Oreochromis niloticus*). *Aquaculture Research* 250: 308-316.
59. Gaye-Sliessegger J, Focken U, Abel HJ, Becker K (2005) Improving estimates of tropic shift in Nile tilapia (*Oreochromis niloticus* L) *Comp Biochem Physiol A Mol Integr Physiol* 140: 117-124.
60. Yones AM (2010) Effect of lupin kernel meal as plant protein sources in diets of red hybrid tilapia (*Oreochromis niloticus* × *O mossambicus*), on growth performance and nutrients utilization. *African Journal Biological Science* 6: 1-16.
61. Metwalli AA (2013) Effects of partial and total substitution of fish meal with corn gluten meal on growth performance, nutrients utilization and some blood constituents of the Nile tilapia (*Oreochromis niloticus*). *Egypt J Aquat Biol & fish* 17: 91-100.
62. Yue YR, Zhou QC (2008) Effect of replacing soybean meal with cottonseed meal on growth, feed utilization and hematological indexes for juvenile hybrid tilapia (*Oreochromis niloticus* × *Oaureus*). *Aquaculture* 284: 185-189
63. Zhu H, Gong G, Wang J, Wu X, Xue H, et al. (2011) Replacement of fish meal with blend of rendered animal protein in diets for Siberian sturgeon (*Acipenser baeri*), results in performance equal to fish meal fed fish. *Aquacult Nutr* 17: e389-e395.

Citation: Yones AMM, Metwalli AA (2015) Effects of Fish Meal Substitution with Poultry By-product Meal on Growth Performance, Nutrients Utilization and Blood Contents of Juvenile Nile Tilapia (*Oreochromis niloticus*). J Aquac Res Development 6: 389. doi:10.4172/2155-9546.1000389

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.com/journals/submission>