

4-((1E)-2-(5-(4-hydroxy-3-methoxystyryl)-1-phenyl-1H-pyrazoyl-3-yl) vinyl)-2-methoxy-phenol) (CNB-001) Does Not Regulate Human Recombinant Protein-Tyrosine Phosphatase 1B (PTP1B) Activity *in vitro*

Paul A Lapchak^{1,2*}, Jacqueline A Lara¹ and Paul D Boitano¹

¹Department of Neurology, Cedars-Sinai Medical Center, Advanced Health Sciences Pavilion, Los Angeles, USA

²Department of Neurosurgery, Cedars-Sinai Medical Center, Advanced Health Sciences Pavilion, Los Angeles, USA

Abstract

Protein-Tyrosine Phosphatase1B (PTP1B) is a negative regulator of the insulin signaling pathway and is a potential therapeutic target for treatment of type 2 diabetes, cardiovascular disease, metabolic syndrome and cancer. It has been postulated that CNB-001 [4-((1E)-2-(5-(4-hydroxy-3-methoxystyryl)-1-phenyl-1H-pyrazoyl-3-yl) vinyl)-2-methoxy-phenol)] may regulate PTP1B activity suggested by a computer-based active site docking recognition model. This possibility was studied using a human recombinant PTP1B assay, and a phospho-peptide fragment of the insulin receptor β subunit domain (IR5). The positive control, suramin, inhibited PTP1B with an IC₅₀ (half minimal (50%) inhibitory concentration) value of 16.34 μ M; CNB-001 did not affect enzyme activity across the range of 1nM-0.1mM. This study suggests that PTP1B inhibition is not involved in the beneficial effects of CNB-001 in obese type 2 diabetic mice.

Keywords: Curcumin analog; Neuroprotection; Metabolic disorder; Neurotrophic; Phosphatases; Pleiotropic

Introduction

CNB-001 [4-((1E)-2-(5-(4-hydroxy-3-methoxystyryl)-1-phenyl-1H-pyrazoyl-3-yl)vinyl)-2-methoxy-phenol], is an important first in class pleiotropic drug candidate being developed to treat neurodegenerative diseases such as acute ischemic stroke and Alzheimer's disease [1-4]. Using phenotypic screening assays directed against some of the exacerbating mechanisms (mitochondrial dysfunction, oxidative stress glutamate toxicity) underlying deficits resulting from activation of the stroke cascade, CNB-001 has been shown to support cell survival [5], CNB-001 also a potent 5-lipoxygenase inhibitor (5-LOX) [4], anti-apoptotic and antioxidant [6], a negative regulator of inflammation [down-regulates, 5-LOX, cyclooxygenase-2 (COX-2), interleukin-6 (IL-6)] [3,7,8], and an activator of brain-derived neurotrophic factor (BDNF) and its signaling pathways [3]. The pleiotropic nature of the drug makes an optimal candidate for cellular protection and repair for a variety of diseases that currently have limited treatment options [2,3,7,9-16].

We recently studied the pharmacological effects of CNB-001 in a diabetic mouse model (7), a model of obesity-associated insulin resistance induced by a high fat diet. The study found that CNB-001 effectively attenuated or reversed many of the deleterious changes including weight gain, serum triglycerides and serum IL-6 levels. Moreover, in obese mice, CNB-001 increased energy expenditure and high-fat insulin resistance, glucose tolerance and reversed gastrocnemius muscle deficits (i.e.: 2-deoxyglucose uptake). Importantly, in the study, Panzhinskiy et al. [12] hypothesized that CNB-001 may regulate Protein-Tyrosine Phosphatase1B (PTP1B) based upon the best fit of the molecule using computer-based active site docking recognition. PTP1B is a negative regulator of the insulin signaling pathway and is a potential therapeutic target, in particular for treatment of type 2 diabetes [12-16]. Since preliminary studies suggested that CNB-001 may potentially bind to PTP1B and regulate PTP1B activity, an effect of CNB-001 on the phosphatase may explain some of the beneficial effects that were observed.

Thus, using a human recombinant PTP1B assay, and a phospho-peptide fragment of the insulin receptor β subunit domain, we tested the

hypothesis that CNB-001 interacts with the enzyme. In this controlled study, we used suramin as the positive control inhibitor [17].

Materials and Methods

Drug preparation: CNB-001 (AQ BioPharma Co. Ltd, Shanghai, China) was dissolved in reagent grade dimethyl sulfoxide (DMSO) (Sigma-Aldrich, Saint Louis, MO) for assay.

Enzyme assay: The PTP1B Colorimetric Assay Kit was purchased from Enzo Life sciences International Inc., Plymouth Meeting, PA. The assay kit contained recombinant PTP1B enzyme (residues 1-322 M.W. 37,400) expressed in E. coli. The phosphopeptide substrate (IR5) used in the sensitive assays contains the amino acids 1142-1153pY1146 (MW-1703 kDa), a sequence from the insulin receptor β subunit domain that must be autophosphorylated to achieve full receptor kinase activation. This "activation loop" is the target of several protein phosphatase regulators of insulin signaling, including, notably, PTP1B. The enzyme has an estimated Km of 85 μ M for the PTP1B phospho-peptide substrate (IR5) which is used to measure PTP1B activity.

Methods: Since the assay endpoint is the measurement of free phosphate, a phosphate standard curve was prepared per the manufacturers protocol; six concentrations of inorganic phosphate were established in duplicate wells: 0, 0.25 nmol, 0.5 nmol, 1.0 nmol,

***Corresponding author:** Paul A Lapchak, Director of Translational Research, Cedars-Sinai Medical Center Professor, Department of Neurology and Neurosurgery, Advanced Health Sciences Pavilion, Suite 8305, 127 S. San Vicente Blvd, Los Angeles, USA, Tel: 310-248-8188; Fax: 310-248-7568, E-mail: Paul.Lapchak@cshs.org

Received August 18, 2014; **Accepted** September 22, 2014; **Published** September 29, 2014

Citation: Lapchak PA, Lara JA, Boitano PD (2014) 4-((1E)-2-(5-(4-hydroxy-3-methoxystyryl)-1-phenyl-1H-pyrazoyl-3-yl) vinyl)-2-methoxy-phenol) (CNB-001) Does Not Regulate Human Recombinant Protein-Tyrosine Phosphatase 1B (PTP1B) Activity *in vitro*. J Neurol Neurophysiol 5: 232. doi:[10.4172/2155-9562.1000232](http://dx.doi.org/10.4172/2155-9562.1000232)

Copyright: © 2014 Lapchak PA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

2.0nmol and 3.0nmol by adding the appropriate amounts of assay buffer and phosphate standard. All test sample/inhibitor assays were performed following the protocol, with variances only in the concentrations of the reagents (as described): PTP1B enzyme was prepared for all assays to allow for a final amount of 2.5 ng/well and IR5 was prepared to have a final assay concentration of 75 μ M. Briefly; 35 μ l of 1X assay buffer (100 mM MES, pH 6.0, 300 mM NaCl, 2 mM EDTA, 2 mM DTT, 0.1% NP-40) was added to each well and incubated for 15 minutes at 39°C; 10 μ l of CNB-001 or DMSO as baseline control (final concentrations 1.0 mM, 0.1 mM, 10 μ M, 1 μ M, 10nM and 1.0nM) were added to each well followed by 5 μ l of PTP1B enzyme (2.5ng per well) dilution. In experiments using suramin, it was dissolved in water, and 10 μ l of suramin or water were added to each well. Reactions were initiated by adding 50 μ l of IR5 substrate (final concentration 75 μ M) and the plate was incubated for 30 minutes at 39°C. Reactions were then terminated by adding 25 μ l of the provided phosphate detection reagent (Red Reagent) and wells were agitated gently to mix. Color was allowed to develop for 25-30 minutes, and absorbance was read at 620 nm using a Spectra Max M2 spectrophotometer (Molecular Devices, Sunnyvale, CA).

Results: In this controlled study, we used suramin, a reversible and competitive inhibitor of PTP1B to establish the characteristics of the assay [17]. Suramin has a K_i of approximately 5.5 μ M and an IC_{50} value in the range of 9.5-11 μ M for PTP1B inhibition, as reported in the literature (17). In the suramin study, the standard curve was linear with a regression equation of $y = -0.013 + 0.2086 B$ and a regression analysis fit of $R^2 = 0.9972$. Thus, phosphate product amounts could easily be calculated.

In Figure 1, we provide data demonstrating the effect of suramin on PTP1B activity. Using suramin concentration of 1.0-nM-0.10 mM we constructed an inhibition curve, we show that low doses of suramin (1.0 nM to 0.1 μ M) had no effect on PTP1B activity. However, higher doses of suramin inhibited PTP1B and phosphate production. In our assay, the IC_{50} (half minimal (50%) inhibitory concentration) value was 16.34 μ M for 50% PTP1B inhibition.

In Figure 2, we provide data for the effects of CNB-001 on PTP1B across the concentration range of 1.0-nM-0.10 mM. In the CNB-001 study, the standard curve was linear with a regression equation of $y =$




Figure 1: Suramin inhibits PTP1B with an IC_{50} of 16.34 μ M; PTP1B assays were run in triplicate with either buffer or suramin (0.1mM-1nM). Significantly different (** $p < 0.01$) from baseline enzyme activity levels measured by phosphate product produced over 30 minutes. IC_{50} value is calculated inhibitory concentration required for 50% enzyme activity inhibition compared to baseline control. Paired-t-test was used for statistical analysis (Graph Pad Inc.).




Figure 2: CNB-001 does not affect PTP1B activity *in vitro*; PTP1B assays were run in triplicate with either DMSO or CNB-001 (0.1mM-1nM). Activity measured in the presence of CNB-001 was not significantly different ($p > 0.05$) from baseline enzyme activity levels measured by phosphate product produced over 30 minutes. Baseline was lowered by the presence of DMSO in the assay. Paired-t-test was used for statistical analysis (Graph Pad Inc.).

$-0.010 + 0.1349 B$ a regression analysis fit of $R^2 = 0.9821$. Thus, even in the presence of DMSO, phosphate product amounts could easily be calculated. Note that in the presence of DMSO, the amount of phosphate detected using the RED reagent was reduced compared to the phosphate detected in the absence of DMSO (i.e.: buffer alone). Nevertheless, for each drug, we constructed standard curves using the vehicles used to solubilize each drug so that regression analysis could be conducted. In this assay, CNB-001 did not have any effect on PTP1B activity.

Conclusion

We tested the hypothesis that CNB-001 interacts with PTP1B to produce the beneficial effects observed in obese diabetic mice. Using a specific recombinant human PTP1B assay, under conditions recommended by the manufacturer, we found that CNB-001 did not regulate PTP1B activity, but suramin effectively inhibited enzyme activity with an IC_{50} similar to that previously documented in the literature [17]. Based upon this data, it does not appear that CNB-001 regulates PTP1B activity, and CNB-001 is not an inhibitor of PTP1B.

References

1. Lapchak PA, McKim JM Jr (2011) CeeTox™ Analysis of CNB-001 a Novel Curcumin-Based Neurotrophic/Neuroprotective Lead Compound to Treat Stroke: Comparison with NXY-059 and Radicut. Transl Stroke Res 2: 51-59.
2. Lapchak PA (2011) Neuroprotective and neurotrophiccurcuminoids to treat stroke: a translational perspective. Expert Opin Investig Drugs 20: 13-22.
3. Lapchak PA, Schubert DR, Maher PA (2011) Delayed treatment with a novel neurotrophic compound reduces behavioral deficits in rabbit ischemic stroke. J Neurochem 116: 122-131.
4. Valera E, Dargusch R, Maher PA, Schubert D (2013) Modulation of 5-lipoxygenase in proteotoxicity and Alzheimer's disease. J Neurosci 33: 10512-10525.
5. Liu Y, Dargusch R, Maher P, Schubert D (2008) A broadly neuroprotective derivative of curcumin. J Neurochem 105: 1336-1345.
6. Jayaraj RL, Tamilselvam K, Manivasagam T, Elangovan N (2013). Neuroprotective effect of CNB-001, a novel pyrazole derivative of curcumin on biochemical and apoptotic markers against rotenone-induced SK-N-SH cellular model of Parkinson's disease. J Mol Neurosci 51: 863-870.
7. Panzhinskiy E, Hua Y, Lapchak PA, Topchiy E, Lehmann TE, et al. (2014) Novel curcumin derivative CNB-001 mitigates obesity-associated insulin

- resistance. J Pharmacol Exp Ther 349: 248-257.
8. Narumoto O, Matsuo Y, Sakaguchi M, Shoji S, Yamashita N, et al. (2012) Suppressive effects of a pyrazole derivative of curcumin on airway inflammation and remodeling. Exp Mol Pathol 93: 18-25.
 9. Elgebaly MM, Ogbi S, Li W, Mezzetti EM, Prakash R, et al. (2011) Neurovascular injury in acute hyperglycemia and diabetes: A comparative analysis in experimental stroke. Transl Stroke Res 2: 391-398.
 10. Moskowitz MA, Lo EH, Iadecola C (2010) The science of stroke: mechanisms in search of treatments. Neuron 67: 181-198.
 11. Lo EH, Rosenberg GA (2009) The neurovascular unit in health and disease: introduction. Stroke 40: S2-3.
 12. Combs AP (2010) Recent advances in the discovery of competitive protein tyrosine phosphatase 1B inhibitors for the treatment of diabetes, obesity, and cancer. J Med Chem 53: 2333-2344.
 13. Shinde RN, Sobhia ME (2013) Binding and discerning interactions of PTP1B allosteric inhibitors: novel insights from molecular dynamics simulations. Journal of molecular graphics & modeling 45: 98-110.
 14. Tang YB, Lu D, Chen Z, Hu C, Yang Y, et al. (2013) Design, synthesis and insulin-sensitising effects of novel PTP1B inhibitors. Bioorg Med Chem Lett 23: 2313-2318.
 15. Liu Z, Chai Q, Li YY, Shen Q, Ma LP, et al. (2010) Discovery of novel PTP1B inhibitors with antihyperglycemic activity. Acta Pharmacol Sin 31: 1005-1012.
 16. Tobin JF, Tam S (2002) Recent advances in the development of small molecule inhibitors of PTP1B for the treatment of insulin resistance and type 2 diabetes. Curr Opin Drug Discov Devel 5: 500-512.
 17. McCain DF, Wu L, Nickel P, Kassack MU, Kreimeyer A, et al. (2004) Suramin derivatives as inhibitors and activators of protein-tyrosine phosphatases. J Biol Chem 279: 14713-14725.

Citation: Lapchak PA, Lara JA, Boitano PD (2014) 4-((1E)-2-(5-(4-hydroxy-3-methoxystyryl)-1-phenyl-1H-pyrazol-3-yl) vinyl)-2-methoxy-phenol (CNB-001) Does Not Regulate Human Recombinant Protein-Tyrosine Phosphatase1B (PTP1B) Activity *in vitro*. J Neurol Neurophysiol 5: 232. doi:[10.4172/2155-9562.1000232](https://doi.org/10.4172/2155-9562.1000232)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>

