

Elasmobranch Distribution and Assemblages in the Southern Tyrrhenian Sea (Central Mediterranean)

Bottari T, Busalacchi B, Profeta A, Mancuso M*, Giordano D and Rinelli P

Institute of Coastal Marine Environment (IAMC) - CNR, Spianata S. Raineri, 86 - 98122 Messina, Italy

Abstract

The aim of this study is to identify elasmobranch species present in the Southern Tyrrhenian Sea, to describe their distribution and abundance, to identify significant spatial or temporal differences between species. 14 bottom trawl surveys were carried out from 1994 to 2007. 16 species of elasmobranchs were recorded. Multidimensional Scaling Ordination (MDS) showed two groups according to the depth gradient: in the first one the stations from upper slope and in the second one the stations from middle slope. Mean biomass indices and frequency of occurrence showed that *Galeus melastomus*, *Etmopterus spinax* and *Scyliorhinus canicula* were the most abundant species. Mean biomass indices for other species were very low. The mean abundance of *G. melastomus* exhibited a positive temporal trend in biomass and density. The mean abundance of *E. spinax* exhibited a negative temporal trend in biomass and in density. For other species abundance varied greatly between years, even if there was no evident trend.

Keywords: Sharks; Ecosystem; Trawl surveys; Multivariate analysis

Introduction

In the Mediterranean Sea there is a high level of exploitation due to a great variety of fishing gears; generally the elasmobranchs are not targeted by commercial fisheries but they are an important by-catch especially of the trawl fishery and deep-water long liners [1-4].

The decreases in abundance and biomass of some elasmobranch species throughout the last decade have been reported in the Gulf of Lions [5,6].

The role of these species as indicators of fishing pressure has been suggested [4,7,8] and management strategies are necessary to minimize significantly the chondrichthyan by-catch.

This paper characterizes the assemblages of demersal elasmobranch on the bottom trawl fishing grounds along the Southern Tyrrhenian Sea. Experimental trawl surveys are analysed for the main species in terms of species composition, community structure and distribution.

This paper could be useful for monitoring future trends in the same area and would allow comparison with other seas.

Materials and Methods

Study area and sampling design

Data here reported come from 14 bottom trawl surveys, carried out during the MEDITS Project, from 1994 to 2007 in the Southern Tyrrhenian Sea. The study area extended from Cape Suvero to Cape S. Vito (Figure 1), within the isobath of 800 metres, for a total area of 7256 km². Only 65% (4716 km²) of the total area studied can be trawled by commercial vessels. The fishing fleet is represented by 80 trawlers providing about 5000 tons of fish, molluscs and crustaceans (IREPA2008).

The bottom of this area is characterized by a narrow continental shelf, sometimes entirely missing and by a steep slope [9]. Sampling procedures were the same in all surveys, according to MEDITS project protocol [10]. Sampling was carried out randomly and the hauls were proportionately distributed in five bathymetric strata: stratum A: 10-50 metres (622 km²); stratum B: 51-100 metres (1003 km²); stratum C: 101-200 metres (1224 km²); stratum D: 201-500 metres (1966 km²); stratum E: 501-800 metres (2441 km²). A total of 360 hauls were carried

out. An experimental sampling gear with a cod-end mesh size of 20 millimetres was used. The fishing speed was 3 knots. The horizontal and vertical openings of the net (on average 18.4 and 1.90 m respectively) were measured using a SCAMMAR system. The haul length was 30 min in the shelf (10-200 m), and 60 min in the slope (>200 m). All elasmobranch species caught were identified, counted and weighed.

Data processing

Spatial and abundance analyses were employed to investigate temporal trend. In particular, two abundance indexes, mean density index (DI; N/km²) and mean biomass indices (BI; kg/km²), were estimated (for each stratum and overall area) according to the swept-area principle [11].

Data were analysed in terms of multivariate analysis using the package Primer v6 [12]. Analyses were carried out on density index. In order to examine the demersal elasmobranch assemblages distribution along space, time and depth, the Multi Dimensional Scaling (MDS) ordination method was employed. The Bray-Curtis similarity index was used on square root transformed data.

The analysis of Similarity (ANOSIM) [13] was applied to detect differences between depths (strata) and time (years). The typifying and discriminating species of the MDS stations were determined using the similarity percentage analysis (SIMPER) [13].

The DI by haul were interpolated and mapped for the three most abundant species. The GIS software ArcMap™ 9.0 (ESRI) was used. An "exact interpolator" (Inverse Distance Weighted) was employed to

***Corresponding author:** Monique Mancuso, Institute of Coastal Marine Environment (IAMC) - CNR, Spianata S. Raineri, 86-98122 Messina, Italy, Tel: 0906015438; E-mail: monique.mancuso@iamc.cnr.it

Received November 26, 2013; **Accepted** December 30, 2013; **Published** January 07, 2014

Citation: Bottari T, Busalacchi B, Profeta A, Mancuso M, Giordano D, et al. (2014) Elasmobranch Distribution and Assemblages in the Southern Tyrrhenian Sea (Central Mediterranean). J Aquac Res Development 5: 216 doi:10.4172/2155-9546.1000216

Copyright: © 2014 Bottari T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Map of the study area.

Species	Family	Order	Common name
<i>Galeus melastomus</i> Rafinesque, 1810	Scyliorhinidae	Carchariniformes	Blackmouth catshark
<i>Scyliorhinus canicula</i> (Linnaeus, 1758)	Scyliorhinidae	Carchariniformes	Small-spotted catshark
<i>Scyliorhinus stellaris</i> (Linnaeus, 1758)	Scyliorhinidae	Carchariniformes	Nursehound
<i>Squalus acanthias</i> Linnaeus, 1758	Squalidae	Squaliformes	Piked dogfish
<i>Etmopterus spinax</i> (Linnaeus, 1758)	Etmopteridae	Squaliformes	Velvet belly
<i>Dalatias licha</i> (Bonnaterre, 1788)	Dalatiidae	Squaliformes	Kitefin shark
<i>Dasyatis pastinaca</i> (Linnaeus, 1758)	Dasyatidae	Rajiformes	Common stingray
<i>Pteroplatytrygon violacea</i> (Bonaparte, 1832)	Dasyatidae	Rajiformes	Pelagic stingray
<i>Myliobatis aquila</i> (Linnaeus, 1758)	Myliobatidae	Rajiformes	Common eagle ray
<i>Dipturus oxyrinchus</i> (Linnaeus, 1758)	Rajidae	Rajiformes	Longnosed skate
<i>Raja clavata</i> Linnaeus, 1758	Rajidae	Rajiformes	Thornback ray
<i>Raja miraletus</i> Linnaeus, 1758	Rajidae	Rajiformes	Brown ray
<i>Raja montagui</i> Fowler, 1910	Rajidae	Rajiformes	Spotted ray
<i>Raja polystigma</i> Regan, 1923	Rajidae	Rajiformes	Speckled ray
<i>Torpedo marmorata</i> Risso, 1810	Torpedinidae	Torpenidiformes	Spotted torpedo
<i>Torpedo torpedo</i> (Linnaeus, 1758)	Torpedinidae	Torpenidiformes	Common torpedo

Table 1: Elasmobranch species identified in the Southern Tyrrhenian Sea during the Medit project from 1994 to 2007.

		Species	Av. abundance	Av. Similarity	sim/SD	
D stratum: 201 - 500 m	Av. Similarity: 48.75	<i>Galeus melastomus</i>	3.1	32.24	1.34	
		<i>Scyliorhinus canicula</i>	1.78	13.2	0.81	
E stratum: 501 - 800 m	Av. Similarity: 72.36	<i>Galeus melastomus</i>	3.28	40.95	2.62	
		<i>Etmopterus spinax</i>	2.41	30.68	2.28	
		Species	D stratum Average abundance	E stratum Average abundance	Av. dissimilarity	Diss/SD
D stratum: 201 - 500 m vs E stratum: 501-800 m	Av. Dissimilarity: 56.6	<i>Etmopterus spinax</i>	0.21	2.41	17.68	2.17
		<i>Galeus melastomus</i>	3.1	3.28	13.64	1.28
		<i>Scyliorhinus canicula</i>	1.78	0.38	13.26	1.27
		<i>Scyliorhinus stellaris</i>	0.75	0.08	6.21	0.54
		<i>Torpedo marmorata</i>	0.46	0	3.8	0.54

Table 2: SIMPER analysis results.

render back the real value in every sample site of the studied area [14]. Mean distribution maps were produced pooling the records of surveys from 1994 to 2007.

The correlation between DI and BI values and years was tested using the Spearman's rank-order correlation (ρ).

The distribution of elasmobranches in relation to depth was analyzed comparing the density index by Kruskal-Wallis test.

Results

16 elasmobranch species were recorded (Table 1). The orders

observed were: Rajiformes (8 species), Squaliformes (3 species), Carcharhiniformes (3 species) and Torpediniformes (2 species).

MDS ordination (Figure 2) showed the stations reporting the stratum. Two main groups were distinguished: in the first one the stations from upper slope (201-500 metres) and in the second one the stations from middle slope (501-800 metres) (ANOSIM test: Global R=0.6; $p < 0.01$).

To individuate the importance of time in order to discriminate the assemblages one-way ANOSIM tests were performed for “year” factor across all “stratum” groups respectively. Each depth stratum was treated separately. There were no differences, across “stratum” groups and between “year” groups (Global R: 0.182 $p > 0.05$).

The results of the SIMPER analysis highlighted the species that mainly contribute as a percentage to similarity within groups “stratum” (Table 2). *Galeus melastomus* and *Scyliorhinus canicula* were important in typifying the demersal fish community of D stratum (201-500 m). The analysis performed on E stratum (501-800 m) showed that, although *G. melastomus* was still present, *Etmopterus spinax* also contributed to typifying the group.

SIMPER analysis indicates dissimilarity between assemblages (average dissimilarity: 56.6). The pool of species responsible for discriminating these groups was mainly constituted by *E. spinax*, *G. melastomus* and *S. canicula*.

In Figure 3 distribution is shown of the three most abundant elasmobranch species in the Southern Tyrrhenian Sea over 14 years. *G. melastomus* and *E. spinax* are more abundant in the eastern and western part of the study area, specifically outside the Gulf of S. Eufemia and in the areas bordering the Gulf of Castellammare; these species were also abundant in an area localized between Sicily and Calabria, in correspondence with the Strait of Messina. *G. melastomus* and *E. spinax* were also present inside and outside the Gulf of Patti respectively.

S. canicula was more equally distributed in the study area and like the other two species, was particularly present outside the Strait of Messina.

The mean abundance of *G. melastomus* exhibited a positive temporal trend in biomass and density (Spearman ρ : 0.543, $p < 0.05$ and ρ : 0.587, $p < 0.05$ respectively). Instead the mean abundance of *E. spinax*




Figure 3: Distribution of (a) *Galeus melastomus*, (b) *Etmopterus spinax*, and (c) *Scyliorhinus canicula* along the Southern Tyrrhenian Sea over 14 years.




Figure 2: Non-metric Multidimensional Scaling Ordination (MDS) of the sampling stations according to depth. Triangles are the hauls in the upper slope (200 – 500 metres); circles are the hauls in the middle slope (500 – 800 metres).

exhibited a negative temporal trend in biomass (Spearman ρ : -0.565, $p < 0.05$) and in density (Spearman ρ : -0.661, $p < 0.05$). For other species, abundance varied greatly between years (Figure 4).

Abundance appeared to be greatest for *G. melastomus* deeper than 400 m, for *E. spinax* deeper than 450 m and for *S. canicula* at 300-450 m. However, differences in abundance were statistically significant only for *G. melastomus* (Kruskal Wallis H: 36.1, $p < 0.01$). and *E. spinax* (Kruskal Wallis H: 57.1, $p < 0.01$). In comparison *S. canicula* was more evenly distributed with respect to depth (Kruskal Wallis H: 14.7, $p > 0.05$) (Figure 5).

Discussion

The analysis of demersal elasmobranch species in the Southern Tyrrhenian Sea has shown that demersal elasmobranch assemblages are aligned with depth. These results are similar to those obtained in the Atlantic Ocean [15] and in the Western Mediterranean [16] The most important boundary, located around 500 m, separates the species of the upper slope from those of the middle slope down to 800 m. The bathymetric boundaries are similar with those obtained in previous




Figure 4: Mean abundance (N/km²) and standard error of the most abundant species from the trawl surveys of 1994-2007.

studies of demersal assemblages carried out in the Southern Aegean [17] and North West Mediterranean [18]. Data reported here suggests that the structure of assemblages does not change in over a period of 14 years.

The upper slope assemblage is characterised by *G. melastomus* and *S. canicula*. The middle slope assemblage is characterised also by *G. melastomus*, a species with a wide depth range, and *E. spinax*, a species present only in this assemblage.

Some papers on slope assemblages demonstrate the importance of factors associated to depth rather than depth itself to affect the assemblages structure [19-21]. The depth in fact, may affect other environmental factors, both biological (trophic factors, interspecific competition, predator-prey relationship) and physical (steepness of the continental slope, substrate type, hydrographic condition, dissolved oxygen, light intensity) [22-24]. Recently, shark assemblages changes, related to temperature and salinity modifications of deep-water masses, have been also reported [25].

The significant increase in density and biomass indices shown by *G. melastomus* has already been reported for the species in the South of Sicily [26,27].

The opposite abundance temporal trends shown by *G. melastomus*

and *E. spinax* could be explained by comparing their life history traits. *Galeus melastomus* is a multiple oviparous whereas *Etmopterus spinax* is a ovoviviparous. The study of length at first maturity has revealed that *E. spinax* is a late-maturing species [28] and this fact makes these species more vulnerable to exploitation. Also, they do not have the same “catchability” to trawl fishing. The size distribution varies with depth for both species with larger specimens occurring at deeper waters and the smaller ones at shallower waters [16,26,29]. Moreover, the wider vertical distribution of *G. melastomus* (lower than 1000 m) might mitigate the fishing pressure as the species lives beyond the usual deepest commercial trawling limit [26,27]. *G. melastomus* shows an increasing abundance in the Southern Tyrrhenian Sea despite the persistent trawling activities [30].

Cartilaginous fish represent a good fraction (about one third) of the by-catch of red-shrimp fishing in the South of Sicily. The most common species are *G. melastomus* and *E. spinax* which are caught in over 90% of the hauls [31]. Generally they are discarded immediately after capture but elasmobranchs may die after capture because of the sudden pressure changes and handling on board [32,33]. We can hypothesize that *E. spinax* may be more sensitive to these events than *G. melastomus*.

Nevertheless, factors such as competition, changes in oceanographic conditions and changes in food abundance could affect species abundance.

There is lack of studies about distribution and structure of fish




Figure 5: Mean abundance (N/km²) and standard error for the most abundant species in each depth interval in the Southern Tyrrhenian Sea from the trawl surveys of 1994-2007.

communities in a long temporal scale in most of Mediterranean regions; such studies are necessary for ecosystem based management. We argue that results here reported could be useful as basis for management of fishery activities in the Tyrrhenian Sea.

Acknowledgement

This paper is a result of the MEDITS Project. We are grateful to all the participants in the Trawl Surveys as well as to the captain and crew of the trawler "Pasquale e Cristina" for their help during the sampling.

References

- Vacchi M, Notabartolo Di Sciara G (2000) The Cartilaginous Fishes in Italian Seas, A Resource that Urges to be Protected. *Biol Mar Medit* 7: 296-311.
- Bertrand J, Gil De Sola L, Papaconstantinou C, Relini G, Souplet A (2000) Contribution on the Distribution of Elasmobranchs in the Mediterranean from the MEDITS Surveys. *Biol Mar Medit* 7 : 1-15.
- Coelho R, Bentes L, Goncalves J, Lino PG, Ribeiro J, Erzini K (2003) Reduction of Elasmobranch By-Catch in the Hake Semipelagic Near-Bottom Longline Fishery in the Algarve (Southern Portugal). *Fish Sci* 69: 293-299.
- Carbonell A, Alemany F, Merella P, Quetglas A, Román E (2003) The By-Catch of Sharks in the Western Mediterranean (Balearic Islands) Trawl Fishery. *Fish Res* 61: 7-18.
- Aldebert Y (1997) Demersal Resources of the Gulf of Lions (Mediterranean). Impact on Fish Diversity. *Vie Milieu* 47: 275-284.
- Bertrand JA, Aldebert Y, Souplet A (1998) Temporal Variability of Demersal Species in the Gulf of Lions from Trawl Surveys (1983-1997). *IFREMER Actes De Colloq*. 26: 153-164.
- Stevens JD, Bonfil R, Dulvy NK, Walker PA (2000) The Effects of Fishing on Sharks, Rays, and Chimaeras (Chondrichthyan), and Implications for Marine Ecosystems. *ICES J Mar Sci* 57: 476-494.
- Rodríguez-Cabello C, Sánchez F, Serrano A, Olaso I (2008) Effects of Closed Trawl Fishery Areas on Some Elasmobranch Species in the Cantabrian Sea. *Journal of Marine Systems* 72:418-428.
- Greco S, Rinelli P, Giordano D, Perdichizzi F (1998) Valutazione Delle Risorse Demersali Da Capo Suvero A San.
- Capo VL (Tirreno Meridionale). *Biol Mar Medit* 5: 74-84.
- Anonymous (2012) International MEDITS-Handbook. Revision N. 6, MEDITS Working Group.
- Gunderson DR (1993) Surveys of Fisheries Resources. John Wiley & Sons, New York: 248 .
- Clarke KR, Gorley RN (2006) Primer V6: User Manual/Tutorial. PRIMER-E, Plymouth.
- Clarke KR (1993) Non-Parametric Multivariate Analyses of Changes in Community Structure. *Aust J Ecol* 18: 117-143.
- Isaaks EH, Srivastava RM (1989) The Effects of Fishing on Marine Ecosystems. *Adv Mar Biol* 34: 201-352.
- Roel BA (1978) Demersal Communities off The West Coast of South Africa. *S Afr J Marine Sci* 5: 575-584.
- Massuti E, Moranta J (2003) Demersal Assemblages and Depth Distribution of Elasmobranchs from the Continental Shelf and Slope off The Balearic Island (Western Mediterranean). *ICES J Mar Sci* 60: 753-766.
- Tserpes G, Peristeraki P, Potamias G, Tsimenides N (1999) Species Distribution in the Southern Aegean Sea Based On Bottom-Trawl Surveys. *Aquat Living Resour* 12 : 167-175.
- Massuti E, Reñones O (2005) Demersal Resource Assemblages in the Trawl Fishing Grounds Off The Balearic Islands (Western Mediterranean). *Sci Mar* 69: 167-181.
- Stefanescu C, Lloris D, Rucabado J (1993) Deep-Sea Fish Assemblages in The Catalan Sea (Western Mediterranean) Below A Depth of 1000 M. *Deep-Sea Res* 40 : 695-707.
- Massuti E, Gordon JDM, Moranta J, Swan SC, Stefanescu C, Merrett NR (2004) Mediterranean and Atlantic Deep-Sea Fish Assemblages: Differences in Biomass Composition and Size-Related Structure. *Sci Mar* 68 : 101-115.
- Cartes JE, Maynou F, Lloris D, Gil De Sola L, Garcia M (2009) Influence of Trawl Type on the Composition and Diversity of Deep Benthopelagic Fish and Decapod Assemblages Off the Catalan Coasts (Western). *Sci Mar* 73 : 725-737.
- Hecker B (1990) Variation in Megafaunal Assemblages on the Continental Margin South of New England. *Deep-Sea Res* 37: 37-57.
- Smale MJ, Roel BA, Badenhorst A, Field F (1993) Analysis of The Demersal Community of Fish and Cephalopods on The Agulhas Bank, South Africa. *J Fish Biol* 43 (Suppl.A): 169-191.
- Sardà F, Cartes JE, Company JB (1994) Spatio-Temporal Variation in Megabenthos Abundance in Three Different Habitats of the Catalan Deep-Sea (Western Mediterranean). *Mar Biol* 120: 211-219.
- Cartes JE, Fanelli E, Lloris D, Matallanas J (2013) Effect of Environmental Variations on Sharks and Other Top Predators in the Deep Mediterranean Sea over the Last 60 Years. *Clim Res* 55: 239-251.
- Ragonese S, Nardone GD, Ottonello D, Gancitano S, Giusto GB, Sinacori G (2009) Distribution and Biology of the Blackmouth Catshark *Galeus Melastomus* in the Strait of Sicily. *Medit Mar Sci* 10: 55-72.
- Ragonese S, Vitale S, Dimech M, Mazzola S (2013) Abundances of Demersal Sharks and Chimaera from 1994-2009 Scientific Surveys in the Central Mediterranean. *Plos ONE* 8: E74865.
- Coelho R, Erzini K (2005) Length at First Maturity of Two Species of Lantern Sharks (*Etmopterus Spinax* and *Etmopterus Pusillus*) Off Southern Portugal. *J Mar Biol Ass UK* 85: 1163-1165.
- Rinelli P, Bottari T, Florio G, Romeo T, Giordano D, Greco S (2005) Observations on Distribution and Biology of *Galeus Melastomus* (Chondrichthyes, Scyliorhinidae) in the Southern Tyrrhenian Sea (Central Mediterranean). *Cybius* 29: 41-46.
- Mangano MC, Kaiser MJ, Porporato EMD, Spanò N (2013) Evidence of Trawl Disturbance on Mega-Epibenthic Communities in the Southern Tyrrhenian Sea. *Mar Ecol Prog Ser* 475: 101-117.
- Ragonese S, Di Stefano L, Bianchini ML (2000) Catture E Selettività Di Pesci Cartilaginei Nella Pesca Dei Gamberi Rossi Nello Stretto Di Sicilia. *Biol Mar Medit* 7: 400-411.
- Enever R, Catchpole TL, Ellis JR, Grant A (2009) The Survival of Skates (*Rajidae*) Caught by Demersal Trawlers Fishing in UK Waters. *Fish Res* 97: 72-76.