

*This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.*



**ISSN: 2329-9088**

## **Tropical Medicine & Surgery**

---

**The International Open Access  
Tropical Medicine & Surgery**

### **Executive Editors**

**Jean-Paul Joseph Gonzalez**

French Research Institute for Development, USA

**Jimmy Thomas**

East Carolina Heart Institute, North Carolina, USA

**Mark Wiser**

Tulane University of Public Health and Tropical Medicine  
USA

**Shi-Bin Cheng**

Brown University, USA

**Dong Zhou**

University of Pittsburgh, USA

---

**Available online at:** OMICS Publishing Group ([www.omicsonline.org](http://www.omicsonline.org))

**T**his article was originally published in a journal by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2329-9088.1000e105>



## Emerging Infectious Diseases: Assessing the Risk of Zoonotic Exposures

Jimmy T Efird\*

Department of Public Health and the Center for Health Disparities, Brody School of Medicine, East Carolina University, Greenville, NC, USA

While nearly 75% of viruses and 50% of bacteria associated with human disease are zoonotic, the emerging disease risk directly attributable to animal exposures remains unclear [1]. Animals and humans have coexisted throughout their existence and adaptive immunity against microbes appears to be the norm. However, a disproportionate burden of infectious diseases occurs in tropical and developing countries, where contact with animals occurs regularly. Furthermore, the transmission route and etiology for over 14% of human pathogens remains unknown, raising the question of species jumping as a potential new source of disease (i.e., evading innate immune surveillance conferred through evolution) [1].

Certainly, zoonotic infection plays a clear and defined role in many illnesses. Occupations such as farming and agricultural work that place individuals in direct contact with animals and their excretions/secretions increase the risk for infection [2]. However, the ease by which viruses and bacteria readily cross species as newly defined emerging infections among humans' likely remains low. Among chicken farm laborers regularly in contact with infected chickens and wildfowl, only 2 in a sample of 27 persons tested positive for antibodies to Rous Sarcoma Virus-Bryan (RSV-B) [3]. Similarly, wildlife workers who handle wild ducks when they are shedding high levels of virus consistently have negative viral and serologic assays [4]. Under experimental conditions,

participants inoculated with high doses of avian subtypes H4N8, H6N1, and H10N7 failed to produce a detectable antibody response, even though ~25% shed virus and displayed mild clinical signs and symptoms [4].

The attributable risk of most lethal microbial agents at the population level has been contained to small groups, who typically die before transmitting disease in large numbers.

Nonetheless, the possibility of a global pandemic cannot be ruled out and routine public health efforts represent the best prophylactic course of action.

### References

1. Taylor LH, Latham SM, Woolhouse ME (2001) Risk factors for human disease emergence. *Philos Trans R Soc Lond B Biol Sci* 356: 983-989.
2. Williams E, Barker I (2001) *Infectious diseases of wild mammals*. Iowa State University Press, Ames, Iowa, USA.
3. Morgan HR (1967) Antibodies for Rous sarcoma virus (Bryan) in fowl, animal, and human populations of East Africa. II. Antibodies in domestic chickens, wildfowl, primates, and man in Kenya, and antibodies for Burkitt lymphoma cells in man. *Journal of the National Cancer Institute* 39: 1229-1234.
4. Beare AS, Webster RG (1991) Replication of avian influenza viruses in humans. *Arch Virol* 119: 37-42.

---

**\*Corresponding author:** Jimmy T Efird, Department of Public Health and the Center for Health Disparities, Brody School of Medicine, East Carolina University, Greenville, NC, USA, E-mail: [efirdj@ecu.edu](mailto:efirdj@ecu.edu)

**Received** September 25, 2013; **Accepted** September 28, 2013; **Published** October 01, 2013

**Citation:** Efird JT (2013) Emerging Infectious Diseases: Assessing the Risk of Zoonotic Exposures. *Trop Med Surg* 1: e105. doi: [10.4172/2329-9088.1000e105](http://dx.doi.org/10.4172/2329-9088.1000e105)

**Copyright:** © 2013 Efird JT. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.