

Evidence from Cochrane Systematic Reviews in Anesthesiology is Insufficient to Support or Reject the Studied Interventions and Highlights the Need of Further Research

Reinaldo S Santos Jr¹, Regina El Dib^{2,3}, Antonio J B Pereira³, Rodrigo L Alves³, Vitor C S da Silva⁴ and Paulo do Nascimento Jr^{5*}

¹Universidade Estadual Paulista, Faculdade de Medicina de Botucatu, Sao Paulo, Brazil

²UNESP, Universidade Estadual Paulista, Faculdade de Medicina de Botucatu, Sao Paulo, Brazil

³McMaster Institute of Urology, McMaster University, St. Joseph's Healthcare, Hamilton, Ontario, Canada

⁴Hospital São Rafael, Salvador, Brazil

⁵Santa Casa de Misericórdia da Bahia, Hospital Santa Izabel, Salvador, Brazil

Abstract

Objective: The present study aimed to classify the evidence of the systematic reviews from the Cochrane Anaesthesia, Critical and Emergency Care Group as sufficient or not to support or reject the interventions studied and their recommendation or not for further research.

Methods: All systematic reviews published by the Cochrane Anaesthesia, Critical and Emergency Care Group up to February 28, 2014, were analyzed regarding the implications of the interventions studied by each review for clinical practice and research, according to their authors' conclusions. The analyzed values were the percentages and 95% confidence intervals (CI) and descriptive statistics of the included studies and meta-analyses of the reviews is also shown.

Results: One hundred fifteen systematic reviews were analyzed and the results were as follows: evidence likely supporting the interventions with authors' recommendations for further research: 32.2% (95% CI 23.7-40.7); evidence supporting the interventions without authors' recommendations for further research: 2.6% (95% CI 0.0-5.5); evidence likely against the interventions with authors' recommendations for further research: 6.1% (95% CI 1.7-10.5); evidence against the interventions without authors' recommendations for further research: 1.7% (95% CI 0.0-4.0); and insufficient evidence with or without authors' recommendations for further research: 57.4% (95% CI 48.4-66.4) and 0%, respectively. Independent of the results, 95.7% (95% CI 92.0-99.4) of the reviews suggested further research. The numbers of included studies and meta-analyses [median; mode (minimum; 1st quartile; 3rd quartile; maximum)] in the reviews were, respectively, 9.5; 4 (0; 4; 18; 737) and 6; 0 (0; 0; 12; 92).

Conclusion: The majority of the systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group resulted in no evidence or insufficient evidence to strongly recommend or discourage specific interventions for clinical practice and their authors did recommend further randomized controlled trials to provide clear evidence in future updates of systematic reviews.

Keywords: Evidence based medicine; Anesthesiology; Systematic review; Meta-analyses; Scientific research

Introduction

Systematic reviews are one of the best methods for mapping, gathering and producing scientific evidence, because they assess several data from different studies simultaneously, based on reproducible and consistent methodology. According to the number and type of included studies in systematic reviews, meta-analyses can be plotted, representing the combined statistical data from the trials that have met the inclusion criteria.

The Cochrane Collaboration has the largest database of systematic reviews and it aims to produce and report systematic reviews with the highest standard of quality [1]. In addition to its methodological accuracy [2], the inferences from its conclusions are translated into implications for clinical practice and research. There are 53 review groups (Cochrane Review Groups) in several medical fields, thus constituting the database. Established in 2000, the Cochrane Anaesthesia Review Group, currently named Cochrane Anaesthesia, Critical and Emergency (ACE) Care Group, publishes systematic reviews in the areas of anesthesiology, critical care, perioperative medicine and emergency medicine, and it has the largest number of the reviews in these areas of all the existing databases and journals [3].

In addition to these several advantages, systematic reviews are often criticized due to the production of data with no statistical significance, the inconsistency of evidence and the absence of solid recommendations for interventions in clinical practice. Only a small proportion of these systematic reviews have been capable of solidly recommending or discouraging specific interventions with no need for further studies, according to their authors' conclusions [4]. Thus, there is an urgent need to perform primary studies, i.e., randomized controlled trials, in larger numbers and with better quality [5]. In anesthesiology,

***Corresponding author:** Paulo do Nascimento Jr, Department of Anesthesiology, Botucatu Medical School, UNESP, Distrito de Rubião Jr s/n, CEP 18.618-970m, Botucatu/SP, Brazil, Tel: 5571991165794; E-mail: pnasc@fmb.unesp.br

Received October 06, 2016; **Accepted** November 23, 2016; **Published** November 30, 2016

Citation: Santos RS Jr, El Dib R, Pereira AJB, Alves RL, da Silva VCS, et al. (2016) Evidence from Cochrane Systematic Reviews in Anesthesiology is Insufficient to Support or Reject the Studied Interventions and Highlights the Need of Further Research. Evid Based Med Pract 2: 111. doi: [10.4172/2471-9919.1000111](https://doi.org/10.4172/2471-9919.1000111)

Copyright: © 2016 Santos RS Jr, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

there is also the necessity of comprehending the results of systematic reviews so that other primary studies and systematic reviews can be performed in different subfields of knowledge. The absence of such comprehensiveness is responsible for the excess of studies in some fields and the lack of studies in others. Moreover, many issues have not been adequately explored by randomized controlled trials, which very often have high levels of heterogeneity, making it difficult to plot them in meta-analyses and to obtain clearer and more precise conclusions.

Previous publications that mapped all the groups of Cochrane database showed that their systematic reviews were invariably inconclusive, concerning the applicability of their results in clinical practice [4,5]. For this reason, a question arises regarding the uncertainties of systematic reviews in anesthesiology, relative to the applicability of their results in clinical practice and research.

In the present study, we proposed the mapping of systematic reviews published in the Cochrane Anaesthesia, Critical and Emergency Care Group and, according to their author's conclusions, classify the evidence as sufficient or not to support or reject the interventions studied. The recommendation or not for further research, based on each author's conclusion and the general characteristics of the reviews relative to the included studies and meta-analyses were also described. So far, this is the first study made under such criteria in the anesthesia field.

Materials and Methods

All systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group published until February 28, 2014 were included. All of the information was extracted from the reviews according to their authors' conclusions. The information concerned the existence or not of benefits from a given studied intervention, as well as the need for further research, according to the evidence to recommend or not the intervention for clinical practice.

This methodology was similar to that of previously published studies that mapped and qualified the evidence from systematic reviews [4,5]. Only complete systematic reviews of randomized controlled trials were included. Protocols and systematic reviews that did not directly involve randomized controlled trials were excluded. Systematic reviews that recommended or did not recommend further research based on cost-benefit analyses were not considered, unless their primary endpoint was the cost of a treatment or intervention.

We used an excel spreadsheet where we indicated the name of the study and one of the three possible categories: "evidence supporting the interventions"; "evidence against the interventions" and "insufficient evidence to support or reject the interventions". The studies were then classified in two subgroups in each category: "recommendation for further studies" and "no recommendation for further studies".

Two investigators working together (RED and RSSJ) identified and extracted all the complete systematic reviews of randomized controlled trials to be included in the study from the Cochrane Anaesthesia, Critical and Emergency Care Group via Cochrane Library database and printed two copies of the full text for further independent analysis. Two investigators independently analyzed and classified each review (RSSJ, PNJ). In cases of disagreement or doubt, a third investigator (RED) would be consulted for consensus.

The classification of the systematic reviews is based on the authors' conclusions according to their overall analysis of the results, regardless of what they considered, i.e., one primary outcome or several secondary outcomes. Then, the systematic reviews were allocated into six possible categories.

When the authors concluded that the evidence supported the intervention and recommended further studies, we defined the category #1, using the word "likely":

- (1) Evidence likely supporting the interventions and the authors recommending further research.

When the authors concluded that the evidence supported the intervention with no recommendation for further research, we defined the category #2:

- (2) Evidence supporting the interventions and the authors not recommending further research.

When the authors concluded that the evidence favored control group and recommended further research, we defined category #3 using the word "likely":

- (3) Evidence likely against the interventions and the authors recommending further research.

When the evidence favored control group and the authors did not recommend further research, we defined category #4:

- (4) Evidence against the interventions and the authors not recommending further research.

Insufficient evidence to draw a conclusion and the authors recommending further research was category #5:

- (5) Insufficient evidence to support or to reject the interventions and the authors recommending further research b And when there was insufficient evidence to draw a conclusion and the authors not recommending further research, we defined category #6:

- (6) Insufficient evidence to support or to reject the interventions and the authors not recommending further research.

Evidence supporting the intervention was defined as effectiveness (the extent to which a specific intervention, when used under ordinary circumstances, does what it is intended to do) or efficacy (the extent to which an intervention produces a beneficial result under ideal conditions) as compared to the control group, whether placebo, standard care or another intervention [6]. On the other hand, evidence against the intervention was defined when effectiveness or efficacy was seen on the control group instead of the intervention group.

The numbers of clinical trials and meta-analyses in each systematic review were also analyzed. Meta-analyses plotted with just one study were not computed.

Statistical methodology


The implications for clinical practice and research are presented as absolute numbers, percentages and 95% confidence intervals (CI) of all of the evaluated systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group. Meta-analyses and the included studies in each review are expressed as medians, minimum numbers, first and third quartiles, maximum numbers and modes.

Results

One hundred eighteen systematic reviews were identified. Two reviews were excluded from the study due to duplication and one due to being a review summary. After the exclusions, 115 reviews were analyzed (Figure 1). The median number of randomized controlled trials per systematic review was 9.5, for a total of 2258 studies in all of

the reviews. The median number of meta-analyses per review was six, for a total of 1072 meta-analyses in all of the reviews (Table 1).

There was 100% inter-rate agreement between investigators on the classification of the systematic reviews. The commonest event observed in this study, occurring in 57.4% of the reviews, was insufficient evidence to support or reject the interventions with the authors recommending more studies. From the total, in only five reviews (4.3%), the authors did not recommend more studies, considering the actual evidence sufficient to recommend or discourage the intervention. Therefore, 95.7% of the reviews recommended the implementation of additional research, independent of the results obtained from a given intervention, i.e., benefit, harm or absence of evidence to support or discourage the



intervention. Data on implications for clinical practice and research are expressed in Table 2.

Discussion

This study demonstrated a lack of solid evidence produced by the majority of the systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group. According to their authors, 110 out of 115 reviews were not considered sufficiently powerful to answer questions regarding clinical practice convincingly.

Evidence-based medicine represents the bond between good clinical research and clinical practice. Therefore, it should offer doctors and health policy-makers the best available evidence, which should be adequate, consistent and straightforward [7]. High-quality methodological research, with internal and external validation, is an essential condition for achieving these goals.

At the top level of the evidence hierarchy are systematic reviews, which are secondary studies built from primary studies. They are considered the best method for documenting, mapping and producing scientific evidence [2], while they play a major role in supporting fields in need of more research. The best primary evidence consists of randomized controlled trials, but depending on the condition to be studied, cohort studies could represent the best evidence. A systematic review of randomized controlled trials can conclude whether a tested intervention is effective, ineffective or harmful or whether there is insufficient evidence for any conclusion. Unfortunately, the inconsistency of results and absence of sufficient evidence to answer a clinical question are the most frequent outcomes found, which has been a subject of criticism and controversy [3].

The mapping of systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group resulted, in general, in absence of evidence to recommend or not a given intervention. Furthermore, their authors suggested the production of more research, similar to what was recently observed when all of the databases were mapped [5]. The same situation was shown in 2007 [4] and there have not been changes so far, demonstrating the need to focus on primary studies with better quality. The recommendation to conduct further randomized controlled trials occurred in 96.9% [4] and 95.4% [5] of the cases, respectively.

The lack of evidence is a consequence of several factors. In general,

Statistical data	Studies	Meta-analyses
	Included in the reviews	
Median (minimum; 1 st quartile; 3 rd quartile; maximum)	9.5 (0; 4; 18; 737)	6 (0; 0; 12; 92)
Mode	4	0
Total (n, considering all systematic reviews)	2258	1072

Table 1: Statistical data of the systematic reviews.

Implications for clinical practice and research	N	Percentage (%)	95% Confidence interval (%)
Evidence supporting the interventions	40	34.8	26.0-43.5
Evidence likely supporting the interventions and the authors recommending further research	37	32.2	23.7-40.7
Evidence supporting the interventions and the authors not recommending further research	3	2.6	0.0-5.5
Evidence against the interventions	9	7.8	2.9-12.7
Evidence likely against the interventions and the authors recommending further research	7	6.1	1.7-10.4
Evidence against the interventions and the authors not recommending further research	2	1.7	0.0-4.0
Insufficient evidence to support or reject the interventions	66	57.4	48.4-66.4
And the authors recommended further research	66	57.4	48.4-66.4
And the authors did not recommend further research	0	0	-
Total number of systematic reviews with recommendations for further research	110	95.7	92.0-99.4

Table 2: Outcomes of the systematic reviews related to clinical practice and research.

it is due to the poor methodological quality of the primary studies, specifically no clear randomization [8], single-center research with small sample sizes, no concealment of patients' distribution into groups, lack of blinding of results for outcome evaluators [9], bias due to pharmaceutical company financing [10] and different types of heterogeneity.

Publication bias can also erroneously affect the clinical care of patients and future research. De Oliveira et al. [11] analyzed 1163 papers in anesthesiology, demonstrating that a positive result, or a result favorable to an intervention, was an independent predictive factor for publication in higher-impact journals. Moreover, a study with negative or unfavorable results took considerably more time to be accepted for publication [12]. In the present study, there were a higher number of systematic reviews with positive or favorable results in comparison to the negative outcomes, similarly to the fact previously described. A recently published meta-epidemiological analysis [13] that included 93 meta-analyses from 735 randomized controlled trials, published in journals from different specialties, showed significant variation in the sample sizes of the studies in their meta-analyses, with the smaller studies having a higher probability of overestimation of the positive effects of the tested treatments. It is known that research with smaller sample sizes is more susceptible to publication bias [14], whereas results from clinical studies with larger sample sizes tend to be more representative, valuable and publishable.

In our investigation, although it was not an objective, we observed important methodological heterogeneity among the various studies included in the reviews, creating more biases and placing the quality and value of the results at risk. Moreover, the majority of the systematic reviews had no meta-analyses (mode=0) and there was great diversity in the numbers of included studies (0 to 737). One of the most important findings was the lack of randomized controlled trials in systematic reviews (mode=4), which curiously were supposed to be systematic reviews from randomized controlled trials. Due to the heterogeneity of their primary studies, many systematic reviews do not present statistical data or meta-analyses. In the present investigation we observed a high number of systematic reviews with no meta-analyses (mode=0). This fact is probably related to both, the absence or the low number of randomized controlled trials included in the systematic reviews. Heterogeneity of included studies may have contributed to this observation.

One of the major principles of evidence-based medicine is the appraisal and classification of evidence. However, such criteria have been questioned by some investigators, who have suggested remodeling the concepts of power and quality in studies, as well as the resultant evidence [15]. They have suggested that additional dimensions not traditionally considered for classifying evidence should be incorporated, assigning weight to each dimension when determining the overall quality of the evidence. From the results we obtained, we believe that such changes have become necessary, not only in anesthesiology but also in all fields.

There were some limitations of the present investigation. The obtained results were based on the conclusions of the authors of each systematic review. We did not interpret the reviews, nor did we question the authors' interpretations. Possible biases were not analyzed. It is possible that scrutiny of each review would have raised different conclusions from those presented by their authors. Even so, we believe that little would have changed in the results, since most of systematic reviews showed no evidence to support or to reject the intervention. We did not include other databases either, which might have contained different information. Nonetheless, the Cochrane Collaboration has an

exclusive profile, gathering most of the published systematic reviews and following a pattern for their conclusions and outlining the efficacy and/or effectiveness of a treatment, according to the primary outcome and the control group. It also suggests the need, or not, for more studies to support or refute a given intervention, thereby enabling responses to clinical questions.

We conclude that a significant number of the systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group lack solid and consistent evidence for decision-making in clinical practice. On the other hand, around 50% of the systematic reviews offer suggestions that could influence clinical practice and this number may be considered reasonable. Nonetheless, the uncertainties observed in the reviews' conclusions and the lack of included studies and meta-analyses reaffirm the need for more high-quality randomized controlled trials, which could improve the conclusiveness of the systematic reviews. Knowledge translations studies are necessary to verify the real impact of these systematic reviews on clinical practice.

Conclusion

The majority of the systematic reviews of the Cochrane Anaesthesia, Critical and Emergency Care Group resulted in no evidence or insufficient evidence to strongly recommend or discourage specific interventions for clinical practice and their authors did recommend further randomized controlled trials to provide clear evidence in future updates of systematic reviews.

Competing Interests

Non-financial competing interests.

Regarding conflict of interests, we inform that one of the authors (RED) is a member of the Cochrane Diagnostic Test Accuracy Working Group and the Cochrane Prognosis Methods Group and the manuscript reflects our opinion, not Cochrane Collaboration's opinion.

References

1. Cochrane AL (1979) A critical review, with particular reference to the medical profession. *Medicines for the year 2000*. London: Office of health economics, pp: 1-11.
2. Petticrew M (2003) Why certain systematic reviews reach uncertain conclusions. *BMJ* 326: 756-758.
3. Moller AM (2012) How to map the evidence: The development of the systematic review in anaesthesia. *Br J Anaesth* 109: 32-34.
4. El Dib RP, Atallah AN, Andriolo RB (2007) Mapping the cochrane evidence for decision making in health care. *J Eval Clin Pract* 13: 689-692.
5. Villas Boas PJ, Spagnuolo RS, Kamegasawa A, Braz LG, Polachini do Valle A, et al. (2013) Systematic reviews showed insufficient evidence for clinical practice in 2004: what about in 2011? The next appeal for the evidence-based medicine age. *Journal of Evaluation in Clinical Practice* 19: 633-637.
6. Higgins JPT, Green S (2011) *Cochrane Handbook for Systematic Reviews of Interventions* Version. The Cochrane Collaboration.
7. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS (1996) Evidence based medicine: What it is and what it is not. *BMJ* 312: 71-72.
8. Kjaergard LL, Villumsen J, Gluud C (2001) Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Ann Intern Med* 135: 982-989.
9. Savovic J, Jones HE, Altman DG, Harris RJ, Juni P, et al. (2012) Influence of reported study design characteristics on intervention effect estimates from randomized, controlled trials. *Ann Intern Med* 157: 429-438.
10. Lexchin J, Bero LA, Djulbegovic B, Clark O (2003) Pharmaceutical industry sponsorship and research outcome and quality: Systematic review. *BMJ* 326: 1167-1170.
11. De Oliveira GS Jr, Chang R, Kendall MC, Fitzgerald PC, McCarthy RJ (2012) Publication bias in the anesthesiology literature. *Anesth Analg* 114: 1042-1048.

12. Stern JM, Simes RJ (1997) Publication bias: Evidence of delayed publication in a cohort study of clinical research projects. *BMJ* 315: 640-645.
13. Dechartres A, Trinquart L, Boutron I, Ravaud P (2013) Influence of trial sample size on treatment effect estimates: Meta-epidemiological study. *BMJ* 346: f2304.
14. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, et al. (2011) Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 343: d4002.
15. Bagshaw SM, Bellomo R (2008) The need to reform our assessment of evidence from clinical trials: A commentary. *Philos Ethics Humanit Med* 3: 23.

Citation: Santos RS Jr, El Dib R, Pereira AJB, Alves RL, da Silva VCS, et al. (2016) Evidence from Cochrane Systematic Reviews in Anesthesiology is Insufficient to Support or Reject the Studied Interventions and Highlights the Need of Further Research. *Evid Based Med Pract* 2: 111. doi: [10.4172/2471-9919.1000111](https://doi.org/10.4172/2471-9919.1000111)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>