

Exposure to Nicotine in Adult, but not Adolescent, Rats Increases Alcohol Self-Administration in Adulthood

Paula C Bianchi^{1,2,#}, Fabio C Cruz^{3,4,#}, Paulo E Carneiro-de-Oliveira^{1,2}, Tarciso T Miguel⁵, Marcelo T Marin^{1,2}, Ricardo L Nunes-de-Souza^{1,2}, Cleopatra S Planeta^{1,2} and Rodrigo M Leão^{1,2,3*}

¹Laboratory of Pharmacology, São Paulo State University (Unesp), School of Pharmaceutical Sciences, Araraquara, SP, Brazil

²Joint Graduate Program in Physiological Sciences, UFSCar/UNESP, Araraquara, São Paulo, Brazil

³Departamento de Biorregulação, Instituto de Ciências da Saúde, Universidade Federal da Bahia, Salvador, BA 40110-902, Brazil

⁴Pharmacology Department, Universidade Federal de São Paulo, UNIFESP, Brazil

⁵Biomedical Sciences Institute, Universidade Federal de Uberlândia, Uberlândia, Minas Gerais, Brazil

#Authors contributed equally to this work

*Corresponding author: Rodrigo M Leão, Departamento de Biorregulação, Instituto de Ciências da Saúde, Universidade Federal da Bahia, Av. Reitor Miguel Calmon, s/n, Vale do Canela, Salvador, BA, Brazil, 40110-902; Tel: 55-71- 3283 8885; Fax: 55-71- 3283 8894; E-mail: rodrigomolini@yahoo.com.br

Received date: September 01, 2017; Accepted date: September 13, 2017; Published date: September 20, 2017

Copyright: © 2017 Bianchi PC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Among drugs of abuse, nicotine and alcohol are the most consumed in the world, and are often co-abused. Evidence has shown that nicotine enhances the consumption and subjective effects of ethanol. How nicotine increases alcohol consumption is an interesting question that is still left unanswered. We assessed whether repeated treatment with nicotine during adolescence or adulthood could increase operant alcohol self-administration. Male adolescent and adult rats (postnatal day [PND] 28 and 60, respectively) were injected with nicotine (1.0 mg/kg, subcutaneously [s.c.], three times a day for 10 days). During this period, the control group received saline injections (0.1 ml/kg; s.c). On PND 72, adolescent and adult rats pretreated with nicotine were subjected to ethanol self-administration procedures. Alcohol oral self-administration patterns were evaluated using a progressive ratio schedule of reinforcement and 24-hour unlimited access to ethanol. Our results showed that nicotine pretreatment in adult rats increases the motivation for alcohol consumption. However, nicotine exposure during adolescence does not affect alcohol-consuming behaviors in adulthood.

Keywords: Alcohol; Nicotine; Self-administration; Adult and adolescent rats

Introduction

Nicotine and alcohol are the most consumed drugs of abuse in the world and are often co-abused [1]. Epidemiological studies indicate that alcoholism is more prevalent in smokers than in non-smokers [2-5]. Evidence has shown that nicotine can increase alcohol consumption in humans [6-8]. In addition, several pre-clinical studies have shown that repeated administration of nicotine concomitantly with ethanol increases alcohol intake in a home-cage bottle choice paradigm, operant self-administration, and voluntary ethanol seeking [9-15].

Initiation of tobacco smoking, in general, occurs during early adolescence [16,17]. It has been shown that tobacco use during adolescence produces enduring effects that may increase the vulnerability to addiction later in life [18,19]. As reported for adults, alcohol and tobacco are frequently consumed together by adolescents [20,21]. Exposure to nicotine during adolescence increases the propensity for ethanol dependence in adulthood [22]. Recently, it was observed that adolescents exposed to nicotine showed increased binge-like ethanol consumption in adulthood [23]. Moreover, adolescent nicotine exposure facilitates the establishment of ethanol conditioned place preference in adult animals [24]. Evidence demonstrated that the administration of nicotine to mice during adolescence, but not during adulthood, increases sensitivity to ethanol withdrawal in adult mice

[25]. In addition, studies have demonstrated that administration of nicotine or alcohol during adolescence increases the self-administration of the same drug in the adulthood [26-28]. Taken together, these studies suggest that adolescence is a period of increased vulnerability in the development of drug addiction.

Despite the high prevalence of tobacco use and abuse among adolescents, and the fact that adolescent nicotine exposure appears to impact adult ethanol self-administration, the influence of nicotine on ethanol consumption has been studied mainly in adult animals. Here we assessed whether repeated nicotine administration during adolescence or adulthood could increase alcohol-consuming behaviour in adult rats.

Materials and Methods

Subjects

Male Wistar rats were obtained from the animal breeding facility of the São Paulo State University-UNESP at postnatal day (PND) 21 and 60. Groups of four animals were housed in plastic cages 32 (width) X 40 (length) X 16 (height) cm in a room maintained at 23 ± 2°C. Rats were continuously maintained on a reversed light cycle (12 hours light, 12 hours dark, lights off at 08:00 a.m.) and free access to food and water were allowed. All experiments were performed during the dark phase. The experimental protocol was approved by the Ethical

Committee for Use of Human or Animal Subjects of the School of Pharmaceutical Science-UNESP (CEP-12/2008).

Drugs

Nicotine: nicotine 99% (Sigma-Aldrich, MO, USA) was dissolved in saline solution (NaCl 0.9%) at a concentration of 1 mg/ml. Ethanol (Sigma-Aldrich, St Louis, MO, USA), Saccharine (Sigma-Aldrich, St Louis, MO, USA).

Alcohol solutions

Alcohol solutions were prepared by diluting ethanol in tap water at different concentrations (2%, 4%, or 6%; v/v) containing 0.05% of saccharine. The self-administration control group had access only to 0.05% saccharine solution.

Nicotine pretreatment regimen

Adolescents (PND 28) and adults (PND 60) were pretreated with nicotine (1.0 mg/kg subcutaneously) or saline (1.0 mL/kg), three times a day, for 10 days. The dose of nicotine was chosen based on previous experiments conducted in our laboratory.

Operant ethanol self-administration

On PND 72, adolescent and adult rats pretreated with nicotine were subjected to ethanol self-administration procedures. The general procedure was adapted from a previous study [29].

Apparatus

For ethanol self-administration, the animals were put individually in experimental chambers (30 x 30.5 x 24.5 cm) and enclosed in light- and sound-attenuating boxes. The floor of the chambers consisted of a Plexiglas tray covered with sawdust. The front wall of the chamber contained one interchangeable panel. The panel was equipped with two levers located 5 cm above the floor, two cue lights (red and green) above each lever, a hole, 5 cm in diameter, located 1 cm above the floor for alcohol delivery, and a session light in the middle of the panel (12 cm above the floor).

Ethanol self-administration acquisition

Rats were initially given unlimited access to saccharine solution (0.05%) self-administration, in which each lever press was reinforced with the delivery of 0.2 ml solution (fixed ratio schedule of reinforcement; FR1), followed by a 20-second time-out. During self-administration sessions, a green cue light above the active lever signaled saccharine availability and a red cue light, also above the active lever, signaled saccharine delivery. During the time-out period, the green and red cue lights were extinguished and lever presses were recorded but had no consequence. Each daily session was terminated after 60 minutes of access. Three sessions of saccharine reinforcement were performed. The next two sessions of 2% alcohol solution (v/v) containing 0.05% of saccharine introduced a schedule of reinforcement (FR1) for the active lever, followed by two sessions in which we used 4% ethanol solution with saccharine (0.05%). We then provided 6% ethanol solution containing 0.05% of saccharine for the final two acquisition sessions. The control group received saccharine solution (0.05%) in all sessions.

Ethanol self-administration maintenance

After two sessions of 6% ethanol solution, the FR schedule was increased to fixed ratio 3 (FR3). Rats were maintained for five interspersed sessions, three hours per day with short unlimited access, on an FR3 schedule before assessing the progressive ratio schedule of reinforcement.

Progressive ratio schedule of drug reinforcement

After the acquisition and maintenance phase, the self-administration progressive ratio (PR) schedule of drug reinforcement was verified. The progression of response requirements followed the algorithm in accordance to [30]: 1, 1, 2, 2, 3, 3, 4, 4, 5, 5, 7, 7, 9, 9, 11, 11, 13, 13, 15, 15, 18, 18, 21, 21, 24, 24, 27, 27, 30, 30, 34, 34, 38, 38, 42, 42, 46, 46, 50, 50... The final infusion delivered was defined as the break point. The PR session was terminated once the rat failed to obtain reinforcement during 60 minutes. The average number of total responses and the last ratio achieved across the three interspersed PR trials for each individual rat (PND 91-97) was calculated.

24-hours long access

One day after the final PR session, each rat (PND 99) was submitted to a 24-hour unlimited access protocol, with unlimited alcohol access (0.2 mL per reinforcement of 6% ethanol solution containing 0.05% of saccharine on an FR1 schedule). The amounts of alcohol self-administered, as well as the pattern of responding, were recorded.

Statistical analysis

Data were analyzed by two-way ANOVA considering pretreatment (saline vs. nicotine) and solution (saccharine vs. ethanol). When a significant ($p < 0.05$) main effect was observed, the Newman-Keuls test was applied for individual post-hoc comparisons.

Results

Progressive ratio schedule of drug reinforcement


Figure 1A-C depicts the number of responses, reinforcements, and the final ratio achieved (break point) in a progressive ratio schedule to saccharine or ethanol in adult animals pretreated with nicotine or saline, during adolescence.

With respect to the number of responses, two-way ANOVA did not show a significant difference between groups [$F(3,56)=2.86$, $p > 0.05$].

Two-way ANOVA for reinforcement indicated a significant interaction between factors [$F(3,62)=4.04$, $p < 0.05$]. Newman-Keuls *post-hoc* comparison showed that pre-treatment with nicotine during adolescence increased the number of reinforcements for saccharine, but not for ethanol.

A similar result was observed with respect to the final ratio achieved: two-way ANOVA found significant interaction between pre-treatment and solution [$F(3,61)=5.62$, $p < 0.05$]. Post-hoc analysis indicated that nicotine pre-treatment increased the final ratio achieved for saccharine, but not for ethanol solution.

Figure 2A-C depicts the number of responses, reinforcements, and the final ratio achieved (break point) in a progressive ratio schedule to saccharine or ethanol in adult animals pretreated with nicotine or saline during adulthood.




Two-way ANOVA of responses indicated a significant interaction between the two factors [$F(3,38)=4.90$; $p < 0.05$]. Post-hoc analysis revealed that pre-treatment with nicotine during adulthood increased the number of responses for ethanol, but not saccharine, during operant self-administration.

Two-way ANOVA revealed a main effect of solution on the number of reinforcements [$F(1,47)=15.3$; $p < 0.05$]. In addition, ANOVA also showed a significant interaction between factors [$F(3,47)=11.0$; $p < 0.05$]. Post-hoc analysis indicated that the pre-treatment with nicotine during adulthood increased the number of responses for ethanol, but not saccharine, during operant self-administration.

For the final ratio achieved, two-way ANOVA indicated a significant interaction between the two factors [$F(3,47)=15.2$; $p < 0.05$]. Post-hoc analysis indicated that the pre-treatment with nicotine during the adulthood increased the number of responses for ethanol, but not saccharine, during operant self-administration.

24-Hour long unlimited access

Figure 3A-C depicts the number of responses, reinforcement, and the total ethanol or saccharine intake during the 24-hour long unlimited access session in adult animals pre-treated with nicotine or saline during adolescence.



Two-way ANOVA on the number of responses [$F(1,53)=9.27$, $p < 0.05$] and reinforcements [$F(1,53)=15.77$, $p < 0.05$] indicated a significant main effect of solution.

Figure 4A-C depicts the number of responses, reinforcements, and the total ethanol or saccharine intake during the 24-hour long unlimited access session in adult animals pre-treated with nicotine or saline during adulthood. Two-way ANOVA on the number of responses and reinforcements indicated a significant main effect of solution [$F(1,47)=29.39$; $F(1,47)=16.51$, $p < 0.05$, respectively].

Discussion

We studied the effects of nicotine pre-treatment during adolescence (PND 28-37) or adulthood (PND 60-69) on a progressive ratio

schedule and binge-like consumption in an operant ethanol self-administration protocol, carried out on PND 75-79 and PND 81-84, respectively. Pre-treatment with nicotine in adult, but not adolescent, rats increased the motivational aspects of alcohol reinforcement in late adulthood as shown by the breaking point results. However, nicotine pre-treatment did not affect the consummatory aspects when animals were allowed to freely intake ethanol for 24 hours.

Although several epidemiological studies have demonstrated that nicotine exposure during adolescence increases alcohol consumption in adulthood [18,19], we found that in adolescent rats, nicotine pre-treatment did not have long lasting effects on alcohol seeking behaviours.





Figure 3: Figure depicts the number of responses, reinforcements, and total ethanol or saccharine intake during a 24-hour long access session in animals pre-treated with nicotine or saline during adolescence (N=15-20). Sacch = saccharine solution. EtOH = ethanol solution.

Figure 4: Figure depicts the number of responses, reinforcements, and total ethanol or saccharine intake during a 24-hour long access session in animals pretreated with nicotine or saline during adulthood (N=9-12). Sacch = saccharine solution. EtOH = ethanol solution.

Nicotine pre-treatment has previously shown to increase alcohol seeking in adult rodents [13,31-34]. In accordance with our results, López-Moreno et al. [35] showed an increase in ethanol self-administration 14 days after cessation of chronic nicotine administration in adult rats. Moreover, Le and colleagues (2003), showed that nicotine-induced enhancement in ethanol operant self-administration depended on the dose and time after the last nicotine injection. However, nicotine pre-treatment has also been associated with decreased operant self-administration of ethanol [36-38]. The reasons for these different results may be due to the nicotine withdrawal period, schedule of administration, or dose of nicotine.

In accordance with our finding, Kempainen and colleagues (2009), using a two-bottle choice paradigm, showed that nicotine pre-treatment in adolescent rats (PND 29-45) did not change ethanol intake when the animals were tested in adulthood (PND 93-121), but ethanol intake was increased when they were re-injected with nicotine in adulthood prior to ethanol access. In the same way, Smith et al. [39] showed that chronic continuous exposure to nicotine during adolescence (PND 35-53) did not modify ethanol intake in adulthood (PND 54-75). In contrast, [23] observed that female adolescent mice (PND 35-41) exposed to nicotine showed increased bottle-drinking ethanol consumption on (PND 41-44). The effects of nicotine exposure during adolescence on ethanol intake seem to depend on the treatment age, duration, strain, sex, and nicotine withdrawal period [23,40].

To our knowledge, this is the first study that has demonstrated increased operant saccharine self-administration while assessing the consequences of nicotine pretreatment during adolescence in later life. Since changes were observed in the motivation for ethanol seeking in adult rats pretreated with nicotine during adolescence, we can speculate that nicotine causes enduring plastic changes in specific pathways associated with natural reinforcement.

Our finding is in agreement with several adult human and non-human studies that showed an increase in consumption of sweet-tasting and high-calorie foods during nicotine withdrawal [41-45]. However, we did not observe any increase in saccharine intake in adult pre-treated rats. Methodological differences between our result and those cited in the literature may justify the different outcomes. For instance, while in our study the animals were tested after a long nicotine withdrawal period, others tested after a short abstinence period.

It is worth pointing out, that although epidemiological studies have demonstrated that nicotine may act as a "gateway" to subsequent alcohol use and abuse later in life [1,3], it is not clear whether this phenomenon is a consequence of the drug's effects per se or the involvement of related social aspects [46]. Thus, more studies should be conducted to answer this important question.

In summary, we have contributed more evidence that nicotine pretreatment increases the motivation to consume alcohol in adult rats. However, nicotine exposure during adolescence does not affect alcohol-taking behaviours in later adulthood.

Acknowledgements

The authors appreciate the excellent technical assistance of Elisabete Z. P. Lepera and Rosana F. P. Silva. This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) fellowships: 2011/14780-2 to PCB, 2008/10691-2 to RML and 2007/08087-7; 2013/24986-2 to FCC, and grant 2011/07469-9 to CSP. CSP is a CNPq fellowship recipient.

References

- Grant BF, Hasin DS, Chou SP, Stinson FS, Dawson DA (2004) Nicotine dependence and psychiatric disorders in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 61: 1107-1115.
- Burling TA, Ziff DC (1988) Tobacco smoking: a comparison between alcohol and drug abuse inpatients. *Addict Behav* 13: 185-190.
- DiFranza JR, Guerrera MP (1990) Alcoholism and smoking. *J Stud Alcohol* 51: 130-135.
- Harrison EL, Desai RA, McKee SA (2008) Nondaily smoking and alcohol use, hazardous drinking, and alcohol diagnoses among young adults: findings from the NESARC. *Alcohol Clin Exp Res* 32: 2081-2087.
- Harrison EL, McKee SA (2008a) Young adult non-daily smokers: patterns of alcohol and cigarette use. *Addict Behav* 33: 668-674.
- Barrett SP, Darredeau C, Pihl RO (2006) Patterns of simultaneous polysubstance use in drug using university students. *Hum Psychopharmacol* 21: 255-263.
- Kouri EM, McCarthy EM, Faust AH, Lukas SE (2004) Pretreatment with transdermal nicotine enhances some of ethanol's acute effects in men. *Drug Alcohol Depend* 75: 55-65.
- Harrison EL, McKee SA (2008b) Young adult non-daily smokers: patterns of alcohol and cigarette use. *Addict Behav* 33: 668-674.
- Alen F, Santos A, Moreno-Sanz G, Gonzalez-Cuevas G, Gine E, et al. (2009) Cannabinoid-induced increase in relapse-like drinking is prevented by the blockade of the glycine-binding site of N-methyl-D-aspartate receptors. *Neuroscience* 158: 465-473.
- Blomqvist O, Ericson M, Johnson DH, Engel JA, Soderpalm B (1996) Voluntary ethanol intake in the rat: effects of nicotinic acetylcholine receptor blockade or subchronic nicotine treatment. *Eur J Pharmacol* 314: 257-267.
- Hausser SR, Getachew B, Oster SM, Dhaher R, Ding ZM, et al. (2012) Nicotine modulates alcohol-seeking and relapse by alcohol-preferring (P) rats in a time-dependent manner. *Alcohol Clin Exp Res* 36: 43-54.
- Le AD, Lo S, Harding S, Juzytsch W, Marinelli PW, et al. (2010) Coadministration of intravenous nicotine and oral alcohol in rats. *Psychopharmacology (Berl)* 208: 475-486.
- Le AD, Wang A, Harding S, Juzytsch W, Shaham Y (2003) Nicotine increases alcohol self-administration and reinstates alcohol seeking in rats. *Psychopharmacology (Berl)* 168: 216-221.
- Leao RM, Cruz FC, Vendruscolo LF, de Guglielmo G, Logrip ML, et al. (2015) Chronic nicotine activates stress/reward-related brain regions and facilitates the transition to compulsive alcohol drinking. *J Neurosci* 35: 6241-6253.
- Olausson P, Ericson M, Lof E, Engel JA, Soderpalm B (2001) Nicotine-induced behavioral disinhibition and ethanol preference correlate after repeated nicotine treatment. *Eur J Pharmacol* 417: 117-123.
- Chen K, Kandel DB (1995) The natural history of drug use from adolescence to the mid-thirties in a general population sample. *Ame J Public Health* 85: 41-47.
- Lantz PM (2003) Smoking on the rise among young adults: implications for research and policy. *Tobacco control* 12 Suppl 1: i60-70.
- Breslau N, Peterson EL (1996) Smoking cessation in young adults: age at initiation of cigarette smoking and other suspected influences. *Am J Public Health* 86: 214-220.
- Chen J, Millar WJ (1998) Age of smoking initiation: implications for quitting. *Health reports* 9: 39-46(Eng); 39-48(Fre).
- Best D, Rawaf S, Rowley J, Floyd K, Manning V, et al. (2000) Drinking and smoking as concurrent predictors of illicit drug use and positive drug attitudes in adolescents. *Drug Alcohol Depend* 60: 319-321.
- Lewinsohn PM, Rohde P, Brown RA (1999) Level of current and past adolescent cigarette smoking as predictors of future substance use disorders in young adulthood. *Addiction* 94: 913-921.
- Spear LP (2015) Adolescent alcohol exposure: Are there separable vulnerable periods within adolescence? *Physiol Behav* 148:122-130.
- Locker AR, Marks MJ, Kamens HM, Klein LC (2016) Exposure to nicotine increases nicotinic acetylcholine receptor density in the reward pathway and binge ethanol consumption in C57BL/6J adolescent female mice. *Brain Res Bull* 123: 13-22.
- Philpot RM, Engberg ME, Wecker L (2014) Ethanol conditioned place preference and alterations in DeltaFosB following adolescent nicotine administration differ in rats exhibiting high or low behavioral reactivity to a novel environment. *Behavioural brain research* 262: 101-108.
- Riley HH, Zalud AW, Diaz-Granados JL (2010) The influence of a chronic adolescent nicotine exposure on ethanol withdrawal severity during adulthood in C3H mice. *Alcohol* 44: 81-87.
- Adriani W, Spijker S, Deroche-Gamonet V, Laviola G, Le Moal M, et al. (2003) Evidence for enhanced neurobehavioral vulnerability to nicotine during periadolescence in rats. *The Journal of neuroscience: the official journal of the Society for Neuroscience* 23: 4712-4716.
- Laviola G, Macri S, Morley-Fletcher S, Adriani W (2003) Risk-taking behavior in adolescent mice: psychobiological determinants and early epigenetic influence. *Neuroscience and biobehavioral reviews* 27: 19-31.
- Alaux-Cantin S, Warnault V, Legastelois R, Botia B, Pierrefiche O, et al. (2013) Alcohol intoxications during adolescence increase motivation for alcohol in adult rats and induce neuroadaptations in the nucleus accumbens. *Neurop* 67: 521-531.
- Cruz FC, Quadros IM, Planeta Cda S, Miczek KA (2008) Maternal separation stress in male mice: long-term increases in alcohol intake. *Psychopharmacology (Berl)* 201: 459-468.

30. Gilpin NW, Smith AD, Cole M, Weiss F, Koob GF, et al. (2009) Operant behavior and alcohol levels in blood and brain of alcohol-dependent rats. *Alcohol Clin Exp Res* 33: 2113-2123.
31. Abreu-Villaca Y, Cavina CC, Ribeiro-Carvalho A, Correa-Santos M, Naiff VF, et al. (2013) Combined exposure to tobacco smoke and ethanol during adolescence leads to short- and long-term modulation of anxiety-like behavior. *Drug Alcohol Depend* 133: 52-60.
32. Britt JP, Bonci A (2013) Alcohol and tobacco: how smoking may promote excessive drinking. *Neuron* 79: 406-407.
33. Doyon WM, Dong Y, Ostroumov A, Thomas AM, Zhang TA, et al. (2013) Nicotine decreases ethanol-induced dopamine signaling and increases self-administration via stress hormones. *Neuron* 79: 530-540.
34. Smith BR, Horan JT, Gaskin S, Amit Z (1999) Exposure to nicotine enhances acquisition of ethanol drinking by laboratory rats in a limited access paradigm. *Psychopharmacology (Berl)* 142: 408-412.
35. Lopez-Moreno JA, Trigo-Diaz JM, Rodriguez de Fonseca F, Gonzalez Cuevas G, Gomez de Heras R, et al. (2004) Nicotine in alcohol deprivation increases alcohol operant self-administration during reinstatement. *Neuropharmacology* 47: 1036-1044.
36. Dyr W, Koros E, Bienkowski P, Kostowski W (1999) Involvement of nicotinic acetylcholine receptors in the regulation of alcohol drinking in Wistar rats. *Alcohol and alcoholism* 34: 43-47.
37. Nadal R, Samson HH (1999) Operant ethanol self-administration after nicotine treatment and withdrawal. *Alcohol* 17: 139-147.
38. Sharpe AL, Samson HH (2002) Repeated nicotine injections decrease operant ethanol self-administration. *Alcohol* 28: 1-7.
39. Smith AM, Kelly RB, Chen WJ (2002) Chronic continuous nicotine exposure during periadolescence does not increase ethanol intake during adulthood in rats. *Alcohol Clin Exp Res* 26: 976-979.
40. Kemppainen H, Hyytia P, Kiianmaa K (2009) Behavioral consequences of repeated nicotine during adolescence in alcohol-preferring AA and alcohol-avoiding ANA rats. *Alcohol Clin Exp Res* 33: 340-349.
41. Grunberg NE, Morse DE, Maycock VA, Kozlowski LT (1985b) Changes in overwrap and butt length of American filter cigarettes. An influence on reported tar yields. *New York State J Med* 85: 310-312.
42. Grunberg NE, Bowen DJ, Maycock VA, Nespor SM (1985a) The importance of sweet taste and caloric content in the effects of nicotine on specific food consumption. *Psychopharmacology (Berl)* 87: 198-203.
43. Grunberg NE (1982) The effects of nicotine and cigarette smoking on food consumption and taste preferences. *Addict Behav* 7: 317-331.
44. Grunberg NE, Bowen DJ, Winders SE (1986) Effects of nicotine on body weight and food consumption in female rats. *Psychopharmacology (Berl)* 90: 101-105.
45. LeSage MG, Burroughs D, Pentel PR (2006) Effects of nicotine withdrawal on performance under a progressive-ratio schedule of sucrose pellet delivery in rats. *Pharmacol Biochem Behav* 83: 585-591.
46. Yuan M, Cross SJ, Loughlin SE, Leslie FM (2015) Nicotine and the adolescent brain. *The J Physiology* 593: 3397-3412.