

Falls in Parkinson Disease

Abraham Lieberman^{1*}, Aman Deep¹ and Thurmon Lockhart²

¹Muhammad Ali Parkinson Center (MAPC) at Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, AZ 85013, USA

²Department of Bio-Engineering, Gait Laboratory, Arizona State University, Tempe, Arizona, USA

Abstract

Falls are a major risk for Parkinson disease patients (PD). We sought to understand why patients fall. In one study we followed 404 PD patients for a year, 204 of who fell at least once. We did not distinguish between single and recurrent fallers. In a second study we followed 249 different patients for a year, 205 of whom fell once and 44 fell more than once (recurrent fallers).

In the first study fallers were significantly older, 72.6 ± 5.8 versus 66.9 ± 6.1 years, had PD significantly longer 9.3 ± 3.4 versus 5.4 ± 2.2 years, had significantly higher (worse) motor Movement Disorder Society (MDS) Unified Parkinson Disease Rating Scale (UPDRS) scores 29.3 ± 10.9 versus 18.9 ± 8.3 . Fallers walked with significantly shorter steps: 0.45 ± 0.09 versus 0.60 ± 0.13 m. Assessing step length, a simple "bedside" test, was a useful aid in predicting who might fall.

In the second study recurrent fallers had PD significantly longer, 12.6 ± 7.0 versus 5.9 ± 4.5 years, had significantly higher motor MDS UPDRS scores 31.2 ± 12.7 versus 19.7 ± 8.3 and walked with significantly shorter steps: 0.37 ± 0.18 m versus 0.52 ± 0.19 m. A major difference between recurrent and single fallers was an inability of recurrent fallers to stand on one leg for <3 s: 95% versus 11%, odds ratio 178 CI 95% 39.5-801.2 Single fallers who are unable to stand on one leg for <3 s are at risk for recurrent falls.

The short step, we believe, is an adaptation to postural instability, inability to maintain an upright position. The inability to stand on one leg, by depriving the patient of roughly half their proprioception, is a measure of risk for falls. Parkinson disease is viewed as a motor disorder, but impaired proprioception, a sensory symptom, is a major cause of recurrent falls.

Keywords: Parkinson disease; Falls; Freezing of gait

Introduction

Falls are a major cause of disability in Parkinson disease (PD) [1,2]. In a study of 761 hospital admissions for PD only 15% were for management of PD, while 39% were for falls [1]. A major risk for recurrent falls is a previous history of a fall. PD patients who fall once may do so because of PD or because of factors such as poor eye-sight, leg weakness and environmental hazards [3,4].

There is variability in the reported prevalence of falls in PD: from 11% to 68% [3-14]. The variability depends on whether specific fall risk factors are excluded. These include visual loss, orthostatic hypotension resulting from anti-hypertensives or imbalance resulting from tranquilizers, sedatives, or alcohol. The variability in reporting also depends upon whether only serious falls (requiring medical attention) or non-serious falls were recorded [3-14]. The variability also includes whether persons with evolving atypical Parkinson disorders such as Progressive Supranuclear Palsy (PSP) or Multiple System Atrophy (MSA) persons but with a high predilection to fall were excluded [6].

Methods

In 2010-2011 we saw 460 PD patients [3]. We excluded patients with atypical Parkinson disorders. We excluded patients with PD and dementia, Mini-Mental Status Examinations, MMSE, <24 . We excluded patients who were legally blind. We excluded patients with orthostatic hypotension. Although orthostatic hypotension can be part of PD, it can also result from the use of anti-hypertensives, diuretics or dehydration [15,16]. As we often could not distinguish between them we excluded such patients. We excluded patients with neuropathy when it resulted in impaired proprioception or weakness. We excluded patients with major orthopedic problems [13,14]. The number of excluded patients was 56.

We analyzed 404 PD patients: 214 males, 190 females. All were examined every 4 months as part of their routine office visit. At each visit patients and care-givers were instructed to call us if they fell. Of the 404 persons with PD, 204, 50.5%, fell: 164 (80%) patients fell once, 41 fell (20%) more than once. We compared fallers with non-fallers.

We analyzed serious falls where all 4 limbs, the skull or buttocks hit the ground, with patients sustaining a fracture or soft tissue injury. All patients were examined using the motor MDS UPDRS [17] and selected subtests. All patients were studied using the BNI balance scale [18]. This included the ability of a patient to stand on one leg for at least 3 s. All patients walked 7.63 m (25 feet). Adjustments were made for height, the number of steps taken was counted, divided by 7.63 m and an average was obtained.

Next we did a second study 2011-2012 [19]. None of the patients were participants in the first study.

We analyzed 205 patients: 113 men, 92 women of whom 161 (79%) fell once and 44 (21%) fell more than once. All patients were informed that the information collected could be used for research but that they personally could not be identified. Approval for the analysis

***Corresponding author:** Abraham Lieberman, Founder, Muhammad Ali Parkinson Center, Director at Bob and Renee Parsons' Fall Reduction Center, Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, AZ 85013, USA, Tel: +1-6024064309; E-mail: Abraham.Lieberman@dignityhealth.org

Received May 26, 2016; Accepted July 11, 2016; Published July 18, 2016

Citation: Lieberman A, Deep A, Lockhart T (2016) Falls in Parkinson Disease. J Alzheimers Dis Parkinsonism 6: 248. doi: [10.4172/2161-0460.1000248](https://doi.org/10.4172/2161-0460.1000248)

Copyright: © 2016 Lieberman A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

was obtained by the St. Joseph's Hospital institutional review board. No patients were compensated. As the evaluations were part of the patient's routine care no special consent other than the standard signed consent obtained from all patients at the time their visit was obtained. Continuous variables were analyzed using t-tests and categorical variables were analyzed using chi-square tests. We used the SAS 8.01 statistical software package. The first study was analyzed separately from the second study.

Results

In the first study of 404 PD patients, 50.5% fell (204). Among the fallers, 112 (54%) were males, 92 were females (46%). Among the non-fallers 112 (56%) were males, 88 (44%) were females. Among the 204 fallers, 41 (20%) fell more than once. In this study we only looked at fallers (single and recurrent) versus non-fallers. Age, duration of PD, MDS UPDRS motor scores were significantly higher (worse) in the fallers (Table 1). *Step length was significantly shorter* (Table 1). There were no differences in BNI subtests of One Legged Stance or Turning. Among fallers 43 patients, 21.3% had FOG, 161 (78.9%) did not. Among non-fallers 21 patients, 10.5% had FOG, 179 patients, 89.5% did not. *Fallers were more likely to have FOG than non-fallers: Odds Ratio 2.28, 95% CI 1.30-3.99, P<0.0042.*

Among fallers 84% were on levodopa, among non-fallers 80% were on levodopa. Forty percent of fallers and 29% of non-fallers had dyskinesias. Although a higher percent of fallers had dyskinesias we cannot comment on the contribution of dyskinesias to falls because we recorded only the presence, not the severity of dyskinesias.

In the second study, 205 PD patients fell, 44 of who fell at least twice. PD duration and MDS UPDRS Motor Score were significantly higher in recurrent fallers (Table 2). Step length was significantly shorter (Table 2). MDS UPDRS sub-tests of Gait, Pull and FOG were significantly worse and One Legged Stance and Turning sub-tests were significantly worse in recurrent fallers (Tables 1 and 2). Sub-tests of Postural Stability including One Legged Stance and Pull sub-test were worse than sub-tests of Locomotion including Gait sub-test and step length.

Among single fallers 87.5% were on levodopa, among recurrent fallers 100% were on levodopa. 40% of single fallers had dyskinesias, 56% of recurrent fallers had dyskinesias.

Discussion

The prevalence of falls, at least 50%, in both studies is comparable to that reported by others [4-10]. It's estimated that 33% of all people

over age 65 years fall at least once a year. The costs associated with fall-related injuries are high and estimated to reach \$30 billion by 2020 by the Rand Corporation [20]. These figures are likely to be higher in PD, because a higher percent of people with PD fall and are more likely to fall repeatedly. If falls, and especially recurrent falls, can be reduced in PD, they possibly can be reduced in the elderly [20-24].

Fallers have PD longer and are more disabled. However, in an individual patient who falls, the duration of PD may be insufficient to predict another fall. And in an individual patient the MDS UPDRS motor score may be insufficient to predict another fall. This is because the MDS UPDRS is weighted toward upper extremity rigidity, tremor and bradykinesia, metrics that are less likely to predict falls.

Fallers take significantly shorter steps. This in our opinion is an adaptation to impaired postural stability. Freezing of gait, the "ultimate short step", may be an extreme example of such an adaptation [24,25].

Recurrent fallers in the second study were significantly more likely to be unable to stand on one foot for <3 s 95.4% of recurrent fallers compared to 10.8% of single fallers, odds ratio 177.9 CI 95% 39.5-801.2, p<0.001. This was the best predictor of recurrent falls (Table 2) [26].

In a seminal study, Purdon-Martin studied evaluated 130 patients with post-encephalitic Parkinson [27]. He wrote:

In the upright position man stands on a base which is narrow and small relative to his height. A manikin or dummy figure is easily knocked over and man himself is stable only as long as his center of gravity remains within the vertical projection of his base.... Some physical instability is widespread among our patients and the more severely affected are remarkable in that they are unable to preserve their equilibrium not only when they are standing but also when they are sitting.

Purdon- Martin attributed postural instability to a combination of proprioceptive and vestibular dysfunction (sensory symptoms).

Locomotion, the rhythmical movement of the extremities, is generated in the spinal cord [28]. A hierarchy of supra-spinal centers in the cortex and basal ganglia signal the spinal cord. This supra-spinal command network is necessary for initiating walking, turning and stopping. A part of this locomotion network is under dopaminergic control. This motor network is impaired in PD and responds, in part, to dopaminergic therapy [28-31].

The locomotor network interacts with a postural stability network that includes inputs from proprioceptors in the muscles, joints, and tendons as well as inputs from the vestibular nuclei and visual system. This sensory network is not under dopaminergic control [29-34].

Proprioceptive inputs from the lower extremities and their integration into the CNS may, as postulated by Purdon-Martin, be most important in determining postural stability. The inability of recurrent fallers to stand on one leg may be related to an approximately 50% reduction of their proprioceptive input from the ground. Future studies of these phenomena will involve inertial monitoring units and a "fall simulator" to clarify the contribution of impaired body sway, dyskinesias and impaired sensory inputs to falls. Parkinson disease, viewed as a "motor disorder", has a prominent sensory component, one refractory to dopaminergic drugs.

References

1. Temlett JA, Thompson PD (2006) Reasons for admission to hospital for Parkinson's disease. Intern Med J 36: 524-526.

	Non-Fallers	Fallers	p-value
Age	66.9 ± 6.1	72.6 ± 5.8	0.019
PD Duration	5.4 ± 2.2	9.3 ± 3.7	0.026
MDS-UPDRS Motor Score	18.9 ± 8.3	29.3 ± 10.9	<0.01
Step Length Meters/Step	0.6 ± 0.13	0.45 ± 0.09	<0.01

Table 1: Study 1: Comparison age, PD duration and MDS UPDRS, motor scores fallers vs. non-fallers.

	Single-Fallers 161 patients	Recurrent Fallers 44 patients	p-value
Age (years)	67.4 ± 8.7	69.7 ± 9.3	NS
PD Duration (years)	5.9 ± 4.5	12.6 ± 7.0	0.001 *
MDS UPDRS Motor Score	19.7 ± 8.3	32.3 ± 12.6	0.01*
Step length Meters/Step	0.52 ± 0.12	0.31 ± 0.12	<0.01*

Table 2: Study 2: Comparison single versus recurrent fallers.

2. Johnell O, Melton LJ, Atkinson EJ, O'Fallon WM, Kurland LT (1992) Fracture risk in patients with parkinsonism: A population-based study in Olmsted County, Minnesota. *Age Ageing* 21: 32-38.
3. Lieberman A, Krishnamurthi N, Dhall R (2014) Comparison of Parkinson disease patients who fell once with patients who fell more than once. *Alzheimer's Disease & Parkinsonism* 4: 1-4.
4. Pickering RM, Grimbergen YA, Rigney U, Ashburn A, Mazibrada G, et al. (2007) A meta-analysis of six prospective studies of falling in Parkinson's disease. *Mov Disord* 22: 1892-1900.
5. Schrag A, Ben-Shlomo Y, Quinn N (2002) How common are complications of Parkinson's disease? *J Neurol* 249: 419-423.
6. Wielinski CL, Erickson-Davis C, Wichmann R, Walde-Douglas M, Parashos SA (2005) Falls and injuries resulting from falls among patients with Parkinson's disease and other parkinsonian syndromes. *Mov Disord* 20: 410-415.
7. Wood BH, Bilclough JA, Bowron A, Walker RW (2002) Incidence and prediction of falls in Parkinson's disease: A prospective multidisciplinary study. *J Neurol Neurosurg Psychiatry* 72: 721-725.
8. Lim I, van Wegen E, Jones D, Rochester L, Nieuwboer A, et al. (2008) Identifying fallers with Parkinson's disease using home-based tests: Who is at risk? *Mov Disord* 23: 2411-2415.
9. Ashburn A, Stack E, Pickering RM (2001) Predicting fallers in a community-based sample of people with Parkinson's disease. *Gerontology* 47: 277-281.
10. Bloem BR, Grimbergen YA, Cramer M, Willemsen M, Zwiderman AH (2001) Prospective assessment of falls in Parkinson's disease. *J Neurol* 248: 950-958.
11. Dennison AC, Noorigian JV, Robinson KM, Fisman DN, Cianci HJ, et al. (2007) Falling in Parkinson disease: Identifying and prioritizing risk factors in recurrent fallers. *Am J Phys Med Rehabil* 86: 621-632.
12. Latt MD, Lord SR, Morris JG, Fung VS (2009) Clinical and physiological assessments for elucidating falls risk in Parkinson's disease. *Mov Disord* 24: 1280-1289.
13. Kerr GK, Worringham CJ, Cole MH, Lacherez PF, Wood JM, et al. (2010) Predictors of future falls in Parkinson disease. *Neurology* 75: 116-124.
14. Paul SS, Canning CG, Sherrington C, Lord SR, Close JC, et al. (2013) Three simple clinical tests to accurately predict falls in people with Parkinson's disease. *Mov Disord* 28: 655-662.
15. Matinolli M, Korpelainen JT, Korpelainen R, Sotaniemi KA, Myllylä VV (2009) Orthostatic hypotension, balance and falls in Parkinson's disease. *Mov Disord* 24: 745-751.
16. Vaugoyeau M, Hakan H, Azulay J-P (2011) Proprioceptive impairment and postural orientation control in Parkinson's disease. *Human Movement Science* 30: 405-414.
17. Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, et al. (2008) Movement disorder society-sponsored revision of the unified Parkinson's disease rating scale (MDS-UPDRS). Scale Presentation and Clinometric Testing Results *Mov Disord* 21: 2129-2170.
18. Lieberman A, Dhanani S, Dhall R (2012) A quick and easy-to-use clinical scale to assess balance in Parkinson's disease. *J Parkinsonism & Restless Legs Syndrome* 2: 67-71.
19. Lieberman A (2014) Falls in Parkinson disease: The relevance of short steps. *J Novel Physiotherapies* 4: 1-4.
20. Shekelle P, Maglione M, Chang J. Evidence report and evidence-based recommendations: Falls prevention interventions in the Medicare population rand corporation for US Department of Health & Human Services Contract Number 500-98-0281 from 30 Sept 1998-29 Sept 2003.
21. Protas EJ, Mitchell K, Williams A, Qureshy H, Caroline K, et al. (2005) Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabilitation* 20: 183-190.
22. Ashburn A, Fazakarley L, Ballinger C, Pickering R, McLellan LD, et al. (2007) A Randomized Trial of a Home Based Exercise Programme to Reduce the Risk of Falling Among People with Parkinson's Disease. *J Neurol Neurosurg Psychiatry* 78: 678-684.
23. Allen NE, Canning CG, Sherrington C, Lord SR, Latt MD, et al. (2010) The effects of an exercise program on fall risk factors in people with Parkinson's disease: a randomized controlled trial. *Mov Disord* 25: 1217-1225.
24. Allen NE, Sherrington C, Paul SS, Canning CG (2011) Balance and falls in Parkinson's disease: A meta-analysis of the effect of exercise and motor training. *Mov Disord* 26: 1605-1615.
25. Lieberman A (2006) Are freezing of gait (FOG) and panic related? *J Neurol Sci* 248: 219-222.
26. Plotnik M, Giladi N, Balash Y, Peretz C, Hausdorff JM (2005) Is freezing of gait in Parkinson's disease related to asymmetric motor function? *Ann Neurol* 57: 656-663.
27. Purdon- Martin JP. *The Basal Ganglia and Posture*, Pitman Medical Publishing Company LTD, 1967, London Pages 1-152
28. Bourane S, Grossmann KS, Britz O, Dalet A, Del Barrio MG, et al. (2015) Identification of a spinal circuit for light touch and fine motor control. *Cell* 160: 503-515.
29. Nova IC, Perracini MR, Ferraz HB (2004) Levodopa effect upon functional balance of Parkinson's disease patients. *Parkinsonism Relat Disord* 10: 411-415.
30. Bohnen NI, Müller ML, Koeppe RA, Studenski SA, Kilbourn MA, et al. (2009) History of falls in Parkinson disease is associated with reduced cholinergic activity. *Neurology* 73: 1670-1676.
31. Woollacott M, Shumway-Cook A (2002) Attention and the control of posture and gait: A review of an emerging area of research. *Gait Posture* 16: 1-14.
32. Do MC, Breniere Y, Brenguier P (1982) A biomechanical study of balance recovery during the fall forward. *J Biomech* 15: 933-939.
33. Hall LM, Brauer S, Horak F, Hodges PW (2010) Adaptive changes in anticipatory postural adjustments with novel and familiar postural supports. *J Neurophysiol* 103: 968-976.
34. De Nunzio AM, Nardone A, Schieppati M (2007) The control of equilibrium in Parkinson's disease patients: Delayed adaptation of balancing strategy to shifts in sensory set during a dynamic task. *Brain Res Bull* 74: 258-270.