

Farmer's Preference and Effect of Feeding Selected Local Forages with Concentrate on the Dry Matter Intake and Weight Gain Performance of Bonga Sheep under Alarigeta Farmer's Management

Zelek M*, Giorgis DH, Rete C and Silase TG

Bonga Agricultural Research Center, P.O.Box 101, Bonga, Ethiopia

Abstract

A study was conducted at Alarigeta kebele, which is located in southwestern Ethiopia of Kaffa Zone, Adiyo woreda. In an effort to address feed problem, this study was carried out with the objective of identifying farmers preference of local forages for sheep and evaluating the effect of feeding selected local forages on the dry matter intake and weight gain performance of Bonga sheep reared under farmers management. Group discussion was conducted to listing and ranking of the preferred plant species was done through a questionnaire that was administered through a reconnaissance survey. Thirty intact male yearling sheep were divided in to six groups of five sheep based on their initial body weight in randomized complete block design (RCBD). Treatments were consist of six different local forages selected by farmers in study area; T1 (*Convolvulus kilimandschari* Engl.), T2 (*Commelina benghalensis* L.), T3 (*Basella alba* L.), T4 (*Brugmansia suaveolens* Bercht.), T5 (*Bothriocline schimperi* Olivo) and T6 (*Triumfetta tomentosa* Boj.). A daily dry matter intake of T3 (4301 g/day) and T (4400 g/day) are significantly higher ($P < 0.05$) than other groups. The greatest body weight of sheep was recorded in T3 (9.6 kg) and T4 (9.4 kg) as compared to other treatment groups ($P < 0.05$).

Keywords: Bonga sheep; Dry matter intake; Local forage *Convolvulus kilimandschari* Engl CL.; *Basella alba* L., *Brugmansia suaveolens* Bercht., *Bothriocline schimperi* Olivo., *Triumfetta tomentosa* Boj

Introduction

Feed shortage is among the prominent setbacks of the livestock sector in Ethiopia resulting in a low contribution of the sector for the nationwide gross domestic product, which is in contrary to the large population of livestock species in the country. The major feed resources in the country are crop residues and natural pasture with agro-industrial manufactured feeds contributing much less [1]. The highland areas of the nation, which have dominantly crop-livestock production systems are recognized to be under stress because of shrinking cultivated areas per household, land degradation and reduced feed availability [2].

Though increased utilization of agro-industrial by-products has been reported [3], they are not available, affordable or feasible for most of the farmers in the highlands of Ethiopia. Specifically to study area, since the bulk of the population is depends on relief for subsistence of life supplementary feeding to animals is either very costly or unavailable. Consequently, it is imperative to search for low cost and alternative feed resources for the poor farmers of the area.

Exotic fodder species are increasingly being used to overcome shortage of fodder, however local forage species have advantages over exotic spp., which can fit with the existing farming system and are well adapted with the environment and with the economic realities of farmers, are best candidates for so. The main features of such plants as a feed resource is their high crude protein content (CP). Previous studies have shown that a range of 5% to 20% CP was recorded for the browse species in tropical countries [4-6]. Current works in Ethiopia have also reported a range of CP content of local forages between 9.7% and 24.5% [7-9].

Southwestern Ethiopia is well known in origin of coffee in the world and huge number of local plants. Thus, it is crucial to ameliorate the severe feed shortage of the area through efficient utilization of local forage by supporting the local feed utilization knowledge of the local farmers with scientific research outputs. This research was done to

investigate the farmer's preference and nutritional qualities of major local forage in the area, which will serve as a launch pad for future propagation and utilization.

Objective

To study the effect of locally available local feeds on weight gain of Bonga sheep in farmers management

To demonstrate and promote cost-benefit, short term and market oriented fattening strategies

Methodology

Location of study

The experiment was conducted in Southwestern Ethiopia, Kaffa Zone Alarigeta. Alarigeta kebele is located in Adiyo wereda 28 Km from Bonga town. Adiyo wereda has huge potential in sheep production.

Group discussion

The listing and ranking of the preferred plant species was done through a questionnaire that was administered through a reconnaissance survey. The questionnaire was administered to households that were selected through a systematic sampling method in Alarigeta Bonga sheep improvement cooperative members. The respondents were asked to mention the most important local feed

*Corresponding author: Zeleke M, Bonga Agricultural Research Center, P.O.Box 101, Bonga, Ethiopia, Tel: +251916286233; E-mail: mulukenforjesus@yahoo.com

Received January 30, 2017; Accepted February 20, 2017; Published February 27, 2017

Citation: Zelek M, Giorgis DH, Rete C, Silase TG (2017) Farmer's Preference and Effect of Feeding Selected Local Forages with Concentrate on the Dry Matter Intake and Weight Gain Performance of Bonga Sheep under Alarigeta Farmer's Management. J Fisheries Livest Prod 5: 222 doi: [10.4172/2332-2608.1000222](https://doi.org/10.4172/2332-2608.1000222)

Copyright: © 2017 Zelek M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

materials in the study area while the ranking was done through the pair wise ranking method [10].

Farmers (Alarigeta sheep cooperative members) will be informed about low-cost, short-term & market-oriented fattening research project. According to willingness to practice fattening, thirty farmers for thirty sheep were selected.

Feeding trial

Experimental animal and management: Thirty intact male local lambs (yearlings, age less than two years) was selected from Alarigeta cooperative to conduct a feeding experiment. The sheep were ejected from breeding ram selection from Alarigeta cooperative. Selection criteria's of sheep are; general alertness, active feeding, rumination, no visible wound mouth, visible sign of disease. The age of the animal was subjectively determined by looking at their record. The sheep was quarantined for 21 days and during quarantine period, all sheep were dewormed using Albendazol against internal parasites, and vaccinated for pasteurellosis and anthrax with 1ml ovine pasteurellosis vaccine and 0.5 ml anthrax vaccine per sheep, respectively.

Feed preparation and feeding: Six outstanding local forages were selected by pervious study and farmer's preference. They were harvested, dried and stored under shed to keep the quality, chopped to approximately a size of 2-3 cm and used as basal diet or control diet throughout the experimental period. Commercial mineral mix was fed 50 g/day/animal based on the recommended level set by manufacturer. Wheat bran (WB) and Peanut (PN) were purchased and mixed at a ratio of 40% PN: 60% WB offered at 300 g DM/head/day. The concentrate was mixed in a way that the CP content of mixture is on average 18% CP on dry matter basis for the fulfillment of protein requirement of growing sheep with body weight of 15-20 kg. Individual feed troughs for natural pasture hay and mineral supplements as well as water troughs were provided separately for each experimental animal.

Experimental design and treatments

Randomized complete block design (RCBD) with six treatments consisting of six sheep per treatment were used to conduct the experiment. After 21 days of quarantine period, the experimental lambs were blocked in to four treatment groups based on initial body weight. Treatments were composed of six local forage and commercial mineral mix (50 g/day/head).

Measurements

Dry matter intake: Following the quarantine period, and after an acclimatization period of 15 days to the experimental pens and diets, the feeding trail was conducted for 90 days. Feeds were offered at 08:00 hours and 16:00 h into equal portion and water offered free choice. Daily feed offered to the experimental animals and the corresponding refusal of every animal were measured and recorded during the experimental period. Both basal and supplement diets were offered separately and intake were determined by the difference between the amount of feed

given and refused every day (24 h). Samples were taken from batches of feed offered and orts, thoroughly mixed and sub-sampled (10%) for dry matter analysis. The sub-samples were kept frozen (-20°C) until analyses (Table 1).

Body weight change and feed conversion efficiency: Initial body weights of each animal were determined by taking mean of two consecutive day's weight after overnight fasting. Body weights of animal were measured during the feeding period of 90 days at ten days interval after overnight fasting. Body weight measurements were taken using spring balance. The weight was recorded for each sheep against its identification number on a weight register book. A body weight change was determined as a difference between the final and initial body weight. Average daily body gain was calculated as the difference between the final and initial body weight of the sheep divided by the number of feeding days.

Body weight gain and average daily body weight gain was calculated as follows:

$$\text{Weight gain} = \text{Final body weights} - \text{Initial body weights}$$

$$\text{Average daily body gain (ADG)} = \frac{\text{Final body weights} - \text{Initial body weights}}{\text{Number of feeding days}}$$

Statistical analysis

Data collected was analysed using the frequency and descriptive tools of the Statistical Package for Social Sciences (SPSS) version 20 (SPSS 2013) and weighted using the Likert scale weighting procedures [11].

A two-way analysis of variance (ANOVA) was followed for the experimental (feeding trial) data using SAS 2008 version 9.2. The ANOVA procedure followed randomized block design (RCBD). The treatment means were separated by least significant difference (LSD). Mean differences were considered significant when $P \leq 0.05$, whereas $0.05 < P < 0.10$ was considered to show a statistical tendency for difference. The appropriate model used for data analysis was depicted here under:

$$Y_{ij} = \mu + T_i + B_j + e_{ij}$$

Where: Y_{ij} = response variable

μ = overall mean

T_i = treatment effect

B_j = block effect

e_{ij} = random error

Result and Discussion

Farmers performance of local forages

In Table 2 shows local forage selected in terms of productivity, adaptation and high nutritive value. According to farmers these local forages have high nutritive value, productivity, adaptability, tolerance

Trt	Local Name (Kafenono)	Family Name	Botanical Name	Concentrate mix(50 g/day/head)
1	Yimbiro	Convolvulaceae	<i>Convolvulus kilimandschari</i> Engl.	CM
2	Nallexo	Commelinaceae	<i>Commelina benghalensis</i> L.	CM
3	Nophoo	Basellaceae	<i>Basella alba</i> L.	CM
4	Mogne Abeba	Solanaceae	<i>Brugmansia suaveolens</i> Bercht	CM
5	Yamesho	Asteraceae	<i>Bothriocline schimperii</i> . Oliv. & Hiern ex Benth.	CM
6	Mogecco	Tiliaceae	<i>Triumfetta tomentosa</i> Boj.	CM

Table 1: Treatment feed (local forage selected by farmers preference).

	Treatments, Mean SEM						Sig	
Intake (g/d)	T1	T2	T3	T4	T5	T6		
CM DM	300	300	300	300	300	300	-	ns
LF DM	3654 ^b	3900 ^b	4301 ^a	4400 ^a	2810 ^c	3700 ^b	0.83	**
Total DM	3954 ^b	4200 ^b	4621 ^a	4715 ^a	3110 ^c	4000 ^b	0.12	**

^{a,b,c,d} means within a row bearing a common superscript are significantly different; ns: non-significant/ $P>0.05$; **= $P<0.05$; SEM: Standard Error of Mean; DM: Dry Matter, CM: Concentrate Mix; LF: Local Forage

Table 2: Daily dry matter intake of Bonga sheep fed local forages with concentrate mix.

	Treatments, Mean SEM						Sig	
Parameters	T1	T2	T3	T4	T5	T6		
IBW (Kg)	32.8	33.4	33.2	32.1	31.8	32.8	0.16	ns
FBW (Kg)	40.6 ^b	40.8 ^b	42.8 ^a	41.5 ^a	38.4 ^c	40.7 ^b	0.33	**
BWG (Kg)	7.8 ^b	7.2 ^b	9.6 ^a	9.4 ^a	6.6 ^c	7.9 ^b	0.12	**
ADG (g/d)	86 ^b	80 ^b	106.6 ^a	104.4 ^a	73.4 ^c	87.7 ^b	1.16	**

^{a,b,c,d} means within a row bearing a common superscript are significantly different; ns: non-significant/ $P>0.05$; **= $P<0.05$; BWC: Body Weight Change; ADG: Average Daily Weight Gain; FBW: Final Body Weight; IBW: Initial Body Weight; Sig: Significance Level; SEM: Standard Error of Mean

Table 3: Body weight parameters of Bonga sheep fed local forages with concentrate mix.

and palatability. They used these forages for a long period of time for fattening, milk production and calf growth.

Dry matter intake

A trend of daily dry matter intake during the feeding period is presented in Table 3. As depicted in the table, intake of T3 (4301 g/day) and T (4400 g/day) are significantly higher ($P<0.05$) than other groups. This might be associated with increment in body weight and associated increase in intake to satisfy nutrient requirement of the animal. Farmers in the study area also say that, *Nophoo* and *Mogne Abeba* are highly palatable to animals. According to Table 3, statistically there is no significant difference ($P>0.05$) between T1 (3954), T2 (3954) and T6 (4000), but higher than ($P<0.05$) T5 (3110).

Weight gain performance

Live weight gain (g/day) of sheep during the experiment period are presented in Table 3. The greatest body weight of sheep was recorded in T3 (9.6 kg) and T4 (9.4 kg) as compared to other treatment groups ($P<0.05$). This could be due to well-balanced mixture of nutrient and palatability of *Basell aalba L.* and *Brugmansia suaveolens Bercht*

Conclusion and Recommendation

The finding of this investigation has shown that farmers prefer local forages according to their palatability and effect on weight gain of animals. Provided that the appropriate entry point is identified, the species investigated can be utilized to ameliorate the ever-increasing feed shortage of the area.

The dry matter intake and live weight gain of animal which feed T3 (*Basell aalba L.*) and T4 (*Brugmansia suaveolens Bercht.*) significantly higher than other groups.

However their future utilization and management needs some complementary and additional investigations on digestibility, agronomic attributes and carcass evaluation should be conducted.

Acknowledgements

We acknowledge the Southern Agricultural Research Institute, for funding and providing resources. We also acknowledge Bonga Agricultural Research Center, Livestock Research Case team members their assistance and encouraging comments.

References

1. Gebremedhin B, Hirpa A, Berhe K (2009) Feed marketing in Ethiopia: Results of rapid market appraisal. Improving Productivity and Market Success (IPMS)

of Ethiopian farmers project Working Paper 15. ILRI (International Livestock Research Institute), Nairobi, Kenya: 64.

2. Aune JB, Bussa MT, Asfaw FG, Ayele AA (2001) The ox ploughing system in Ethiopia: can it be sustained? *Outlook on Agriculture* 30: 275-280
3. Benin S, Ehui S, Pender J (2004) Policies affecting changes in ownership of livestock and use of feed resources in the highlands of northern Ethiopia. *J Afr Econmic* 13: 166-194.
4. Heneidy S T (1996) Palatability and nutritive value of some common plant species from the Aqaba gulf area of Sinai, Egypt. *J Arid Environ* 34: 115-123.
5. Le Houerou HN (1980) Chemical composition and nutritive value of browse in tropical West Africa. In: Le Houerou, HN (ed.) *Browse in Africa: the current state of knowledge*. International Livestock Center for Africa (ILCA), Addis Ababa, Ethiopia: 261-297.
6. Shayo CM, Uden P (1999) Nutritional uniformity of neutral detergent solubles in some tropical browse leaf and pod diets. *Animal Feed Sci Tech* 82: 63-73.
7. Abdulatife M (2009) Assessment of pastoral perceptions, range condition and chemical composition of major feed resources in Chifra district of Afar Regional State, Ethiopia. An MSc Thesis presented to Haramaya University: 125p.
8. Wondatir Z (2010) Livestock production systems in relation with feed availability in the highlands and central rift valley of Ethiopia. An M Sc. Thesis Presented to Haramaya University.
9. Shimelis S (2010) Vegetation ecology, forage biomass and evaluation of nutritive value of selected browse species in Nechisar National Park, Ethiopia. An M Sc. Thesis Presented to Addis Ababa University.
10. Defoer T, Budelman A (2000) Managing soil fertility. A resource guide for Participatory Learning and Action Research (PLAR). Royal Tropical Institute. Amsterdam.
11. Vagia WM (2006) Likert-type scale response anchors. Clemson International Institute for Tourism and Research Development, Department of Parks, Recreation and Tourism Management. Clemson University.