

Fetal Megacystis

Maria G. Trevino Alanis¹, Areli S. Lopez Sanchez¹, Aaron A. Morin Juarez¹, Hector R. Martinez Menchaca², and Gerardo Rivera Silva^{1*}

¹Department of Basic Sciences, University of Monterrey, Mexico

²Department of Orthodontics, Pediatric Dentistry and Special Care, University of Louisville, USA

*Corresponding author: Gerardo Rivera Silva, Department of Basic Sciences, Division of Health Sciences, University of Monterrey, Mexico, Av. Ignacio Morones Prieto 4500, Pte, San Pedro Garza García, NL, Mexico, Tel: 1-52-8182151446; E-mail: gerardo.rivera@udem.edu

Received date: May 17, 2016, Accepted date: Jun 28, 2016, Published date: Jul 01, 2016

Copyright: © 2016 Trevino-Alanis MG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Clinical Image



Figure 1: Diagnosis of fetal megacystis by ultrasonography.

Mrs. LAL, 29 years old, G1P0A0, with gestational age of 15 weeks, with singleton pregnancy in breech presentation, slow movement just one time and heart rate of 158 -167 beats per minute. Her ultrasonography showed a gestational age by fetal body measurements of 15.4 weeks with a fetal weight of 123 grams. Furthermore, oligohydramnios, incipient renal pyelectasis, and megacystis with a volume of 10 cc (Figure 1). The diagnosis of fetal megacystis was confirmed. The patient was treated with intrauterine surgical correction with resolution of this malformation.