

Frailty and Survival in Patients with End-Stage Renal Disease

Ryota Matsuzawa¹ and Atsuhiko Matsunaga^{2*}

¹Department of Rehabilitation, Kitasato University Hospital, Sagami-hara, Japan

²Department of Rehabilitation, School of Allied Health Sciences, Kitasato University, Sagami-hara, Japan

Abstract

Frailty is highly prevalent in patients with end-stage renal disease who require hemodialysis therapy. Frailty has been proposed by Fried and colleagues, and predicts poor outcomes in hemodialysis patients. Muscle weakness is a major component of frailty. We previously performed a single-center, prospective, observational, cohort study among 190 Japanese patients undergoing hemodialysis and examined the association of muscle weakness in lower extremity with survival. As a result, muscle weakness in the lower extremities had a significant negative effect on survival in these patients. Muscle weakness is reversibly activated by exercise training or physical therapy in patients undergoing hemodialysis. Therefore, exercise training for patients with end-stage renal disease could improve their prognosis via correction of muscle weakness. In this paper, we reviewed muscle weakness, which is one of the components of frailty, in hemodialysis patients based on our previous study.

Keywords: Dialysis; Kidney; Renal; CKD; Frailty; Muscle weakness; Physical therapy; Rehabilitation; Exercise

Introduction

Frailty is highly prevalent in patients with end-stage renal disease who require hemodialysis therapy. Frailty predicts poor outcomes in community-dwelling elderly populations, patients with chronic kidney disease, and patients undergoing hemodialysis [1-7]. Frailty has been proposed by Fried and colleagues, and the gold standard for identifying frailty is based on the following five criteria: shrinking, weakness, poor endurance and energy, slowness, and a low physical activity level [8].

Weakness is one of the five components in frailty and is determined by handgrip strength. However, handgrip strength of patients undergoing hemodialysis is decreased by the presence of a hemodialysis shunt in the arm. In some patients, handgrip strength is reduced because of carpal tunnel syndrome, cubital tunnel syndrome, or destructive cervical spondylosis. Therefore, handgrip strength potentially fails to predict generalized muscle weakness in hemodialysis patients. However, lower extremity muscle strength is not directly influenced by a hemodialysis shunt or these comorbidities in the upper extremities. Furthermore, lower extremity muscle strength directly affects walking ability, standing balance function, activity of daily living, and quality of life, and is a target of intervention for physical therapists.

We review muscle strength, which is one of the components of frailty, in hemodialysis patients based on our previous study [9].

Lower extremity muscle strength and survival

We previously performed a single-center, prospective, observational, cohort study among 190 Japanese patients undergoing hemodialysis and examined the association of muscle weakness in lower extremity with survival [9]. Although all of the patients in our study population walked unassisted, approximately half of them had muscle strength lower than the cut-off value predicting whether people need assistance with walking. Over 7 years of follow-up, a total of 30 deaths occurred, with 17 attributed to cardio-cerebrovascular disease. The estimated cumulative survival was markedly different between the two groups, with 92.0% in the non-weakness group versus 75.6% in the weakness group (Figure 1). After adjustment for patients' characteristics and severity of comorbidity, muscle weakness in the lower extremities had a significant negative effect on survival. The hazard ratio in the weakness group was 2.73 (95% confidence interval = 1.14–6.52, P = 0.02) compared with the non-weakness group. Several possibilities can be considered regarding our results. Lower extremity muscle strength and physical activity level appear to interact. A sedentary lifestyle increases the risk of mortality in patients undergoing hemodialysis [10]. However, the underlying mechanisms remain to be determined.

Strength exercises in hemodialysis patients

Resistance training successfully enhances skeletal muscle and improves muscle strength in frail elderly populations and in those with chronic diseases. Cheema et al. investigated the effectiveness of resistance training on skeletal muscle quantity and strength in patients




Figure 1: Muscle weakness and survival in hemodialysis patients (Modified from reference 9). Muscle weakness: Isokinetic knee extensor muscle strength (kg) was divided by dry weight and expressed as percentage (%). A cut-off point of muscle weakness was 40% in Japan. People with values under this cut-off point may need assistance with walking.

***Corresponding author:** Atsuhiko Matsunaga, Department of Rehabilitation, School of Allied Health Sciences, Kitasato University, 1-15-1 Kitasato, Sagami-hara, Kanagawa 252-0373, Japan, Tel: +81-42-778-9970; E-mail: atsuhikonet@gmail.com

Received February 09, 2016; **Accepted** February 23, 2016; **Published** February 29, 2016

Citation: Matsuzawa R, Matsunaga A (2016) Frailty and Survival in Patients with End-Stage Renal Disease. J Yoga Phys Ther 6: 227. doi:10.4172/2157-7595.1000227

Copyright: © 2016 Matsuzawa R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

who received maintenance hemodialysis [11]. They found that resistance training improved muscle strength, but not muscle quantity. A meta-analysis showed the effect of exercise training on muscle strength in hemodialysis patients in 2014 [12]. In this meta-analysis, analysis was performed using 10 previously reported randomized, controlled trials, and exercise training was associated with improved muscle strength in these patients.

However, because almost all hemodialysis patients participated in these randomized, controlled trials of exercise training were younger than those in the clinical setting, future clinical trials for frail elderly patients undergoing hemodialysis are required to apply the evidence of exercise training in a clinical setting. Furthermore, we must encourage an active participate of frail elderly patients in exercise training program by not only physical therapist but also care manager [13].

Summary and Conclusion

Frailty is associated with poor outcomes in patients undergoing hemodialysis. Muscle weakness is a major component of frailty and is reversibly activated by exercise training or physical therapy in patients undergoing hemodialysis. Therefore, exercise training for patients with end-stage renal disease could improve their prognosis via correction of muscle weakness.

Financial Support

Kitasato University Research Grant.

References

1. Drubbel I, de Wit NJ, Bleijenberg N, Eijkemans RJ, Schuurmans MJ, et al. (2013) Prediction of adverse health outcomes in older people using a frailty index based on routine primary care data. *J Gerontol A Biol Sci Med Sci* 68: 301-308.
2. Rothman MD, Leo-Summers L, Gill TM (2008) Prognostic significance of potential frailty criteria. *J Am Geriatr Soc* 56: 2211-2216.
3. Rockwood K, Howlett SE, MacKnight C, Beattie BL, Bergman H, et al. (2004) Prevalence, attributes, and outcomes of fitness and frailty in community-dwelling older adults: report from the Canadian study of health and aging. *J Gerontol A Biol Sci Med Sci* 59: 1310-1317.
4. Roshanravan B, Khatri M, Robinson-Cohen C, Levin G, Patel KV, et al. (2012) A prospective study of frailty in nephrology-referred patients with CKD. *Am J Kidney Dis* 60: 912-921.
5. Roshanravan B, Robinson-Cohen C, Patel KV, Ayers E, Littman AJ, et al. (2013) Association between physical performance and all-cause mortality in CKD. *J Am Soc Nephrol* 24: 822-830.
6. Bao Y, Dalrymple L, Chertow GM, Kaysen GA and Johansen KL (2012) Frailty, dialysis initiation, and mortality in end-stage renal disease. *Arch Intern Med* 172: 1071-1077.
7. Johansen KL, Dalrymple LS, Delgado C, Kaysen GA, Kornak J, et al. (2014) Comparison of self-report-based and physical performance-based frailty definitions among patients receiving maintenance hemodialysis. *Am J Kidney Dis* 64: 600-607.
8. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, et al. (2001) Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 56: M146-156.
9. Matsuzawa R, Matsunaga A, Wang G, Yamamoto S, Kutsuna T, et al. (2014) Relationship between lower extremity muscle strength and all-cause mortality in Japanese patients undergoing dialysis. *Phys Ther* 94: 947-956.
10. Matsuzawa R, Matsunaga A, Wang G, Kutsuna T, Ishii A, et al. (2012) Habitual physical activity measured by accelerometer and survival in maintenance hemodialysis patients. *Clin J Am Soc Nephrol* 7: 2010-2016.
11. Cheema B, Abas H, Smith B, O'Sullivan A, Chan M, et al. (2007) Progressive exercise for anabolism in kidney disease (PEAK): a randomized, controlled trial of resistance training during hemodialysis. *J Am Soc Nephrol* 18: 1594-1601.
12. Heiwe S and Jacobson SH (2014) Exercise training in adults with CKD: a systematic review and meta-analysis. *Am J Kidney Dis* 64: 383-393.
13. Ciccone MM, Aquilino A, Cortese F, Scicchitano P, Sassara M, et al. (2010) Feasibility and effectiveness of a disease and care management model in the primary health care system for patients with heart failure and diabetes (Project Leonardo). *Vasc Health Risk Manag* 6: 297-305.