

## From Understanding the Sources of Pollution to Control the Quality of Coastal Sediment from Non-Point Pollution Sources

Shao-Wei Liao\*

Department of Tourism Management, Tajen University, Pingtung, Taiwan

The end-point treatment of point pollution sources is the best way to improve the quality of water and prevent coastal pollution, all over the world. Nevertheless, non-point pollution sources that influence the quality of surface water are serious concerns. Terrestrial pollution was deposited and accumulated over the estuary and coastal areas by adhering to the suspended particles in the rivers. Therefore, the pollution status and its history can be understood by analyzing and monitoring the sediments in the estuary as well as in the lagoon area. Sediments not only act as the carrier of pollution, but also as a potential secondary source of pollution in aquatic systems [1,2].

High concentrations of PAHs in sediments are usually high in highly industrialized and urbanized locations [3,4]. Moreover, highly anthropogenic activities are generally recognized to be the most important source of PAHs release into the environment [5]. At coastal areas, anthropogenic polycyclic aromatic hydrocarbons are introduced via urban runoff [6], industrial processes, vehicle exhausts, and spillage of fossil fuels [7]. Meanwhile, heavy metals discharged into a river system by natural or anthropogenic sources during their transport are distributed between the aqueous phase and bed sediments. Therefore, sediments are regarded as the ultimate sinks for heavy metal cations [8]. Heavy metal residing in contaminated habitats may accumulate in microorganisms, aquatic flora and fauna, which in turn may enter the human food chain and result in health problems [9,10].

Multivariate approaches have been used successfully to support the interpretation of complex field measurements, and to extract meaningful information from such databases [11,12]. Canonical Discriminate Analysis (CDA) determines how a set of quantitative variables may differentiate among many known classes. Finally, which group the unclassified values (samples in the lagoon) are classified into can be predicted correctly by using CDFs [13,14]. The approach also allows for relationships among the groups to be graphically represented by plotting the canonical scores of sample observations and have been used popularly to find sources of pollution.

Effective pollution control and water-resource management, especially in lagoons, depend upon identifying the main sources of pollution. The results might yield useful information concerning estuary recovery and water resources management and might be applicable to other basins with similar characteristics that are experiencing similar coastal environmental issues.

### References

1. Calmano W, Wolfgang A, Forstner U (1990) Exchange of heavy metals between sediment components and water. *Metal speciation in the environment* 23: 503-522.
2. Sin SN, Chua H Lo W, Ng LM (2001) Assessment of heavy metal cations in sediments of ShingMun River Hong Kong. *Environ Int* 26: 297-301.
3. Witt G (1995) Polycyclic aromatic hydrocarbons in water and sediment of the Baltic Sea. *Mar Pollut Bull* 31: 237-248.
4. Ashley JTF, Baker JE (1999) Hydrophobic organic contaminants in surficial sediments of Baltimore harbor: inventories and sources. *Environ Toxicol Chem* 18: 838-849.
5. Culotta L, De Stefano C, Gianguzza A, Mannino MR, Orecchio S (2006) The PAH composition of surface sediments from Stagnone coastal lagoon, Marsala (Italy). *Mar Chem* 99: 117-127.
6. Mc Cready S, Slee DJ, Birch GF, Taylor SE (2000) The distribution of polycyclic aromatic hydrocarbons in surface sediments of Sidney Harbor, Australia. *Mar Pollut Bull* 40: 999-1006.
7. Pettersen H, Naf C, Broman D (1997) Impact of PAHs outlets from an oil refinery on the receiving water area sediment trap fluxes and multivariate statistical analysis. *Mar Pollut Bull* 34: 85-95.
8. Gibbs RJ (1973) Mechanisms of trace metal transport in rivers. *Science* 180: 71-73.
9. Cook JA, Andrew SM, Johnson MS (1990) Lead, zinc, cadmium and fluoride in small mammals from contaminated grass-land established on fluorspar tailings. *Water Air and Soil Pollut* 51: 43-54.
10. Deniseger J, Erickson J, Austin A, Roch M, Clark MJR (1990) The effects of decreasing heavy metal concentrations on the biota of Buttle Lake, Vancouver Island, British Columbia. *Water Res* 24:403-416.
11. Simeonov V, Stratis JA, Samara C, Zachariadis G, et al. (2003) Assessment of the surface water quality in Northern Greece. *Water Res* 37: 4119-4124.
12. Singh KP, Malik A, Mohan D, Sinha S (2004) Multivariate statistical techniques for the evaluation of spatial and temporal variations in water quality of Gomti River (India)-a case study. *Water Res* 38: 3980-3992.
13. Liao SW, Gau HS, Lai WL, Chen JJ, Lee CG (2008) Identification of pollution of Tapeng Lagoon from neighbouring rivers using multivariate statistical method. *J Environ Manage* 88: 286-292.
14. Chung CY, Chen JJ, Lee CG, Chiu CY, Lai WL, et al. (2011) Integrated Estuary Management for Diffused Sediment Pollution in Dapeng Bay and Neighboring Rivers (Taiwan). *Environ Monit Assess* 173: 499-517.

\*Corresponding author: Shao-Wei Liao, Department of Tourism Management, Tajen University, Pingtung, Taiwan, Tel: +886 87624002 ext 3600; E-mail: [swliao@tajen.edu.tw](mailto:swliao@tajen.edu.tw)

Received October 19, 2013; Accepted October 21, 2013; Published October 25, 2013

Citation: Liao SW (2013) From Understanding the Sources of Pollution to Control the Quality of Coastal Sediment from Non-Point Pollution Sources. *J Pollut Eff Cont* 1: e103 doi: [10.4172/jpe.1000e103](http://dx.doi.org/10.4172/jpe.1000e103)

Copyright: © 2013 Liao SW. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.