

## Gene Silencing with Herbal Compounds against Bacteria

Moradi Fateme, Esmaeili Davoud\*, Hossein Saghi and Parya Baban Zadeh

Department of Microbiology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

### Abstract

Quorum sensing (QS) is an important process involved in bacterial survival and infections, recent research has focused on the development of therapeutic agents which prevent or manage bacterial pathogenesis by inhibiting bacterial QS. Inhibition of quorum sensing offers an alternative to antibiotic mediated bactericidal or bacteriostatic approach and reduces the risk for development of resistance. Inhibition of bacterial quorum sensing by attenuating the signals can prevent the development of bacterial and fungal virulence and successful establishment of infections. Understanding the quorum sensing inhibition activity of natural bioactive phytochemicals can lead to the discovery of novel compounds and development of more effective strategies in preventing and managing microbial infections. In this study the effect of *Satureja khuzestanica* extract, an endemic plant of Iran, on the expression level of QS gene in *P. aeruginosa* and *A. baumannii* were investigated.

**Keywords:** RT-PCR; *Satureja khuzestanica*; *Pseudomonas aeruginosa*; MIC; Carvacrol

### Introduction

For this purpose first is necessary expose organism with MIC concentration of herbal extracts. Then, microorganisms were treated with extract before and after treatment using genomic and proteomic techniques. Surprisingly, in this study the expression level of QS gene was decreased in the presence of *S. Khuzestanica*. However, the expression of gyrA gene (Housekeeping gene) that was used as an internal control was not altered before and after treatment with this herb. Based on the results, *S. Khuzestanica* could play a major role in lowering the *P. aeruginosa* resistance to drugs, by reducing QS gene expression. According to results of current research we hope in future be used it to the clinic with a wider range as a complementary therapy and also for surgery operation.

### Materials and Methods

For many years, control of bacterial infections by inhibiting microbial growth has been a primary approach of antimicrobial chemotherapy. An emerging problem associated with continual in discriminant use of this therapeutic strategy is the selection of resistant bacteria with higher levels of tolerance against broad-spectrum antibiotics.

Development of novel antibiotics that interfere with metabolism coupled with continued in discriminant use of antibiotics will only lead to evolution of new resistance mechanisms and pathways by bacteria. Recently, it has been recognized that there is a need for a strategy that can block very basic mechanisms of bacterial communication that appear to control bacterial virulence factors leading to pathogenicity. Emerging research has suggested that functions including swarming, biofilm formation, secretion of virulence factors and acquiring competency play an important role in successful and recurrent establishment of bacterial infections in living systems. These processes common to several bacterial diseases are now shown to be related to a cell density-dependent regulation of gene expression mediated by the recognition of signalling molecules called auto inducers (AI). This process sometimes referred to as 'quorum sensing' (QS) has been shown to modulate the expression of genes involved in processes related to survival, virulence and pathogenicity [1-3] and is mediated by small signalling molecules such as acetylated- homoserine lactones (AHLs).

### Results

Opportunistic Gram negative bacteria including *Pseudomonas* sp. have been shown to employ this type of AHL-mediated quorum sensing to regulate the expression of genes related to survival, virulence and competency [4]. Recently, several inhibitors of QS have been discovered from natural sources and have been produced synthetically. Some natural sources include *Australian macroalga*, *Deliseapulchra* and *green alga*, *Ulvalactua* that produce halogenated furanones and furanone-like pigments which interfere with quorum sensing [5,6]. These halogenated furanones synthesized in the laboratory have a high degree of AHL inhibition activity and were able to repress a number of genes controlled by QS. In addition, analogs of *S-adenosylmethionine* (SAM), such as *S-adenosylcysteine* and sinefungin can inhibit LuxI mediated synthesis of AHL but can interfere with eukaryotic pathways that use SAM. Recent studies have shown that interference with *P. aeruginosa* quorum signalling by metabolites from *Penicillium species* can increase susceptibility to these agents [7]. Unfortunately, all these compounds are highly reactive and toxic to the host which has limited their therapeutic applications [8] and potential usage in vivo. The current quest for new antimicrobials is therefore aimed at discovering nontoxic inhibitors of QS that can be used for treatment of bacterial infections in humans [8].

### Discussion

Natural products especially plants used in traditional medicines are a promising source for deriving molecules that can potentially inhibit quorum sensing [9-11]. These plants can offer a large and attractive repertoire for the discovery of quorum sensing inhibitors. They are of particular importance as these have been used for thousands of years for the treatment and management of diseases and may have few

**\*Corresponding author:** Esmaeili Davoud, Department of Microbiology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran, Tel: 98 21 805 2941; E-mail: [esm114@gmail.com](mailto:esm114@gmail.com)

Received June 04, 2015; Accepted July 06, 2015; Published July 09, 2015

**Citation:** Fateme M, Davoud E, Saghi H, Zadeh PB (2015) Gene Silencing with Herbal Compounds against Bacteria. J Med Microb Diagn 4: 196. doi:10.4172/21610703.1000196

**Copyright:** © 2015 Fateme M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

side-effects and toxicity issues as with many antibiotic regimens and currently known QS inhibitors. Herbs, Spices and Medicinal Plants HSMP used in Hispanic cultures have been used for several centuries to treat common ailments, are well known for their antimicrobial effects on a variety of human pathogens [12,13]. However, few studies have investigated their QS related antivirulent activities including in *P. aeruginosa*. The body of literature discussing HSMP from Central/South American continent as sources of QSI is very limited [14-17]. We believe that some of the antimicrobial properties of HSMP may be contributed by the QS inhibiting phytochemicals present in it. *Satureja khuzestanica*, from Lamiaceae family is an Iranian endemic plant, famous for its medical uses as an analgesic and antiseptic in folk medicine [18]. It is mostly found in western and southern part of Iran [19]. Recently, antiviral, antibacterial, antifungal, and antiprotozoal effects were investigated from various species of *Satureja* [19,20]. However, the possible effect of *Satureja khuzestanica* on decreasing the resistance of *P.aerogenase* against antibiotics and the mechanisms involved have not yet been studies.

## Conclusion

The antibacterial activity of the *S. Khuzestanica*'soil might be due to main phenolic components, Carvacrol and Thymol [21]. Carvacrol is also found in Thyme, However, its high ratio in *S. khuzistanica* has discriminated this plant from other herbs with antimicrobial effects [22,23]. In view of this and with regard to the antimicrobial effect of *S. Khuzestanica* against *p. aerogenase* and also resistance of this strain to variety of antibiotics, this study aimed to test this hypothesis that *S. Khuzestanica* extract may alter the expression of QS gene ,and thus may lead to a lower susceptibility of this strain to antibiotics.

## References

1. Schauder S, Bassler BL (2001) The languages of bacteria. *Genes Dev* 15(12): 1468- 1480.
2. Lyon GJ, Muir TW (2003) Chemical signaling among bacteria and its inhibition. *Chem Biol* 10(11): 1007-1021.
3. Raffa RB, Iannuzzo JR, Levine DR, Saeid KK, Schwartz RC, et al. (2005) Bacterial communication (quorum sensing) via ligands and receptors: a novel pharmacologic target for the design of antibiotic drugs. *J Pharmacol Exp Ther* 312(2): 417-423.
4. Pesci EC, Milbank JBJ, Pearson JP, McKnight S, Kende AS, et al. (1999) Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 96: 11229-11234.
5. Givskov M, de Nys R, Manefield M, Gram L, Maximilien R, et al. (1996) Eukaryotic interference with homoserine lactone-mediated prokaryotic signalling. *J Bacteriol* 178: 6618-6622.
6. Egan S, James S, Kjelleberg S (2002) Identification and characterization of a putative transcriptional regulator controlling the expression of fouling inhibitors in *Pseudoalteromonastunicata*. *Appl Environ Microbiol* 68: 372-378.
7. Rasmussen TB, Skindersoe ME, Bjarnsholt T, Phipps RK, Christensen KB, et al. (2005) Identity and effects of quorum-sensing inhibitors produced by *Penicillium* species. *Microbiology* 151(Pt 5): 1325-13240.
8. Hentzer M, Givskov M (2003) Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections. *J Clin Invest* 112(9): 1300-1307.
9. Cowan MM (1999) Plant products as antimicrobial agents. *Clinical Microbiological Reviews* 12: 564-582.
10. Wallace RJ (2004) Antimicrobial properties of plant secondary metabolites. *Proc Nutr Soc* 63: 621-629.
11. Mihalik K, Crixell SH, McLean RJC, Vatter DA (2007) Dietary Phytochemicals as Quorum Sensing Inhibitors. *Fitoterapia. Journal for the Study of Medicinal Plants* 78: 302-310.
12. Rojas A, Hernandez L, Pereda-Miranda R, Mata R (1992) Screening for antimicrobial activity of crude drug extracts and pure natural products from Mexican medicinal plants. *J Ethnopharmacol* 35(3): 275-283.
13. Quinlan MB, Quinlan RJ, Nolan JM (2002) Ethnopharmacology and herbal treatments of intestinal worms in Dominica, West Indies. *J Ethnopharmacol* 80(1): 75-83.
14. Gomez-Beloz A, Chavez N (2001) The botanica as a culturally appropriate health care option for Latinos. *J Altern Complement Med* 5: 537-46.
15. Huxtable RJ (1983) Herbs along the western Mexican-American border. *Proc West Pharmacol Soc* 26: 185-191.
16. Mikhail N, Wali S, Ziment I (2004) Use of alternative medicine among Hispanics. *J Altern Complement Med* 10 (5): 851-859.
17. Lewis WH, Elvin-Lewis MPF (2003) *Medical Botany: Plants Affecting Human Health*, 2nd Edition, Wiley. New Jersey.
18. Scully C, Gorsky M, Lozada-Nur F (2003) The diagnosis and management of recurrent aphthous stomatitis: a consensus approach. *J Am Dent Assoc* 134(2): pp. 200-207.
19. Van Baren, C (2006) Triterpenic acids and flavonoids from *Satureja parvifolia*. Evaluation of their antiprotozoal activity. *Z Naturforsch C* 61(3-4): p. 189-192.
20. Tampieri MP (2005) The inhibition of *Candida albicans* by selected essential oils and their major components. *Mycopathologia* 159(3): p. 339-45.
21. Deans SG, S KP (1989) Antibacterial activity of summer savory (*Satureja hortensis* L) essential oil and its constituents. *J. Hortic. Sci.*, 64: p. 205-210.
22. Dorman HJD, Deans SG (2000) Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl. Microbiol.*, 88: p. 308-316.
23. Lambert RJW (2001) A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *J Appl Microbiol.* 91: p. 453-462.