

Genitalia Malformations in *Stramonita haemastoma* (Gastropoda: Muricidae) from Atlantic and Mediterranean Coast

Tahani El Ayari*, Anwar Mleiki and Najoua Trigui El Menif

Department of Sciences of Bizerta, University of Carthage, Laboratory of Environment Bio-monitoring, Group of Fundamental and Applied Malacology (LEB/GFAM), Bizerta, Tunisia.

Abstract

Stramonita haemastoma is a gonochoric marine gastropod. The examination of the genital tract has revealed the occurrence of malformations affecting the penis of males *S. haemastoma*, as for the females only aphyllia was recorded. Investigations of the genital tract malformations were made in snails sampled monthly from Bizerta Channel (Northern Tunisia), over one-year period of sampling (June 2009 to May 2010). *S. haemastoma* was sampled in 2013 from sixteen sites located in the Mediterranean and Atlantic coast. The development of such type of malformations was associated to the use of antifouling paints. However, this assumption should be supported by further bioassays.

Keywords: *Stramonita haemastoma*; Penis malformations; Aphyllia; Mediterranean coast; Atlantic coast

Introduction

Malformations in marine gastropods, often associated with pollution, may have severe consequences on the marine system [1,2]. For instance, imposex had caused reproductive failure and in consequence population decline of several gastropod species [1-5]. Undeniably, imposex is the most common malformation reported in marine gastropods; it affects the genital tract of normal female by imposing the development of male sexual characters onto female [6]. According to the literature, imposex is induced following an exposure to tributyltin (TBT) used in antifouling paints [7]. Several marine gastropod species were found affected by this abnormality, in 2005 Shi et al. [8] published a list with 260 species affected by imposex [8,9]. This list has considerably increased since this date with the addition of new affected species as *Heleobia australis* [10], *Nassarius mutabilis* [11] and *Plicopurpura pansa* [12].

TBT may induce another genital tract malformation called Dumpton Syndrome (DS), corresponding to the under-development of male sexual characters on both sexes (male and imposex affected female) [13]. The Dumpton syndrome is also known as aphyllia in marine gastropods, this deficiency was firstly observed in the snail *Nucella lapillus* sampled from Dumpton Gap (England) [13] and then in four other localities: Brest (France) [14], Brittany (France) [14], Spain [15] and Portugal [16]. DS was also recorded in other gastropods as *Ocenebrina erinacea* [17]. Lima [17] noted a high rate of aphyllic females *S. haemastoma* (42.6%) during an experimental induction of imposex. After that, [18] Queiroz et al. noted a high occurrence of aphyllia in females *S. haemastoma* sampled from the Northeast of Brazil and suggested the possibility of the presence of the allele responsible for the Dumpton Syndrome at high frequencies in *S. haemastoma* population. Male affected by DS may have an undersized penis, an under-developed vas deferens, in the extreme cases they may present a split prostate, and a completely absence of the penis [13]. Female affected by DS developed or not an excrescence instead of a penis, which is considered an advantage in severely TBT-polluted site [13].

Besides to those genital abnormalities, morphological malformations were reported to affect the penis [7,14,19-24], vas deferens [25] or both penis and vas deferens [26-28]. Penis malformations were reported in *Hexaplex trunculus* [22,24,26,28], *Bolinus brandaris* [26,29], *Hinia reticulata* [27], *Nucella lapillus* [14], *Cyclope neritea* [30,31], *Leucozonia nassa* and *Leucozonia ocellata* [21].

There has been some report on imposex in *Stramonita haemastoma* in any areas of the world but not much was known in this part of the country. Contrary to its congeneric species *H. trunculus* [7,22,24] and *B. brandaris* [26,29]. In the present study, we reported malformations affecting the genital tract of males and females imposex-affected *S. haemastoma* collected from both Mediterranean and Atlantic coast.

Materials and Methods

Adult gastropod, *S. haemastoma*, were sampled monthly during 12 months from June 2009 to May 2010 in a rocky site located in Bizerta Channel. In totality, a sample of 1035 gastropod/month with shell length 28 mm to 80 mm is collected. Sampling was performed by SCUBA and some time by hand (snorkelling) at a station located in the artificial channel linking the Bizerta lagoon to the Mediterranean Sea (Figure 1). The length of the channel of Bizerta is about 1500 m, 240 m broad and 12 m depth. It is characterized by strong currents, an intense maritime traffic and chemical and urban pollution [32-34]. The direct and indirect discharges of urban and industrial wastes have contaminated the lagoon by various toxic compounds such as organo-chlorinated pesticides [35,36], halogenated aromatic compounds like polychlorinated biphenyls (PCBs) [2], polycyclic aromatic hydrocarbons (PAHs) [37], heavy metals and organotin compounds [38,39].

30 to 45 individuals were collected in 2013 from 5 and 11 North-Eastern Atlantic and Western Mediterranean coasts, respectively (Figure 1 and Table 1). In the laboratory, specimens were anaesthetized in a 1:1 mixture of 7.5% MgCl₂ and seawater. Shell and penis length were measured using a vernier calliper to the nearest 0.1 mm. The genital tract was examined under a binocular microscope.

*Corresponding author: Tahani El Ayari, Department of Sciences of Bizerta, University of Carthage, Laboratory of Environment Bio-monitoring, Group of Fundamental and Applied Malacology (LEB/GFAM), Bizerta, Tunisia, Tel: +216 58 83 54 83; Fax: +216 72 59 05 66; E-mail: tahaniayari@yahoo.fr

Received May 15, 2017; Accepted July 10, 2017; Published July 12, 2017

Citation: El Ayari T, Mleiki A, El Menif NT (2017) Genitalia Malformations in *Stramonita haemastoma* (Gastropoda: Muricidae) from Atlantic and Mediterranean Coast. J Aquac Res Development 8: 493. doi: [10.4172/2155-9546.1000493](https://doi.org/10.4172/2155-9546.1000493)

Copyright: © 2017 El Ayari T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Sampling localities of *Stramonita haemastoma*.

| Areas | Sample code | Sample name | Year of collection | GPS coordinate | |
|-----------------------------|-------------|-----------------|--------------------|----------------|-------------|
| | | | | Latitude | Longitude |
| North-Eastern Atlantic | AB | Arcachon Bay | 2013 | 44°592602 N | -1°214601 E |
| | RF | Ria Formosa | 2013 | 36°988717 N | -7°983696 E |
| | BO | Bouznilca | 2013 | 33°828000 N | -7.145958 E |
| Western Mediterranean basin | DB | Dar Bouaaza | 2013 | 33°534499 N | -7°827555E |
| | ES S | Essaouira | 2013 | 31°692465 N | -9°815664 E |
| | THA | Etang de Than | 2013 | 43°393343 N | 3°702986 E |
| | ORA | Can | 2013 | 35°901409 N | -0°331933 E |
| | ALG | Algiers | 2013 | 36°45873 N | 2°50838 E |
| | ANN | Annaba | 2013 | 36°898241 N | 7°771432 E |
| | TAB | Tabarca | 2013 | 37°514243 N | 9°869430 E |
| | BC | Bizerta channel | 2010 to 2013 | 36136030 N | 11°101338 E |
| | SB | Sidi Bechir | 2013 | 37°345694 N | 9°735649 E |
| | KOR | Korbous | 2013 | 36°816849 N | 10°566951 E |
| | KEL | Kelibia | 2013 | 38°126280 N | 12°793723 E |
| | EH | El Houaria | 2013 | 37.033306 N | 11°059370 E |
| | PAL | Palermo | 2013 | 43°332742 N | 3°612416 E |

Table 1: Sample location of *Stramonita haemastoma*.




Figure 2: Penis malformations in males *Stramonita haemastoma*. a: normal penis; b–g: abnormal penises observed in Bizerta Channel (Tunisia); h–i: abnormal penises observed in Bouznikha (Morocco). ex 1: excrescence 1; ex 2: excrescence 2; ex 3: excrescence 3; f: foot; ot: ocular tentacle; p: penis; p1: penis 1; p2: penis 2; vd: vas deferens. Scale bars=1 mm.

Results

Penis malformations in males *S. haemastoma*

In male *S. haemastoma*, the normal penis has a bent shape, a large triangular tip, tapering gradually toward the snail head and without

flagellum (Figure 2a). At the center of the penis there is a penis duct which is a curved closed tube beginning from the vas deference and reaching the tip of the penis. This tube ensures the passage of the sperm from the prostate to the female during the period of reproduction. The length of the penis increases considerably during the period of

reproduction extending from May to October, increases in penis length may exceed 5 mm to 6 mm during this period. Malformations affecting the male penis were observed in 2 out of 16 stations. The first station is located in the Western Mediterranean basin (Bizerta Channel, Tunisia), the second one is in the North-Eastern Atlantic Ocean (Bouznika, Morocco). Penis malformations were mainly observed in Tunisian population, 9 males were found with malformed penis in Bizerta Channel, against 1 male in Bouznika station. Malformations observed in Bizerta Channel consist in the development of a bud tissue (excrescence) close to the tip (Figures 2b-2d), or at the half of the total penis length always at the posterior side (Figure 2e). The number of excrescences may reach three at the level of the tip of the main penis (Figure 2d). Another malformation consists in the differentiation of a supplementary small penis measuring 2.3 mm in length and without a penis duct (biphallid penis) (Figure 2f). In addition to the main penis which appear undersized (3.5 mm in length), filamentous and with an incomplete penis duct (Figure 2f). Another specimen was found with two penises, the main penis has a normal shape and length (17.6 mm), the second penis is more developed compared to the last case and with a complete penis-duct (3.7 mm in length) (Figure 2g). Four Aphallic males were found in Bizerta Channel. The malformation observed in Bouznika station corresponds to the development of a long penis (23 mm) in the shape of the Z letter (Figures 2h and 2i). The rates of malformed males were 0.86% and 2.38% in Tunisian population and Moroccan population, respectively.

Aphallia in imposex-affected females *S. haemastoma*

Aphallia is reported for the first time in females *S. haemastoma*, aphallic females were observed only in the Western Mediterranean basin (Figure 3). Aphallic females were found in 7 out of 11 imposex-affected populations. Females without penis and with only a vas deferens were found in three Tunisian sites (Tabarca, 15.38% and El Houaria, 6.66%) and in one Algerian site (Oran, 12.5%) (Figure 3). While, females *S. haemastoma* with a vas deferens and a papilla were found in four stations, length of the papilla varied between 0.11 mm in Algiers (Algeria) and 0.33 mm in Korbous (Tunisia). Rates of females showing only a papilla varied between 5.88% and 50% in Korbous and Sidi Bechir, respectively.

Discussion

The present work highlights the occurrence of morphological malformations affecting the genital tract of males and imposex-affected females *S. haemastoma*. Malformations described in the literature are essentially related to the differentiation or the under-development of a part or a whole male genital tract in both sexes [22,26,40]. In Tunisia, penis malformations were first reported by Lahbib et al. [22] in males and imposex affected females *H. trunculus*, while Abidli et al. [26]

detected malformations affecting the penis and/or the vas deferens of two muricids (*H. trunculus* and *B. brandaris*) sampled from Bizerta Channel (Table 2). In *S. haemastoma* only imposex was signalled in specimens sampled from Bizerta Channel [11,38]. Penis excrescences observed in 3 males *S. haemastoma* were previously observed in 1 male *H. trunculus* collected from Bizerta Channel [22] and in three other gastropod species (*Hexaplex trunculus*, *Bolinus brandaris* and *Hinia reticulata*) collected from the Atlantic coast [24,27,29]. Aphallia and biphallia were the most common malformations recorded in the literature (Table 2). Whereas, the observation of male with abnormal shaped penis (Z-shape) is newly reported in *S. haemastoma* collected from an Atlantic shore (Morocco). Abnormal shaped penis was only reported by Lahbib et al. [30] in 1 Male *Cyclope neritea*. In contrast, aphallia was widely reported in previous studies (Table 2), in the present study aphallia was detected in *S. haemastoma* in both sexes. Aphallic females were more frequent than males, similar finding was recorded in *Nucella lapillus* [16]. Usually, aphallia is reported as cases of DS, in other cases it was considered a path of imposex development [18]. However, there is a doubt about the occurrence of Dumpton Syndrome in this population, since the rate of aphallic males was low. Indeed, Queiroz et al. [18] did not found any aphallic male, while they noted a very high rate of aphallic imposexed females (46.2%). Same finding was reported by Santos et al. in *Nucella lapillus* 0.1% of aphallic males against 23.8% of aphallic females. Quintela et al. [18] noted 7% of aphallic males and 31% imposexed females. That's why those authors recommended thus the use of the diameter of the vas deferens instead of the size of the penis in studying the Dumpton Syndrome. As for imposex development, the aphallic path observed here was not observed previously in *S. haemastoma* sampled from Tunisian coast [30,38]. This path was only observed in the Brazilian population [18].

The mechanism implicated in the development of penis malformations was widely discussed but still unknown; some authors supported the hypothesis linking genitalia malformations to the presence of oragnotin compounds in the aquatic environment. In this context, [41] observed penis malformations in juveniles *Buccinum undatum* exposed to a very high concentration of TBT (500 ng/L). Later, Lahbib et al. [22] supported this idea based on the fact that penis malformations were only found in highly TBT-polluted site. However, the low rates of penis malformations recorded in previous studies: (0.32%) in Portuguese *H. trunculus* [24], (1.34%) in Tunisian *H. trunculus* [26], (0.56%) in Tunisian *B. brandaris* [26], (1.69%) in Spanish *B. brandaris* [29], (0.2%) and (2.44%) in Brazilian *Leucozonia nassa* and *Leucozonia ocellata*, respectively [21]. Penis malformations reported in the present study were very low (0.86%) in Tunisian *S. haemastoma* against (2.38%) in Moroccan population. Factors other than tributyltin were probably implication in the development of those malformations. This assumption is supported by the observation of biphallid males among museum specimens of *H. trunculus* collected before the use of the biocide TBT (tributyltin) [7]. Generally, malformations are evidence of environmental problems generating negative impacts on live organisms [42,43], but they may also appear naturally. Indeed, further investigations are needed in order to elucidate the causal factor generating such malformations in marine gastropods.

Conclusion and Recommendations

The present work elucidates the occurrence of malformations affecting the genital tract of males and imposex-affected females *S. haemastoma*. The mechanism implicated in the development of penis malformations is supported by the hypothesis linking genitalia malformations to the presence of oragnotin compounds in the aquatic




Figure 3: Rates of aphally in females *Stramonita haemastoma*.

| Studies | Species | Locations | Penis malformation | | |
|-----------------------------|------------------------------|-----------------------------|--------------------|-----|----------------------------|
| | | | N | Nmm | Malformation |
| Present study | <i>Stramonita haemastoma</i> | Bizerta Channel | 1035 | 3 | Penis excrescences |
| | | -- | -- | 1 | Biphallid penis |
| | | -- | -- | 1 | Biphallid penis/undersized |
| | | -- | -- | 4 | Aphallic |
| | | Bouznika | 42 | 1 | Penis Z-shape |
| Sanchez-Marin et al. (2015) | <i>Nucella lapillus</i> | Portuguese coast | 1433 | 2 | Aphallic |
| | | -- | -- | - | Undersized penis |
| | | -- | -- | 6 | Biphallid penis |
| Abdili et al. (2009) | <i>Hexaplex trunculus</i> | Gulf of Tunis | 522 | 6 | Biphallid penis |
| | | -- | -- | 1 | Triphallic penis |
| | | <i>Bolinus brandaris</i> | Gulf of Tunis | 531 | 3 |
| Lahbib et al. (2008) | <i>Hexaplex trunculus</i> | Steg Tunis North Lake | 14 | 1 | Bifurcated tip |
| | | Small Gulf of Tunis | 48 | 1 | Biphallid penis |
| | | Bizerta Channel | 42 | 2 | Penis excrescences |
| Terlizzi et al. (1998) | <i>Hexaplex trunculus</i> | Italian coast | 600 | -- | Bifurcated penis |
| | | -- | -- | 1 | Aphallic |
| Terlizzi et al. (1999) | <i>Hexaplex trunculus</i> | Italian coast | 3000 | 4 | Aphallic |
| Barreiro et al. (1999) | <i>Nucella lapillus</i> | Spanish coast | 1602 | 39 | Aphallic |
| Vasconcelos et al. (2006) | <i>Hexaplex trunculus</i> | Ria Formosa | 621 | 2 | Penis excrescences |
| Ramon and Amor (2001) | <i>Bolinus brandaris</i> | Spanish coast | 59 | 1 | Penis excrescences |
| Stroben et al. (1992) | <i>Hinia reticulata</i> | Brittany and Normandy Coast | 2760 | 4 | Penis excrescences |
| Huet et al. (2008) | <i>Nucella lapillus</i> | French coast | 1847 | 19 | Aphallic |
| Costa et al. (2014) | <i>Leucozonia nassa</i> | Southeast Brazil | 493 | 1 | Biphallid penis |
| | | -- | 58 | 2 | Biphallid penis |
| Gibbs (1993) | <i>Nucella lapillus</i> | UK | -- | -- | Aphallic |
| | | -- | -- | -- | Undersized penis |

Table 2: Penis malformations in *Stramonita haemastoma* with comparison to data reported in the literature. N: number of specimens; Nmm: number of malformed males.

environment generating negative impacts on live organisms. Indeed, further investigations are needed in order to clarify the causal factor generating this malformation in marine gastropods.

References

- Axiak V, Micallef D, Muscat J, Vella A, Mintoff B (2003) Imposex as a biomonitoring tool for marine pollution by tributyltin: Some further observations. *Environ Int* 28: 743-749.
- Rilov G, Gasith A, Evans M, Benyahu Y (2000) Unregulated use of TBT based antifouling paints in Israel (Eastern Mediterranean): High contamination and imposex levels in two species of marine gastropods. *Mar Ecol Prog Ser* 192: 229-238.
- Bryan G, Gibbs P, Hummerstone L, Burt G (1986) The decline of the gastropod *Nucella lapillus* around South-West England: Evidence for the effect of tributyltin from antifouling paints. *J Mar Biol Assoc UK* 66: 611-640.
- Horiguchi T, Takiguchi N, Cho H, Kojima M, Kaya M, et al. (2000) Ovo-testis and disturbed reproductive cycle in the giant abalone, *Haliotis madaka*: possible linkage with organotin contamination in a site of population decline. *Mar Environ Res* 50: 223-229.
- Smith P (1996) Selective decline in imposex levels in the dogwhelk *Lepsiella scobina* following a ban on the use of TBT antifoulants in New Zealand. *Mar Poll Bull* 32: 362-365.
- Gibbs P, Bryan G, Pascoe P, Burt G (1987) The use of the dog-whelk, *Nucella lapillus*, as an indicator of tributyltin (TBT) contamination. *J Mar Biol Assoc UK* 67: 507-523.
- Garaventa F, Faimali M, Terlizzi A (2006) Imposex in pre-pollution times. Is TBT to blame? *Mar Pollut Bull* 52: 701-702.
- Shi H, Huang C, Zhu S, Yu X, Xie W (2005) Generalized system of imposex and reproductive failure in female gastropods of coastal waters of mainland China. *Mar Ecol Prog Ser* 204: 179-189.
- Titley-O'Nea LC, Munkittrick K, Macdonald B (2011) The effects of organotin on female gastropods. *J Environ Monit* 13: 2360-2388.
- Neves R, Echeverria C, Pessoa L, Paiva P, Paranhos R, et al. (2013) Factors influencing spatial patterns of molluscs in a eutrophic tropical bay. *J Mar Biol Assoc UK* 93: 577-589.
- Lahbib Y, Abidli S, Trigui El MN (2013a) TBT pollution in Tunisian coastal lagoons as indicated by imposex in *Hexaplex trunculus* (Gastropoda: Muricidae). *Transit Water Bull* 6: 17-24.
- Domínguez-Ojeda D, Patrón-Soberano O, Nieto-Navarro J, De Lourdes Robledo-Marengo M, Velázquez-Fernández J (2015) Imposex in *Plicopurpura pansa* (Neogastropoda: Thaididae) in Nayarit and Sinaloa, Mexico. *Rev Mex Biodivers* 86: 531-534.
- Gibbs PE (1993) A male genital defect in the dog-whelk, *Nucella lapillus* (Neogastropoda), favouring survival in a TBT-polluted area. *J Mar Biol Assoc UK* 667-678.
- Huet M, Le Goïc N, Gibbs P (2008) Appearance of a genetically-based pollution resistance in a marine gastropod, *Nucella lapillus*, in the southwest Brittany: A new case of Dumpton Syndrome. *J Mar Biol Assoc UK* 88: 1475-1479.
- Barreiro R, Quintela M, Ruiz J (1999) Aphally and imposex in *Nucella lapillus* from Galicia (NW Spain): Incidence, geographical distribution and consequences for the biomonitoring of TBT contamination. *Mar Ecol Prog Ser* 185: 229-238.
- Sánchez-Marín P, Oliveira IB, Sousa AC, Takahashi S, Tanabe S, et al. (2016) Evaluation of female aphally in imposex-affected populations of *Nucella lapillus* at the southernmost distributional limit of the species in Europe. *J Mollus Stud* 82: 144-153.
- Lima Afad, Castro ÍBD, Rocha-Barreira CDA (2006) Imposex induction in *Stramonita haemastoma floridana* (Conrad, 1837) (Mollusca: Gastropoda: Muricidae) submitted to an organotin-contaminated diet. *Braz J Oceanogr* 54: 85-90.
- Queiroz LR, Castro ÍBD, Rocha-Barreira CDA (2007) New imposex development index (IDI) for *Stramonita haemastoma* (Mollusca: Muricidae): A transplantation experiment in the Brazilian Northeast.
- Cardoso R, Caetano C, Cabrini T (2009) Biphallia in imposexed females of marine gastropods: New record for *Nassarius vibex* from Brazil *Braz J Biol* 69: 223-224.
- Castillo V, Brown D (2012) One Case of Triphallia in the Marine Snail

- Echinolittorina peruviana (Caenogastropoda: Littorinidae). Int J Morph 30: 791-796.
21. Costa M, Mardegan B, Zamprogno G, Pedrucci F, De Morais L, et al. (2014) Penis Malformations in *Leucozonia nassa* (Gmelin, 1792) and *Leucozonia ocellata* (Gmelin, 1791) in a TBT Contaminated Region from Brazil. Aquatic Science and Technology (AST) 2: 52-66.
 22. Lahbib Y, Abidli S, Trigui El MN (2008) Imposex level and penis malformation in *Hexaplex trunculus* from the Tunisian coast. Am Malacol Bull 24: 79-89.
 23. Meirelles C, Castro I, Pinheiro J (2007) A first record of biphallia in imposexed male of *Leucozonia nassa* (Caenogastropoda: Fascioliidae). J Mar Biol Assoc UK 1-3.
 24. Vasconcelos P, Gaspar M, Castro M (2006) Imposex in *Hexaplex* (*Trunculariopsis*) *trunculus* (Gastropoda: Muricidae) from the Ria Formosa lagoon (Algarve coast-southern Portugal). Mar Pollut Bull 52: 337-341.
 25. Gibbs P, Bryan G, Pascoe P, Burt G (1990) Reproductive abnormalities in female *Ocenebra erinacea* (Gastropoda) resulting from tributyltin-induced imposex. J Mar Biol Assoc UK 70: 639-656.
 26. Abidli S, Lahbib Y, Trigui El Menif N (2009) Imposex and genital tract malformations in *Hexaplex trunculus* and *Bolinus brandaris* collected in the Gulf of Tunis. B Mar Sci 85: 11-25.
 27. Stroben E, Oehlmann J, Fioroni P (1992) The morphological expression of imposex in *Hinia reticulata* (Gastropoda: Buccinidae): A potential indicator of tributyltin pollution. Mar Biol 113: 625-636.
 28. Terlizzi A, Geraci S, Minganti V (1998) Tributyltin (TBT) pollution in the coastal waters of Italy as indicated by imposex in *Hexaplex trunculus* (Gastropoda, Muricidae). Mar Poll Bull 36: 749-752.
 29. Ramon M, Amor M (2001) Increasing imposex in populations of *Bolinus brandaris* (Gastropoda: Muricidae) in the northwestern Mediterranean. Mar Environ Res 52: 463-475.
 30. Lahbib Y, Abidli S, Chiffolleau J, Averty B, Trigui El MN (2010) Imposex and butyltin concentrations in snails from the lagoon of Bizerte (Northern Tunisia). Mar Biol Res 6: 600-607.
 31. Lahbib Y, Abidli S, El Menif N (2011) Spawning and intracapsular development of *Stramonita haemastoma haemastoma* (Gastropoda: Muricidae) collected in northern Tunisia. Mar Biol Res 7: 719-726.
 32. Barhoumi B (2014) Biosurveillance de la pollution de la lagune de Bizerte (Tunisie) par l'analyse comparée des niveaux de contamination et de l'écotoxicité des sédiments et du biote. Dissertation, Université Michel de Montaigne-Bordeaux III, France.
 33. Hamdi H, Jedidi N, Yoshida M, Mosbahi M, Ghrabi A (2002) Some Physio-Chemical Properties of Lake Bizerte Sediments. RPP-SEPMCL Initial Report 49-54.
 34. Louiz I, Kinani S, Gouze M, Ben-Attia M, Menif D, et al. (2008) Monitoring of dioxin-like, estrogenic and anti-androgenic activities in sediments of the Bizerte lagoon (Tunisia) by means of *in vitro* cell-based bioassays: Contribution of low concentrations of polynuclear aromatic hydrocarbons (PAHs). Sci Total Environ 402: 318-329.
 35. Barhoumi B, Lemenach K, Devier M, Ben Ameer W, Etcheber H, et al. (2014) Polycyclic aromatic hydrocarbons (PAHs) in surface sediments from the Bizerte Lagoon, Tunisia: Levels, sources, and toxicological significance. Environ Monit and Assess 186: 2653-2669.
 36. Hassine S, Ameer W, Gandoura N, Driss M (2012) Determination of chlorinated pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers in human milk from Bizerte (Tunisia) in 2010. Chemosphere 89: 369-377.
 37. Ameer W, Trabelsi S, Bedoui B, Driss M (2011) Polychlorinated biphenyls in sediments from Ghar El Melh lagoon, Tunisia. B Environ Contam Tox 86: 539-544.
 38. El Ayari T, Abidli S, Lahbib Y, Rodríguez GP, García AJ, et al. (2015) The effect of size and epibiotic barnacles on imposex in *Stramonita haemastoma* collected from the northern coast of Tunisia. Mar Biol Res J 11: 313-320.
 39. Lahbib Y, Abidli S, Trigui El MN (2009) Relative growth and reproduction in Tunisian populations of *Hexaplex trunculus* with contrasting imposex levels. J Shellfish Res 28: 891-898.
 40. Gibbs P (1996) Oviduct malformation as a sterilising effect of tributyltin (TBT)-induced imposex in *Ocenebra erinacea* (Gastropoda: Muricidae). J Mollus Stud 62: 403-413.
 41. Mensink B, Kralt A, Vethaak D, Ten Hallers-Tjabbes C, Koeman J, et al. (2002) Imposex induction in laboratory reared juvenile *Buccinum undatum* by tributyltin (TBT). Environ Toxicol Pharmacol 11: 49-65.
 42. Rodríguez J, Tueros I, Borja Á, Franco J, Alonso J, et al. (2009) Butyltin compounds, sterility and imposex assessment in *Nassarius reticulatus* (Linnaeus, 1758), prior to the 2008 European ban on TBT antifouling paints, within Basque ports and along coastal areas. Cont Shelf Res 29: 1165-1173.
 43. Schulte-Oehlmann U, Oehlmann J, Fioroni P, Bauer B (1997) Imposex and reproductive failure in *Hydrobia ulvae* (Gastropoda: Prosobranchia). Mar Biol 128: 257-266.

Citation: El Ayari T, Mleiki A, El Menif NT (2017) Genitalia Malformations in *Stramonita haemastoma* (Gastropoda: Muricidae) from Atlantic and Mediterranean Coast. J Aquac Res Development 8: 493. doi: [10.4172/2155-9546.1000493](https://doi.org/10.4172/2155-9546.1000493)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>