

Genome silencing, Cell Division and Phytohormone Biosynthesis in Winter Wheat (a possible relationship)

Fedora Sutton*

Department of Plant Sciences, South Dakota State University, USA

It is not clear to what extent gene silencing plays a role in polyploid gene expression nor is it clear whether there is a mechanism to silence similar genes or all genes involved in a particular pathway. It is known that epigenetic marks such as methylation and deacetylation can influence gene expression resulting in silencing of groups of genes.

One group of genes for which deacetylation has been reported to play a key role is the set of ribosomal RNA (rRNA) genes. These genes are expressed at the Nucleolar Organizer Regions (NORs). Of the *Arabidopsis* histone deacetylases (HDACs), HDAC6 has been reported as playing a significant role in silencing of the rRNA genes by modifications of those genes [1]. In Figure 1, HDAC6 can be seen localized to the nucleolus where rRNA genes are normally transcribed. In this case HDAC6 switched off the rRNA genes. Early et al. [1] have also described a model that predicts the key events that allow switching from active transcription to repression of transcription. These key events involve de novo cytosine methylation and histone deacetylation.

Rather than turning off whole sets of genes, epigenetic marks can be limited to regions of one sister chromatid and not another. Such differential epigenetic marks on sister chromatids can lead to the generation of daughter cells in which one cell expresses particular genes while in the other daughter cell the genes are silenced [2]. Assuming no other controls are placed on the expression or lack of expression of these genes, it means that there will be cells in the same tissue that are expressing a particular protein and other cells that do not express that protein. As a result the cell fates are different. An additional aspect of this control is the role of stimuli on influencing the epigenetic changes leading to differential gene expression.

At the transcriptional level, there are many studies utilizing DNA microarrays and next generation RNA sequencing to identify gene networks responsive to various stimuli. This is a very complex and vast area of study. For now, I will focus the discussion on plants and response to stimuli to withstand abiotic stress. Our studies have focused




Figure 1: *Arabidopsis thaliana* nucleus. DNA is stained with DAPI (blue). HDAC6 which is Flag tagged (HDAC-Flag) is immunolocalized with anti-Flag monoclonal primary antibody to the nucleolus (red signal) where rRNA genes are located [1].

on deciphering the mechanism(s) that allow plants, particularly cereals, to withstand freezing conditions. For most organisms, prior exposure to an attenuated version of the stress is known to prepare the organism to withstand the stress (acclimation). In studying freeze survival, many studies have focused on what occurs during that attenuation period termed cold acclimation. In nature, cold acclimation occurs during the autumn and can be mimicked in the laboratory with low non-freezing temperatures under dehydration conditions and low light intensity [2].

Triticum aestivum is a hexaploid (AA, BB and DD). Therefore in our system we have to be aware as to whether we are observing gene expression from all three genomes or from a subset. This is important for many reasons including understanding dosage compensation in wheat. Firstly, I will describe the experimental system and then we will return to the role of the genomes in cold acclimation.

Having an experimental system that is very closely related and differing significantly only in survival of the stress is very valuable. We have such a system. Utilizing winter wheat lines generated by azide mutagenesis and varying in freeze survival and not vernalization, we have been able to compare expression of the cold acclimation regulon consisting of transcription factors known to map to the long arm of chromosome 5, the frost resistant (Fr) loci [3,4]. These transcription factors belong to the group known as CBF/DREB based on identification of the binding of those proteins to C-repeat/dehydration-responsive elements (CRT/DRE) [5,6]. We identified two of these genes *Cbf-12* and *Cbf-14*; from the *Cbf* central cluster of the Fr loci [7] as associated with the difference in freeze survival between the lines. *Cbfs-3,-5,-6* are significantly differentially regulated between FR and FS (Figure 2). However, these *Cbf* genes are not located within the central cluster.

As stated earlier, it is of great value when studying gene expression in a polyploid to be able to ascertain which if any of the genomes is a key player. As seen in figure 2, *Cbf-D22* is silenced in response to cold acclimation in both lines. However, *Cbf-A22* and *Cbf-B22* are upregulated almost to the same extent in response to cold acclimation in both lines. With respect to *Cbf-22*, these results suggest silencing to achieve dosage compensation to limit its expression. How and why the *Cbf-D22* is silenced is not known.


The results depicted in Figure 2, also reveal that silencing is not limited to one genome. Although *Cbf-B22* is activated in both lines another gene on the B genome *Cbf-B10* is silenced in both lines.

*Corresponding author: Fedora Sutton, Department of Plant Sciences, South Dakota State University, USA, Tel: 605-688-6286; E-mail: Fedora.Sutton@sdstate.edu

Received October 29, 2013; Accepted October 30, 2013; Published November 06, 2013

Citation: Sutton F (2013) Genome silencing, Cell Division and Phytohormone Biosynthesis in Winter Wheat (a possible relationship). J Plant Biochem Physiol 1: e115. doi:10.4172/2329-9029.1000e115

Copyright: © 2013 Sutton F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Information garnered from the response of the genomes and the levels of transcription for individual genes is very important. However, from a system's point of view, it is important to move beyond the single genes and define the pathways that must function together to manifest physiological changes. To that end, we utilized out transcriptomics datasets and performed pathway analyses [8] with the Pathway Studio software [9]. The pathway "neighbors of DREB2A," was up regulated (positive) with the highest median fold change in response to cold acclimation in the line with greater freeze survival [8], and as expected during cold acclimation and for a non-photosynthesizing tissue (the crown, the most freeze resistant part of the plant) four "photosynthetic-associated" pathways reflected reduction in expression (negative) in response to cold acclimation. Similar negative median fold changes were obtained for the E2F pathway which controls cell division and the "Jasmonic acid biosynthesis pathway". Of significance is the finding that the degree of shut down of these pathways in response to cold acclimation was greater for the freeze resistant line than that for the freeze susceptible line.

Critical to the understanding of response to acclimation in preparation to withstand stresses, is the identification of the final gene products. These are the metabolites and the proteins. From the pathway analyses, we know that absence or reduction of Jasmonic acid biosynthesis is important to maintaining the "relative dormant" state of the crown tissue. It is well known that ABA is important for plants to withstand various stresses. Our studies along with the well described

Citation: Sutton F (2013) Genome silencing, Cell Division and Phytohormone Biosynthesis in Winter Wheat (a possible relationship). J Plant Biochem Physiol 1: e115. doi:10.4172/2329-9029.1000e115

ABA dependent systems suggest an ABA/Jasmonic acid antagonistic system to maintain the meristematic crown tissue in a dormant state. Lack of cell division as reflected by the shutdown of the E2F pathway and the absence of growth during cold acclimation [3] leads to the design of a model that predicts that deacetylation and methylation events may occur to silence genes involved in cell division and biosynthetic pathways (pathways of Jasmonic Acid but not that of ABA).

References

1. Earley K, Lawrence RJ, Pontes O, Reuther R, Encisco AJ, et al. (2006) Erasure of histone acetylation by Arabidopsis HDA6 mediates large-scale gene silencing in nucleolar dominance. *Genes Dev.* 20:1283-1293.
2. Lansdorp PM, Falconer E, Tao J, BrindAmour J, Naumann U (2012) Epigenetic differences between sister chromatids? *Ann NY Acad Sci* 1266: 1-6.
3. Kenefick DG, Koepke JA, Sutton F (2002) Plant water uptake by hard red winter wheat (*Triticum aestivum* L.) genotypes at 2 degrees C and low light intensity. *BMC Plant Biol* 2: 8.
4. Sutton F, Chen D, Ge X, Kenefick D (2009) Cbf genes of the Fr-A2 allele are differentially regulated between long-term cold acclimated crown tissue of freeze-resistant and - susceptible, winter wheat mutant lines. *BMC Plant Biology* 9:34.
5. Yamaguchi-Shinozaki K, Shinozaki K (1994) A novel cis-acting element in an Arabidopsis gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* 6: 251-264.
6. Baker SS, Wilhelm KS, Thomashow MF (1994) The 5'-region of Arabidopsis thaliana *cor15a* has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol Biol* 24: 701-713.
7. Knox AK, Li C, Vágújfalvi A, Galiba G, Stockinger EJ, et al. (2008) Identification of candidate CBF genes for the frost tolerance locus Fr-Am2 in *Triticum monococcum*. *Plant Mol Biol* 67: 257-270.
8. Karki A, Horvath DP, Sutton F (2013) Induction of DREB2A pathway with repression of E2F, jasmonic acid biosynthetic and photosynthesis pathways in cold acclimation-specific freeze-resistant wheat crown. *Funct Integr Genomics* 13: 57-65.
9. Bogner V, Leidel BA, Kanz KG, Mutschler W, Neugebauer EA, Biberthaler P (2011) Pathway analysis in microarray data: a comparison of two different pathway analysis devices in the same data set. *Shock* 35: 245-251.

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit/>