

Gravity and Electromagnetism

Paul TE Cusack*

Independent Researcher, BSc E, DULE, 1641 Sandy Point Rd, Saint John, NB, Canada E2K 5E8, Canada

Abstract

Here is a paper that uses the theory of electromagnetism to model gravity and the universe. The standard magnetic flux, capacitance, conductivity, dielectrics are considered. The mathematics includes the golden mean parabola, Euler's Identity, Matrices, Eigenvectors, and Differential equations.

Keywords: Gravity; Magnetic flux; Capacitance dielectric; Permeability; Mass gap

Introduction

Here we provide a solution to the physical universe that shows that it can be modelled as an electromagnetic flux using well know electrical engineering mathematics. The Then gravity is modelled as an eigen vector. The standard Golden Mean parabola and the derivative equals the function is used once again to help solve the problem of why gravity exists. We begin with magnetism and flux.

Magnetic flux density

$$L = \mu_0 N^2 A / l = 0.666 = 1.15 * n^2 * 0.1646 / 0.8415 = \sqrt{3}$$

$$B = \mu_0 N i / l = 1.15 * \sqrt{3} * 1.3 / 0.8415 = 3.05 = c = M\rho$$

$$1/L = B / [\partial E / \partial t] \partial M / \partial t$$

$$\partial E / \partial t = BL (\partial M / \partial t)$$

$$2t - 1 = BL (2)$$

$$t = 5/2 = 2.5b = T = \text{PERIOD}$$

$$x^3 - x - 1 = 2^2 - 2 - 1 = 1 = E.$$

Conductivity

$$R = \rho * l / A$$

$$1.618 = \rho * 0.8415 / 0.1646$$

$$\rho = 0.316$$

$$\sigma = 1/\rho = 3.14 = \pi = E$$

$$d = 0.8415$$

$$d/2 = 0.42 = \pi - e$$

$$\ln e = 1$$

$$\pi - \ln e = 2.14$$

$$2 * \partial M / \partial t = 1 * 2 = 2.$$

Universal capacitor resistance

$$Q = 1 / [\omega C R c]$$

$$1.3 = 1 / [0.8415 * 1.48 * R c]$$

$$R c = 1.619 \text{ cf } 1.618.$$

Euler's identity

$$A = E * t = E^2 = t^2 = 1$$

$$i * \int \sin \theta - \int \cos \theta = A = 1 = -e^\pi$$

$$i^* \cos \pi - \sin \pi = 1$$

$$\theta = \pi = 180^\circ$$

This is half a cycle.

$$\text{Area Under cosine} = 0$$

$$\text{Area Under sine} = \pi = E.$$

Capacitance continued

$$C = q/V$$

$$1.5 = q/0.86$$

$$q = 1.3 = \partial i / \partial t \text{ where } t = 1$$

$$\text{Now, } C = \epsilon r = \epsilon_0 A/d$$

$$1.5 = 0.86 (8.854)(A/d)$$

$$A/d = \sqrt{3}/8.854$$

$$d = 0.86 \text{ because } \sin \theta = 0.86/1$$

$$A = 0.1956 (0.8415)$$

$$A = 0.1646$$

$$\sqrt{A} = 0.4057 = t$$

$$d_0 = 1/2$$

$$0.4057/2 = 0.2028 = Y = E \text{ cf } Y = 0.203 \text{ [Dampened Cosine Curve]}$$

$$t = s - c$$

$$t = s - \partial s / \partial t$$

Integrate (Time is a Constant with respect to t)

$$1 = s^2/2 - s$$

$$s^2 - s - 1 = 0$$

$$x^2 - x - 1 = 0$$

*Corresponding author: Cusack P, Independent Researcher, BSc E, DULE, 1641 Sandy Point Rd, Saint John, NB, Canada E2K 5E8, Canada, Tel: (506) 214-3313; E-mail: st-michael@hotmail.com

Received December 10, 2016; Accepted March 18, 2017; Published March 28, 2017

Citation: Cusack PTE (2017) Gravity and Electromagnetism. Fluid Mech Open Acc 4: 155. doi: 10.4172/2476-2296.1000155

Copyright: © 2017 Cusack PTE. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$t=s-c$
 $t=s-\partial s/\partial t$
 Integrate: $t^2/2=s^2/2-s$
 $t=1$
 $t^2/2=s^2/2-s$
 $2=s^2-s$
 $s^2-s-2=0$
 Quadratic: $(s-2)(s+1)=0$.
 $s=2, -1$
 $-\partial s/\partial t=-1$
 $\partial s/\partial t=1$
 $\partial t/\partial t=1$
 $\partial s/\partial t=s$
 $y=y'$
 The Derivative Equals the Function
 $d=v_i t+1/2 at^2$ (same equation)
 $d=s_i+s/2$
 $d=d/2+d/2$
 $s_i=d/2$
 The initial distance= $1/2$
 $\partial E/\partial t=2t-1$
 $\partial E/\partial t=2(1/2)-1=0$ (Minimum)
 $F=v$
 $M=R$
 $V=L \partial i/\partial t F=V=v$
 $0.86=0.666 \partial i/d\partial t$
 $3/2 * 0.666=\partial i/\partial t$
 $1.3=\partial i/\partial t$
 $1-\partial i/\partial t=1-1.3=-0.3=-c/10=-\partial s/\partial t$
 current i =speed of light c .
 $s=t-i 0.1334=t-1.3$
 $t=1.16666$
 $E=1/t=1/1.166=0.86$
 $1-\partial i/\partial t=\partial s/\partial t$
 $t-i=s$
 $t-c=s$
 $s=t-c$
 $t=s-c$
 time=distance-speed of light
 distance=time-speed of light/10

distance= $0.4+0.3=0.7$
 $1/s=1.4285$
 $s=0.85$
 $V=L \partial i/\partial t$
 But $F=G M_1 M_2/R^2$
 $\partial i/\partial t=M_1 M_2/R^2$
 when M_1 is very large and M_2 is very small, $=1/R^2$.
 $\int \partial i/\partial t=i=\int 1/R^2=-1/R$
 $i=-1/R$
 $F^2/2=-1/R$
 $1/2R^2 * R=-1$
 $1/2 vM=K.E.=1$
 Therefore, $F=v R=M$.

Dielectric Constant and Permeability

$L'C'=\epsilon \mu$
 $(0.666)(1.5)=1 1=\epsilon(15\%)$
 $\epsilon=1/1.15=0.86$
 Capitanace= 1.48
 $C*M=1.5*2=3=Mp$
 Mass Gap
 $L=1/1.5=0.6666=G$.

So the inductance of our universe is G . The energy of the mass with velocity is converted to a Gravitational field. $G=0.666$ is the result. Gravity is modelled like a magnetic field. Like magnets, matter is attracted to matter. We call it gravity. The Earth has a magnetic field because it has matter or mass. The magnetic field is proportional to $g=9.81$.

$5.972/9.81=0.609=1/1.642$
 (The Mass of the Earth is closer to 6.060108)
 Better still
 $1/1.618=0.618 0.618*9.806=6.06=2.02*3=Y*c=Ec$
 But $E=Mc^2$
 So $Ec=Mc^3=g/M$
 $M^2c^3=g$
 (This is why $g=9.81m/s$)
 $g=M^2Mp^3$
 $g=M^2 * (\partial M/\partial t)^3$
 $g=E * Mc$
 $a=E * M \partial s/\partial t$
 $a=E*M v$
 $a/v=1=E*M$

$$E \cdot M = 1 = t$$

$$M = t^2.$$

Conservation of Momentum

$$Mv_1 = Mv_2$$

$$v = \sin \theta$$

$$M v_1 = M \sin 1$$

$$\partial M / \partial t \cdot v_1 = M \cdot 0.8415$$

$$2 \cdot v_1 = 0.8415 M$$

$$v_1 = 0.8415 / 2 \cdot M$$

$$= 0.4207 M$$

$$= \text{cuz } M$$

$$0.8415 / \text{cuz} = M$$

$$M = 2$$

$$E = Mc^2$$

$$\pi = 2 \cdot c^2$$

$$c = \sqrt{\pi / 2}$$

$$c = 1.2533 = \text{MINIMUM OF THE ENERGY PARABOLA}$$

$$2t - 1 = 0$$

$$t = 1/2$$

$$P = Mv = 2 \cdot (0.866) = 1.73 = \sqrt{3}$$

$$K.E. = 1/2 Mv^2$$

$$1 = 1/2 M \cdot (0.866)^2$$

$$M = 2.666$$

$$M = -\partial M / \partial t + G$$

And

$$M = F = 26.666$$

Now,

$$K.E. = 1/2 Mv^2$$

$$Pi = 1/2 (2) v^2$$

$$v^2 = \pi$$

$$v = \sqrt{\pi} = 1.77$$

$$\text{Bernoulli: } mgh + 1/2 mv^2 + p$$

$$0.666bh + 1/2 (1.77)^2 + 0$$

$$1/h = 0.4240 = \text{cuz}$$

$$E/h = \text{cuz Hooke's Law.}$$

$$\lim_{y \rightarrow 0} \int x \, dx = y$$

$$\lim_{y \rightarrow 0} \int x \, dy = y$$

$$e^x = y$$

$$y = y' = e^x$$

$$y = y' = y'' = y''' \dots = e^x$$

$$\lim_{y \rightarrow \infty} y = e^x$$

$$e^{\text{Infinity}}$$

$$x = \text{Ln } \infty$$

$$y = y' = y'' = \dots = e^x = \infty$$

The universe is robust [1-4].

Similar Matrices

$$B = S^{-1} A S$$

$$\Omega A = \sqrt{(3/2)}$$

$$\sqrt{3} / \sqrt{4} = [2 \ 0 \ 0 \ 2] = [\sqrt{3} \ v_0 \ 0 \ \sqrt{3}] [|A| = 3$$

$$\det |A^{-1}| = 1 / [\det |A|].$$

$$\begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{4} \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} =$$

$$|D| = 1/4$$

$$\text{So, eigenvalues } \begin{bmatrix} \lambda - 2 & 0 \\ 0 & \lambda - 2 \end{bmatrix} \lambda d = x$$

$$x^2 - x + 1/4 = 4$$

Quadratic (Note the $\sqrt{(-1)} = 0.618$)

$$\sqrt{3} = |A|, 0.7360 = \sqrt{0.858} = \sqrt{\Omega A}.$$

Mass Gap

$$E \cdot t \cdot x \cdot s = E/t$$

$$t \cdot E \cdot t \cdot s / E = \sqrt{3} / \sqrt{4}$$

$$t^2 \cdot s = \sqrt{3/2}$$

$$(t^2 \cdot s)^2 = (\sqrt{3/2})^2$$

$$t^4 \cdot s^2 = 3/E$$

$$E \cdot t^4 \cdot s^2 = c$$

$$\text{But } E = Mc^2$$

$$M = 1 / [E \cdot t^4 \cdot s^2] = 1 / [\sqrt{3} \cdot t^2 \cdot s] = 1.5 \text{ MASS GAP}$$

$$\text{Aside: } E \cdot t \cdot x \cdot s / E / t = t^2 \cdot s = (0.4083)(0.866) = 0.1444 = 0.856$$

C. G. E. = Cusack Gravitational Equation

$$1/G = M \cdot \rho / E \cdot \rho + \partial M / \partial t$$

The mass and energy density is wrt time, not space.

$$\text{So, } \partial^2 E / \partial t^2 = dE/dt / dM/dt + 1 / \partial M / \partial t$$

$$(\partial E / \partial t)^2 = [\partial E / \partial t + 1] / \partial M / \partial t$$

$$x^2 \partial M / \partial t - x - 1 = 0$$

$$\partial M / \partial t = 2$$

$$[\partial M / \partial t] / 2 = 1$$

$$E = Mc^2$$

$$\text{Mass} = T. E = P.E. + K. E. + 18 = 2(3^2)$$

$$(0.618)^3 = 0.2303$$

$$1/(0.618^3)=4.236=cuz *10$$

$$cuz=\pi-e.$$

Eigenvalues and Eigenvectors

Eigenvalue of the

$$C \cdot U \cdot E = \det \begin{bmatrix} 2-\lambda & 0 \\ 0 & 2-\lambda \end{bmatrix}$$

$$4-4\lambda+\lambda^2=0$$

$$x^2-4x-4=0$$

$$(x-2)(x-2)=0$$

$$x=2=\lambda$$

$$A=[2 \ 0, \ 0 \ 2]$$

$$|A|=4=E\rho$$

$$\det \begin{bmatrix} G & 0 \\ 0 & G \end{bmatrix} = G^2 = 4$$

$$\partial^2 E/\partial t^2 = E \rho$$

$$\partial E^2/\partial t^2 - E \cdot E \rho = 0 = \text{Ln } t$$

$$t=1 \text{ or } x=1 \text{ radian}$$

$$\text{Now, going back: } [2 \ 0, \ 0 \ 2][x \ y] = [\lambda \ x \ y]$$

$$[G \ 0, \ 0 \ G] [t \ E] = G [t \ E]$$

$$\text{Multiply by the diagonal of the unit cube } \sqrt{3}/G = 1.73/2 = 0.86$$

$$0.86 [2 \ 0 \ 0, \ 0 \ 2 \ 0, \ 0 \ 0 \ 2] [1 \ 1 \ 1] = G [t \ s \ E]$$

$$\sqrt{3} = 2t$$

$$t = \sqrt{3}/2 = 0.86$$

$$s = \sqrt{3}/2 = 0.86$$

$$E = \sqrt{3}/2 = 0.86$$

$$\partial^2 E/\partial t^2 - E = \text{Ln } t$$

$$2-1 = \text{Ln } t$$

$$e^2/e^1 = t$$

$$t = e^t$$

$$\text{Ln } t = t$$

$$\text{Ln } t = e^t$$

Derivative

$$1/t = e^t$$

So

$$\text{Ln } t = 1/t \text{ Or the derivative} = \text{the function}$$

$$y = y'$$

$$M = mgh + 1/2 mv^2$$

$$1 = gh + 1/2 v^2$$

$$\text{If } v = a$$

$$1/2 a^2 + 2a - 2 = 0$$

$$(a-1)(a-1) = 0$$

$$a = 1$$

$$M = P.E. + K.E.$$

$$M = mgh + 1/2 mv^2$$

$$v = 1 = a$$

$$\text{But } a = E = 2G$$

$$g = 1/2$$

$$2 = 2(1/2)(1) + 1/2(2)(1^2) \quad 2 = 1 + 1.$$

$$G = 2$$

$$\partial^2 E/\partial t^2 = 2 = \partial M/\partial t$$

$$\int \partial^2 E/\partial t^2 = M$$

$$\partial E/\partial t = M$$

$$\int \partial E/\partial t = \int M$$

$$E = M^2/2$$

$$E = 1/2 bM^2$$

$$1/2 M^2 = Mc^2$$

$$1/2 M = c^2$$

$$M = 2c^2 = 18 = P.E. + K.E.$$

$$1 - 0.1415 = 0.8575 = 0.$$

$$y'' + y' + y = 0$$

$$\partial^2 E/\partial t^2 + y' + E = \text{Ln } t$$

$$y' = \text{Ln } t$$

$$y = 1/t$$

$$E = 1/t$$

$$d_2 E/dt_2 (\text{Ln } t)^{2/2}$$

$$dE_2/dt_2 = y''.$$

So,

$$(\text{Ln } t)^2/2 + \text{Ln } t + 1/t = 0$$

When $a = v$, this is

$$1/2 a^2 + a + v = 0$$

$$a^2 + 2a + 2v = 0$$

$$(y'')^2 + 2y'' = 2y' = 0$$

$$G^2/2 + 2G + 2(E^1) = 0$$

$$2(2G-1) = -G^2/2 + 2G$$

$$4G-2 = G^2/2 + 2G$$

$$8G-4-4G = G^2$$

$$4G-4 = G^2$$

$$G^2-4G+4 = 0$$

$$(G-2)(G-2) = 0$$

$$(G-2)^2 = 0$$

$$G=2=\partial^2 E/\partial t^2.$$

Second Order Liners Equations

$$ay''+by'+cy=0$$

$$\partial^2 E/\partial t^2 - (1) E = \ln t$$

$$by' = \ln t \quad 0 = \ln t$$

$$t=1$$

$$ar^2+br+c=0$$

$$(1)r^2=0r+(-1)=0$$

$$r^2=1$$

$$r=\pm 1$$

$$(1)(1)=0(1)=(-1)=0$$

$$0=0 \text{ true}$$

The Material Universe Exists Where the Gravity Eigenvector.

$$\text{Eigenvector} = 1/1.5 = 0.667 \text{ i.e.}$$

$$1/(3/2) G = E/(M\rho \partial M/\partial t)$$

$$\partial^2 E/\partial t^2 = E/[M \rho /(\partial M/\partial t)]$$

$$\text{where } E = x^2 - x - 1$$

Why does E=1?

$$2x-1 = \int x^2 - x - 1$$

$$2x-1 = x^3/3 - x^2/2 - x - 1 - C$$

$$C = -1 \text{ @ } x=0$$

$$E = (-1)^2 - (-1) - 1$$

$$E = 1$$

$$E = x^2 - x - 1 = \int E$$

$$E = E'$$

$$x^2 - 1 = x^3/3 - 1.5x^2 - x - 1 = 3/2$$

$$x^3 - 2.5x^2 - x - 0 = 1.5$$

$$x^3 - 2.5x^2 - x - 1.5 = 0$$

$$x = 3 = M \rho = c$$

$$x = t$$

This is the solution to our physical universe.

Conclusion

We see that the physical universe can be thought of as an electrical flux and gravity as an eigenvector.

References

1. Disaster M (2010) undergrad Electricity & Magnetism (E&M) Text Book, Physics Forums and Missions, USA.
2. Tatum JB (2017) Electricity and Magnetism, Jason Stumpf, ITT Technical Institute-Fort Lauderdale Glen Allen, Virginia.
3. Chakraborty P (2016) Electricity and Magnetism. New age International, USA.
4. Cusack P (2016) Astro-Theology, Cusacks Universe. J Phys Math 7: 174.

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>