

5-HT_{5A} Receptors during Ontogeny

Berumen LC and García AG*

Facultad de Química, Universidad Autónoma de Querétaro, Centro Universitario, Querétaro 76010, México

*Corresponding author: Garcia AG, Faculty of Chemistry, Autonomous University of Querétaro, Cerro de las Campanas, Querétaro, 76010, Mexico, Tel: 52(442)1073116; Fax: 52(442)1921302; E-mail: leguga@email.com

Rec Date: January 29, 2014; Acc Date: February 18, 2014; Pub Date: February 28, 2014

Copyright: ©2014 Garcia AG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Serotonin is one of several neurotransmitters that are small molecules acting in developing organisms also as morphogens and growth factors; small metabolites are often regulatory signals to control gene expression [1,2]. The action of these molecules is led by the activation of their receptors but it might be considered also that some of them are biosynthesized from amino acids and can be reactive species, participating in other ways in the metabolism and signaling pathways, as serotonylation for example [3].

They act specifically through the activation of its own receptors but they can also be modulators of other signaling pathways; serotonin is an allosteric modulator of nicotinic receptor for example [4].

Our group has been working in the expression of the receptors during ontogeny for glycine, GABA, purinergic signals and, particularly, serotonin [5-9].

Our group had reported some findings of 5-HT_{5A} receptor during ontogeny of the rat as well as in the chicken [10]. Serotonin signaling during neurulation can be of particular interest, considering that neural tube defects are a frequent type of congenital malformations; the exposure of mouse embryos to 5-HT₂ ligands results in craniofacial and cardiac malformations. Trophic actions of 5-HT_{2B} is well recognized, and the participation of other serotonin receptors may be involved in a fine-tuning action for this molecule. 5-HT_{5A} was found to be expressed differentially in somite's, caudal region and encephalon, suggesting a role for this subtype of receptor in the development of several systems, including the central nervous system [10].

In the rat we have found changes in the expression of 5-HT_{5A} receptor particularly in hippocampus, which is one of the well-known areas that have adult neurogenesis. It is differentially expressed compared to 5-HT_{2C} and 5-HT₇ [9].

In conclusion, members of the 5-HT receptors, especially 5-HT_{5A} are expressed during early development, suggesting a possible role during morphogenesis.

References

1. Berumen LC, Rodríguez A, Mileli R, García-Alcocer G (2012) Serotonin receptors in hippocampus. *ScientificWorldJournal* 2012: 823493.
2. Buznikov GA, Lambert HW, Lauder JM (2001) Serotonin and serotonin-like substances as regulators of early embryogenesis and morphogenesis. *Cell Tissue Res* 305: 177-186.
3. Walther DJ, Peter JU, Winter S, Hölte M, Paulmann N, et al. (2003) Serotonylation of small GTPases is a signal transduction pathway that triggers platelet alpha-granule release. *Cell* 115: 851-862.
4. Pandya AA, Yakel JL (2013) Activation of the $\alpha 7$ nicotinic ACh receptor induces angiogenic effects in rats which is blocked by a 5-HT_{5A} receptor antagonist. *Neuropharmacology* 70: 35-42.
5. García-Alcocer G, Mejía C, Berumen LC, Mileli R, Martínez-Torres A (2008) Developmental expression of glycine receptor subunits in rat cerebellum. *Int J Dev Neurosci* 26: 319-322.
6. Mejía C, García-Alcocer G, Berumen LC, Rosas-Arellano A, Mileli R, et al. (2008) Expression of GABA_A subunits during rat cerebellum development. *Neurosci Lett* 432: 1-6.
7. García-Alcocer G, Padilla K, Rodríguez A, Mileli R, Berumen LC (2012) Distribution of the purinergic receptors P2X(4) and P2X(6) during rat gut development. *Neurosci Lett* 509: 92-95.
8. García-Alcocer G, Sarabia-Altamirano G, Martínez-Torres A, Mileli R (2005) Developmental expression of 5-HT_{5A} receptor mRNA in the rat brain. *Neurosci Lett* 379: 101-105.
9. García-Alcocer G, Segura LC, García Peña M, Martínez-Torres A, Mileli R (2006) Ontogenetic distribution of 5-HT_{2C}, 5-HT_{5A}, and 5-HT₇ receptors in the rat hippocampus. *Gene Expr* 13: 53-57.
10. Mora Loyola E, Moreno Layseca P, Escobar Cabrera J, Berumen LC, Rodríguez Torres A, et al. (2010) Distribution of serotonin receptors 5-HT_{5A} during neurulation. *J Pathol* 220(S1): S8.