

Idiopathic Pulmonary Hypertension Induced Thrombocytopenia - A Case Report

Sananta Kumar Dash*, Venugopal Kulkarni, Rajendra Kumar Sahoo, Gopi Macherla and Ravikiran M

Department of Anesthesia and Critical Care Medicine, Citizens Hospitals and American Oncology Institute, Hyderabad, India

Abstract

Idiopathic pulmonary hypertension (PH) is a diagnosis of exclusion for any patient presenting with pulmonary hypertension (PH). Patient with PH may present with thrombocytopenia along with other signs of PH. We report a case of PH who presented with thrombocytopenia and breathlessness. The possible causes of thrombocytopenia were evaluated and excluded prior to concluding that her thrombocytopenia is due to PH. Various pathogenic mechanisms have been described for thrombocytopenia associated with PH.

Keywords: Idiopathic pulmonary hypertension; Thrombocytopenia; Primary respiratory alkalosis

Introduction

Idiopathic PH is a rare but potentially fatal disease. Multiple pathogenic mechanisms have been described for thrombocytopenia in patients with PH.

Case Report

A 46 year female who was diagnosed to have carcinoma of the breast presented with progressive breathlessness to a state of New York Heart Association functional class IV over 15 days. She was hypotensive (90/64 mmHg) and hypoxic (SpO₂ 88% on room air) on presentation. Initial resuscitation with 500 ml of crystalloid and high flow oxygen with a non-rebreathing mask was done. On clinical evaluation the positive findings included a raised jugular venous pressure (JVP), a loud pulmonary component of second heart sound (P2), pan systolic murmur in tricuspid area and clear lung fields on auscultation. There was no organomegaly and no clinical findings suggestive of deep venous thrombosis. Laboratory investigations revealed thrombocytopenia (platelet counts 38000/cc), raised D Dimer (1.5 µg/ml, normal <0.5 µg/ml), primary respiratory alkalosis in arterial blood gas analysis, prominent common pulmonary and right pulmonary artery on X-ray chest anterior-posterior view. The electrocardiogram (ECG) showed Q wave and inverted T wave in lead III. Immediate computerized tomographic pulmonary angiography (CTPA) ruled out any major thromboembolic event, mediastinal compression or mass. A prominent common pulmonary artery (diameter 2.97 cm) (Figure 1) and a dilated RV was marked on CT scan (Figure 2). A two dimension transthoracic echocardiography (2D Echo) showed a dilated right atrium (RA) and right ventricle (RV) and Doppler showed moderate to severe tricuspid regurgitation with a peak velocity of 3.8 m/s. A triple lumen catheter was inserted in the right internal jugular vein and the central venous pressure reading was 21 mmHg which estimated the right ventricular systolic pressure to be 81 mmHg. A working diagnosis of idiopathic pulmonary hypertension was made after exclusion of possible differential diagnoses and treatment with Sildenafil (50 mg orally twice daily) and Bosentan (62.5 mg orally twice daily) started. She received a transfusion of single donor platelet (as the platelet count dropped further to 20000/cc). She became progressively more dyspnoeic and hypotensive. Vasopressor (Noradrenaline) and inotrope (Dobutamine) were started and dose titrated to maintain a mean arterial blood pressure around 65 mmHg. She was stable for the next three hours after which she complained of worsening breathlessness, became irritable and disoriented. The heart rate suddenly dropped to 40/min with no palpable pulse. Cardiopulmonary resuscitation was started, her trachea was intubated and ventilated with 100% oxygen. The endotracheal suction showed frank hemorrhage and the rhythm was now asystole.




Figure 1: Prominent pulmonary artery in CTPA.

There was no response to all the resuscitation efforts and we were not able to revive her.

Discussion

Our patient could be categorized as Idiopathic pulmonary hypertension (previously categorized as Primary Pulmonary Hypertension (PPH) [1,2]. Idiopathic PH is a rare disease, with varying incidence from 1 to 5.9 cases per million population. Twelve percent of the diagnosed cases are inherited as autosomal dominant [3].

In a cohort study including 576 patients of PH of various etiology, 307 (53%) died during follow-up (the median follow-up time was 3.9 years). The reported 1, 3 and 5 yr survival rates for the PH were 86, 69 and 61%, respectively [4].


Differential diagnosis considered were medical conditions

*Corresponding author: Sananta Kumar Dash, MD, FNB (Critical Care Medicine), EDIC, FCCP, Department of Anesthesia and Critical Care Medicine, Citizens Hospitals and American Oncology Institute, Hyderabad, India, Tel: +919963322755; E-mail: drsananta@yahoo.co.in

Received May 12, 2014; Accepted February 29, 2016; Published March 03, 2016

Citation: Dash SK, Kulkarni V, Sahoo RK, Macherla G, Ravikiran M (2016) Idiopathic Pulmonary Hypertension Induced Thrombocytopenia - A Case Report. J Pulm Respir Med 6: 322. doi:10.4172/2161-105X.1000322

Copyright: © 2016 Dash SK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



associated with PH (including portal hypertension, [5] and connective tissue disease (most commonly systemic sclerosis) [6]. Other possible etiologies considered include human immunodeficiency virus infection [7], chronic hemolytic anemias [8], and congenital heart disease (Eisenmenger syndrome). The diagnosis of idiopathic PH was made by careful exclusion of the above possibilities by detailed clinical history, family history, clinical examinations and specific laboratory parameters.

Pathology behind the worse outcome in PH includes the following. RV is a thin walled chamber structured to handle a low pressure pulmonary circulation [9,10]. Unlike the normal physiology, in PH the perfusion to the RV is gradually restricted only to diastole with the progression of severity of PH resulting in progressive ischemia and failure of RV [11]. Though there may be a disparity of 10-20 mmHg of PASP measurement invasively as compared to 2D Echo, the latter is the preferred initial modality to diagnose, quantify and follow-up PH [12].


Expert panel recommends invasive measurement of pulmonary artery systolic pressure (PASP) as the gold standard but feasibility in each case needs evaluation [13].

In our case PH was diagnosed and quantitated by 2D Echo.

The recommendations for the use of Pulmonary vasodilators (Sildenafil and Bosentan) are restricted to WHO group I pulmonary hypertension (i.e., primary PH or idiopathic PH) probably because, this is the most commonly studied group in clinical trials [14].

The recommended therapies for PH fall into 3 classes of pulmonary vasodilators: phosphodiesterase-5 inhibitors, endothelin receptor antagonists, and prostacyclin analogues. The appropriate choice of the agent should be made judiciously as an erroneous administration of vasodilators in pulmonary venous hypertension may lead to worsening of venous pressure and possible pulmonary edema [14].

Mortality in PH is directly related to RV dysfunction. Hence optimal management of RV dysfunction in all subsets of PH is important. PH with chronic right heart failure is managed by diuresis, sodium and water restriction. Aggressive fluid resuscitation is avoided. Early administration of vasopressors and inotropes are indicated for correction of hypotension and shock [14]. Treatment measures are targeted to augment the supply and decrease the demand for myocardial oxygen. These include oxygen supplementation, maintenance of adequate hematocrit, avoidance of acidosis, reduction in the RV afterload using specific pulmonary vasodilators like inhaled nitric oxide and reducing RV volume load by diuresis and/or ultrafiltration in refractory cases [14]. Our patient received a fluid resuscitation of



500 ml followed by early administration of Noradrenaline (to increase the coronary perfusion pressure), Dobutamine (to support the compromised RV), pulmonary vasodilators-Sildenafil and Bosentan (to reduce RV afterload) and oxygen supplementation. We were unable to use NO as it was not available with us and inhaled prostacyclin was not considered because of thrombocytopenia. We transfused one unit of single donor platelet though there was no evidence of bleeding clinically in view of decreasing trend of platelet count (38000 to 20000/cc).

The possible pathogenic mechanisms for thrombocytopenia in our patient are as described (Figure 3). Blood flow through the pulmonary vasculature in presence of PH can lead to activation of thrombin which binds to protease-activated receptor-1 (PAR1) on platelets. It leads to platelet degranulation and release of factors like adenosine diphosphate (ADP) leading to platelet aggregation [15]. Nitric oxide (NO) prevents premature activation of platelets. Due to defective endothelial nitric oxide synthases (eNOS), the production of NO is decreased in idiopathic PH. It causes a decreased threshold for platelet activation in patients with idiopathic PH [15].

Pathological “Plexiform” lesions in vasculature of lung have been described in patients of idiopathic PH. Fibrin deposition occurs in these plexiform lesions. Red blood cells (RBCs) and platelets experience injury to their cell membranes while passing through these lesions leading to increased cell damage [16-19]. Moreover, concurrent association of diseases such as portal hypertension leading to splenic sequestration, idiopathic thrombocytopenic purpura (ITP) and antiphospholipid antibody syndrome may present with thrombocytopenia along with idiopathic PH [20].

Conclusion

PPH presenting with thrombocytopenia though rare and reported anecdotally, they are known to have worse prognosis. Other causes of possible thrombocytopenia needs to be excluded prior to concluding idiopathic PH as the cause for it. The treatment modality for PH is still not clearly defined.

References

1. Simonneau G, Gatzoulis MA, Adatia I, Celermajer D, Denton C, et al. (2013) Updated Clinical Classification of Pulmonary Hypertension. J Am Coll Cardiol 62: 34-41.
2. Simonneau G, Robbins IM, Beghetti M, Channick RN, Delcroix M, et al. (2009) Updated clinical classification of pulmonary hypertension. J Am Coll Cardiol 54: S43-54.
3. Nichols WC, Koller DL, Slovis B, Foroud T, Terry VH, et al. (1997) Localization of the gene for familial primary pulmonary hypertension to chromosome 2q31-32. Nat Genet 15: 277-280.

4. Thenappan T, Shah SJ, Rich S, Tian L, Archer SL, et al. (2010) Survival in pulmonary arterial hypertension: a reappraisal of the NIH risk stratification equation. *Eur Respir J* 35: 1079-1087.
5. Li W, Dunmore BJ, Morrell NW (2010) Bone morphogenetic protein type II receptor mutations causing protein misfolding in heritable pulmonary arterial hypertension. *Proc Am Thorac Soc* 7: 395-398.
6. Denton CP, Black CM (2003) Pulmonary hypertension in systemic sclerosis. *Rheum Dis Clin North Am* 29: 335-349.
7. Almodovar S, Hsue PY, Morelli J, Huang L, Flores SC (2011) Pathogenesis of HIV-associated pulmonary hypertension: potential role of HIV-1 Nef. *Proc Am Thorac Soc* 8: 308-312.
8. Farmakis D, Aessopos A (2011) Pulmonary hypertension associated with hemoglobinopathies: prevalent but overlooked. *Circulation* 123: 1227-1232.
9. Haddad F, Hunt SA, Rosenthal DN, Murphy DJ (2008) Right ventricular function in cardiovascular disease, part I: Anatomy, physiology, aging, and functional assessment of the right ventricle. *Circulation* 117: 1436-1448.
10. Haddad F, Doyle R, Murphy DJ, Hunt SA (2008) Right ventricular function in cardiovascular disease, part II: pathophysiology, clinical importance, and management of right ventricular failure. *Circulation* 117: 1717-1731.
11. van Wolferen SA, Marcus JT, Westerhof N, Spreeuwenberg MD, Marques KM, et al. (2008) Right coronary artery flow impairment in patients with pulmonary hypertension. *Eur Heart J* 29: 120-127.
12. Rich JD, Shah SJ, Swamy RS, Kamp A, Rich S (2011) Inaccuracy of Doppler echocardiographic estimates of pulmonary artery pressures in patients with pulmonary hypertension: implications for clinical practice. *Chest* 139: 988-993.
13. Galiè N, Hoeper MM, Humbert M, Torbicki A, Vachiery JL, et al. (2009) Guidelines for the diagnosis and treatment of pulmonary hypertension: the Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS), endorsed by the International Society of Heart and Lung Transplantation (ISHLT). *Eur Heart J* 30: 2493-2537.
14. Shah SJ (2012) Pulmonary hypertension. *JAMA* 308: 1366-1374.
15. Aytekin M, Aulak KS, Haserodt S, Chakravarti R, Cody J, et al. (2012) Abnormal platelet aggregation in idiopathic pulmonary arterial hypertension: role of nitric oxide. *Am J Physiol Lung Cell Mol Physiol* 302: 512-520.
16. Jubelirer SJ (1991) Primary pulmonary hypertension. Its association with microangiopathic hemolytic anemia and thrombocytopenia. *Arch Intern Med* 151: 1221-1223.
17. Fischer EG, Marek JM, Morris A, Nashelsky MB (2000) Cholesterol granulomas of the lungs associated with microangiopathic hemolytic anemia and thrombocytopenia in pulmonary hypertension. *Arch Pathol Lab Med* 124: 1813-1815.
18. Stuard ID, Heusinkveld RS, Moss AJ (1972) Microangiopathic hemolytic anemia and thrombocytopenia in primary pulmonary hypertension. *N Engl J Med* 287: 869-870.
19. Bjornsson J, Edwards WD (1985) Primary pulmonary hypertension: a histopathologic study of 80 cases. *Mayo Clin Proc* 60: 16-25.
20. Horn EM, Barst RJ, Poon M (2000) Epoprostenol for treatment of pulmonary hypertension in patients with systemic lupus erythematosus. *Chest* 118: 1229-1230.

Citation: Dash SK, Kulkarni V, Sahoo RK, Macherla G, Ravikiran M (2016) Idiopathic Pulmonary Hypertension Induced Thrombocytopenia - A Case Report. *J Pulm Respir Med* 6: 322. doi:[10.4172/2161-105X.1000322](https://doi.org/10.4172/2161-105X.1000322)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>